ISSN 0021-9673

PUBLISHED
WEEKLY

CUMULATIVE AUTHOR AND SUBJECT INDEXES VOLUMES 211–220

Supplied free of charge to the 1981 subscribers of the Journal of Chromatography

JOURNAL OF

CHROMATOGRAPHY

INTERNATIONAL JOURNAL ON CHROMATOGRAPHY, ELECTROPHORESIS AND RELATED METHODS

EDITOR, Michael Lederer (Switzerland)

ASSOCIATE EDITOR, K. Macek (Prague)

EDITORIAL BOARD
W. A. Aue (Halifax)
V. G. Berezkin (Moscow)
V. Betina (Bratislava)
A. Bevenue (Honolulu, HI)
P. Boulanger (Lille)
A. A. Boulton (Saskatoon)
G. P. Cartoni (Rome)
G. Duyckaerts (Liège)
L. Fishbein (Jefferson, AR)
R. W. Frei (Amsterdam)
A. Frigerio (Milan)
C. W. Gehrke (Columbia, MO)
E. Gil-Av (Rehovot)
G. Guiochon (Palaiseau)
I. M. Hais (Hradec Králové)
J. K. Haken (Kensington)
E. Heftmann (Berkeley, CA)
S. Hjertén (Uppsala)
E. C. Horning (Houston, TX)
Cs. Horváth (New Haven, CT)
J. F. K. Huber (Vienna)
A. T. James (Sharnbrook)
J. Janák (Brno)
E. sz. Kováts (Lausanne)
K. A. Kraus (Oak Ridge, TN)
E. Lederer (Gif-sur-Yvette)
A. Liberti (Rome)
H. M. McNair (Blacksburg, VA)
Y. Marcus (Jerusalem)
G. B. Marini-Bettòlo (Rome)
Č. Michalec (Prague)
R. Neher (Basel)
G. Nickless (Bristol)
J. Novák (Brno)
N. A. Parris (Wilmington, DE)
P. G. Righetti (Milan)
O. Samuelson (Göteborg)
G.-M. Schwab (Munich)
G. Semenza (Zürich)
L. R. Snyder (Tarrytown, NY)
A. Zlatkis (Houston, TX)

EDITORS, BIBLIOGRAPHY SECTION
K. Macek (Prague), J. Janák (Brno), Z. Deyl (Prague)

COORD. EDITOR, DATA SECTION
J. Gasparič (Hradec Králové)

ELSEVIER SCIENTIFIC PUBLISHING COMPANY
AMSTERDAM

JUN 15 7 54 AM '82
UNIVERSITY OF HAWAII
LIBRARY

PERIODICAL
QD
271
.J6
Indexes
v.211-220
U.H. LIBRARY

JOURNAL OF CHROMATOGRAPHY

Scope. The *Journal of Chromatography* publishes papers on all aspects of chromatography, electrophoresis and related methods. Contributions consist mainly of research papers dealing with chromatographic theory, instrumental development and their applications. The section *Biomedical Applications,* which is under separate editorship, deals with the following aspects: developments in and applications of chromatographic and electrophoretic techniques related to clinical diagnosis (including the publication of normal values); screening and profiling procedures with special reference to metabolic disorders; results from basic medical research with direct consequences in clinical practice; combinations of chromatographic and electrophoretic methods with other physicochemical techniques such as mass spectrometry. In *Chromatographic Reviews,* reviews on all aspects of chromatography, electrophoresis and related methods are published.

Submission of Papers. Papers in English, French and German may be submitted, in three copies. Manuscripts should be submitted to: The Editor of *Journal of Chromatography,* P.O. Box 681, 1000 AR Amsterdam, The Netherlands, or to: The Editor of *Journal of Chromatography, Biomedical Applications,* P.O. Box 681, 1000 AR Amsterdam, The Netherlands. Review articles are invited or proposed by letter to the Editors and will appear in *Chromatographic Reviews* or *Biomedical Applications.* An outline of the proposed review should first be forwarded to the Editors for preliminary discussion prior to preparation. Submission of an article is understood to imply that the article is original and unpublished and is not being considered for publication elsewhere. For copyright regulations, see below.

Subscription Orders. Subscription orders should be sent to: Elsevier Scientific Publishing Company, P.O. Box 211, 1000 AE Amsterdam, The Netherlands. The *Journal of Chromatography* and the *Biomedical Applications* section can be subscribed to separately.

Publication. The *Journal of Chromatography* (incl. *Biomedical Applications, Chromatographic Reviews* and *Cumulative Author and Subject Indexes, Vols. 221–230, 231–240* and *241–250*) has 25 volumes in 1982. The subscription prices for 1982 are:

J. Chromatogr. (incl. *Chromatogr. Rev.* and *Cum. Indexes Vols. 221–230, 231–240 and 241–250*) + *Biomed. Appl.* (Vols. 227–251):
Dfl. 3625.00 plus Dfl. 500.00 (postage) (total *ca.* US$ 1650.00)

J. Chromatogr. (incl. *Chromatogr. Rev.* and *Cum. Indexes Vols. 231–240* and *241–250*) only (Vols. 234–251):
Dfl. 2826.00 plus Dfl. 360.00 (postage) (total *ca.* US$ 1274.50)

Biomed. Appl. (incl. *Cum. Indexes Vols. 221–230*) only (Vols 227–233):
Dfl. 1050.00 plus Dfl. 140.00 (postage) (total *ca.* US$ 476.00).

Journals are automatically sent by airmail to the U.S.A. and Canada at no extra costs, and to Japan, Australia and New Zealand, with a small additional postal charge. Back volumes of the *Journal of Chromatography* (Vols. 1 through 226) are available at Dfl. 173.00 (plus postage). Claims for issues not received should be made within three months of publication of the issue. If not, they cannot be honoured free of charge. For customers in the U.S.A. and Canada, wishing additional bibliographic information on this and other Elsevier journals, please contact Elsevier Science Publishing Company Inc., Journal Information Center, 52 Vanderbilt Avenue, New York, NY 10017. Tel: (212) 867-9040.

Abstracts/Contents Lists published in Analytical Abstracts, Biochemical Abstracts, Biological Abstracts, Chemical Abstracts, Chemical Titles, Current Contents/Physical, Chemical & Earth Sciences, Current Contents/Life Sciences, Index Medicus, Mass Spectrometry Bulletin, and Science Citation Index.

See page 3 of cover for Publication Schedule, Information for Authors, and information on the News Section and Advertisements.

© ELSEVIER SCIENTIFIC PUBLISHING COMPANY — 1982

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without the prior written permission of the publisher, Elsevier Scientific Publishing Company, P.O. Box 330, 1000 AH Amsterdam, The Netherlands.

Submission of an article for publication implies the transfer of the copyright from the author(s) to the publisher and entails the authors' irrevocable and exclusive authorization of the publisher to collect any sums or considerations for copying or reproduction payable by third parties (as mentioned in article 17 paragraph 2 of the Dutch Copyright Act of 1912 and in the Royal Decree of June 20, 1974 (S. 351) pursuant to article 16 b of the Dutch Copyright Act of 1912) and/or to act in or out of Court in connection therewith.

Special regulations for readers in the U.S.A. This journal has been registered with the Copyright Clearance Center, Inc. Consent is given for copying of articles for personal or internal use, or for the personal use of specific clients. This consent is given on the condition that the copier pays through the Center the per-copy fee stated in the code on the first page of each article for copying beyond that permitted by Sections 107 or 108 of the U.S. Copyright Law. The appropriate fee should be forwarded with a copy of the first page of the article to the Copyright Clearance Center, Inc., 21 Congress Street, Salem, MA 01970, U.S.A. If no code appears in an article, the author has not given broad consent to copy and permission to copy must be obtained directly from the author. All articles published prior to 1980 may be copied for a per-copy fee of US$ 2.25, also payable through the Center. This consent does not extend to other kinds of copying, such as for general distribution, resale, advertising and promotion purposes, or for creating new collective works. Special written permission must be obtained from the publisher for such copying.

Special regulations for authors in the U.S.A. Upon acceptance of an article by the journal, the author(s) will be asked to transfer copyright of the article to the publisher. This transfer will ensure the widest possible dissemination of information under the U.S. Copyright Law.

Printed in The Netherlands

SIEMENS

New horizons in GC

Easy adaptation to new tasks

Take a glance at the new SiCHROMAT® GC: All chromatographic elements are assembled in one compact block.

Then look at it more closely: You will find that the fittings for the injector and detector modules are standardized and interchangeable. This enables you to assemble any combination to suit your specific requirements; and future modules will have the same interfaces.

Further advantages of the analysis units are:

- the high-grade steel, heat-cleaned pipes for carrier and auxiliary gases are extremely short – hence less contamination and high trace sensitivity.
- the new gasket system can be loosened as often as required without loss of seal. With it, you can use glass, quartz, plastic and metal pipes at will, and above all: pipes with straight ends.

The high-grade steel micro-diaphragms used in the SiCHROMAT are also a new development. They control the flow of carrier and auxiliary gases very reliably.

Whether you use packed columns, capillary columns or any combination of the two – the Siemens SiCHROMAT gives you the flexibility you need.

For a detailed description of the advantages of the new Siemens SiCHROMAT, ask for our 24-page brochure now.



with the modular system of the Siemens SiCHROMAT

E 699/8103-101

To: Siemens AG, Infoservice, Postfach 156, D-8510 Fuerth.
Please send me your 24-page SiCHROMAT brochure
1234

Order your desk copy of:

CUMULATIVE AUTHOR AND SUBJECT INDEXES OF THE JOURNAL OF CHROMATOGRAPHY

covering **Volumes 1-50.**
1972 282 pages.
Price: US $56.00/Dfl.115.00.

covering **Volumes 51-100.**
1975 354 pages.
Price: US $78.00/Dfl. 160.000.

covering **Volumes 101-110.**
1975 126 pages.
Price: US $26.75/Dfl. 55.00.

covering **Volumes 111-120.**
1976 128 pages.
Price: US $26.75/Dfl. 55.00.

covering **Volumes 121-130.**
1977 84 pages.
Price: US $19.50/Dfl. 40.00.

covering **Volumes 131-140.**
1978 88 pages.
Price: US $19.50/Dfl. 40.00.

covering **Volumes 141-160.**
1978 168 pages.
Price: US $34.25/Dfl. 70.00.

covering **Volumes 161-180.**
1980 179 pages.
Price: US $35.00/Dfl. 70.00.

One copy of the Cumulative Author and Subject Index is supplied free of charge to subscribers of the Journal of Chromatography. Additional copies can be purchased.

ELSEVIER

P.O. Box 211, Amsterdam.
The Netherlands.

The Dutch guilder price is definitive. US $ price are subject to exchange rate fluctuations.

7051A

For advertising information please contact our advertising representatives

U.S.A./CANADA

Leni Ellinger
3133 Connecticut Ave, NW
Suite 712
WASHINGTON, DC 20008
Tel.: (202) 232-3366

GREAT BRITAIN

T.G. Scott & Son Ltd.
Mr. M.L. White
30-32 Southampton Street
LONDON, WC2E 7HE
Tel.: (01) 379-7264

OR

General Advertising Department

Elsevier/ Excerpta Medica/ North-Holland
Ms W. van Cattenburch
P.O. Box 211
1000 AE AMSTERDAM
The Netherlands
Tel.: (020) 5803.714/715
Telex: 18582 ESP NL

631EVe

Reader service card

JOURNAL OF

CHROMATOGRAPHY

INTERNATIONAL JOURNAL ON CHROMATOGRAPHY, ELECTROPHORESIS AND RELATED METHODS

use this card for more information on advertisements and news items

I would like to receive, without any obligation, further information on

News items: _____

Advertisements: _____

I am especially interested in the fields as indicated below:

- ☐ 6040 Analytical Chemistry
- ☐ 6050 Chromatography
- ☐ 6150 Mass Spectrometry
- ☐ 6140 Spectroscopy
- ☐ 6240 Pharmaceutical Chemistry
- ☐ 6241 Medicinal Chemistry
- ☐ 6242 Clinical Chemistry
- ☐ 6260 Environmental Chemistry

Name: _____

Position: _____

Address: _____

_____

please use envelope; mail as printed matter

ELSEVIER SCIENCE PUBLISHERS

Advertising department

P.O. BOX 211,

1000 AE AMSTERDAM – THE NETHERLANDS

please use envelope; mail as printed matter

ELSEVIER SCIENCE PUBLISHERS

Advertising department

P.O. BOX 211,

1000 AE AMSTERDAM – THE NETHERLANDS

Reader service card

JOURNAL OF

CHROMATOGRAPHY

INTERNATIONAL JOURNAL ON CHROMATOGRAPHY, ELECTROPHORESIS AND RELATED METHODS

use this card for more information on advertisements and news items

I would like to receive, without any obligation, further information on

News items: _____

Advertisements: _____

I am especially interested in the fields as indicated below:

- ☐ 6040 Analytical Chemistry
- ☐ 6050 Chromatography
- ☐ 6150 Mass Spectrometry
- ☐ 6140 Spectroscopy
- ☐ 6240 Pharmaceutical Chemistry
- ☐ 6241 Medicinal Chemistry
- ☐ 6242 Clinical Chemistry
- ☐ 6260 Environmental Chemistry

Name: _____

Position: _____

Address: _____

_____

JOURNAL *of* CHROMATOGRAPHY

INTERNATIONAL JOURNAL ON CHROMATOGRAPHY, ELECTROPHORESIS AND RELATED METHODS

Cumulative Author and Subject Indexes Volumes 211–220 (1981)

ELSEVIER SCIENTIFIC PUBLISHING COMPANY
AMSTERDAM
1982

Electronic data processing:
Büro für Satztechnik W. Meyer KG
Hanau, G.F.R.

© ELSEVIER SCIENTIFIC PUBLISHING COMPANY — 1982

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without the prior written permission of the publisher, Elsevier Scientific Publishing Company, P.O. Box 330, 1000 AH Amsterdam, The Netherlands.

Submission of an article for publication implies the transfer of the copyright from the author(s) to the publisher and entails the authors' irrevocable and exclusive authorization of the publisher to collect any sums or considerations for copying or reproduction payable by third parties (as mentioned in article 17 paragraph 2 of the Dutch Copyright Act of 1912 and in the Royal Decree of June 20, 1974 (S. 351) pursuant to article 16 b of the Dutch Copyright Act of 1912) and/or to act in or out of Court in connection therewith.

Special regulations for readers in the U.S.A. This journal has been registered with the Copyright Clearance Center, Inc. Consent is given for copying of articles for personal or internal use, or for the personal use of specific clients. This consent is given on the condition that the copier pays through the Center the per-copy fee stated in the code on the first page of each article for copying beyond that permitted by Sections 107 or 108 of the U.S. Copyright Law. The appropriate fee should be forwarded with a copy of the first page of the article to the Copyright Clearance Center, Inc., 21 Congress Street, Salem, MA 01970, U.S.A. If no code appears in an article, the author has not given broad consent to copy and permission to copy must be obtained directly from the author. All articles published prior to 1980 may be copied for a per-copy fee of US$ 2.25, also payable through the Center. This consent does not extend to other kinds of copying, such as for general distribution, resale, advertising and promotion purposes, or for creating new collective works. Special written permission must be obtained from the publisher for such copying.

Special regulations for authors in the U.S.A. Upon acceptance of an article by the journal, the author(s) will be asked to transfer copyright of the article to the publisher. This transfer will ensure the widest possible dissemination of information under the U.S. Copyright Law.

Printed in The Netherlands

AUTHOR INDEX

VOLUMES 211—220

The volume number is placed in parentheses before the page number

A

Aaberg, A., see Barth, T. (214) 83

Abbott, S., Achener, P., Simpson, R. and Klink, F.
Effect of radial thermal gradients in elevated temperature HPLC (218) 123

Abidi, S.L.
Speciation of aryloxyethoxyethyl benzyl dimethyl ammonium salts by glass capillary GC and HPLC (213) 463

Achener, P., see Abbott, S. (218) 123

Agnello, S.A., see Poole, C.F. (217) 39

Ahmed, M.S., Fong, H.H.S., Soejarto, D.D., Dobberstein, R.H., Waller, D.P. and Moreno-Azorero, R.
HPLC separation and quantitation of maytansinoids in *Maytenus ilicifolia* (213) 340

Akhtar, M.H., see Foster, T.S. (216) 303

Akiyoshi, D.E., see MacDonald, E.M.S. (214) 101

Albertsson, P.-Å. and Andersson, B.
Separation of membrane components by partition in detergent-containing polymer phase systems. Isolation of the light harvesting chlorophyll *a/b* protein (215) 131

Alder, J.F., Drew, P.K.P. and Fielden, P.R.
Improved high-frequency permittivity detector for the chromatography of alcohols and other polar species (212) 167

Alexander, G. and Hazai, I.
Chromatographic fingerprinting of coal extracts (217) 19

Alexander, S.P., see Gill, R. (218) 639

Alfredson, T.V.
HPLC column switching techniques for rapid hydrocarbon group-type separations (218) 715

Ali, S.N.
PC separation of phosphate esters, tricarboxylic cycle acids and amino acids in extracts from malaria parasites (214) 111

Alkharasani, M.A. and McCoy, B.J.
GLSC: effects of adsorption and partitioning (213) 203

Alonso-Fernández, J.R., Bóveda, M.D., Parrado, C., Peña, J. and Fraga, J.M.
Continuous TLC of sugars of clinical interest in samples of urine impregnated on paper (217) 357

Anderson, A. and Donald, A.S.R.
Improved method for the isolation of 2'-fucosyllactose from human milk (211) 170

Anderson, A., Goodall, E., Bliss, G.W. and Woodhouse, R.N.
Analysis of the anti-coccidial drug, halofuginone, in chicken tissue and chicken feed using HPLC (212) 347

Andersson, B., see Albertsson, P.-Å. (215) 131

Andersson, L.
Non-linear calibration by cubic splines in gel permeation chromatography (216) 23

Andersson, L.
Experimental evaluation of the precision in time-based high-performance gel permeation chromatography measurements (216) 35

Andersson, L.-O.
Purification and studies of components of the haemostatic system by affinity chromatography (215) 153

Ando, T., see Terabe, S. (212) 295

Arakawa, Y., Wada, O., Yu, T.H. and Iwai, H.
Simultaneous determination of trialkyltin homologues in biological materials (216) 209

Ardrey, R.E. and Moffat, A.C.
GLC retention indices of 1318 substances of toxicological interest on SE-30 or OV-1 stationary phase (Review) (220) 195

Arkel, J. van, see Van Arkel, J. (214) 349

Arrendale, R.F., see Severson, R.F. (211) 111

Arx, E. von, see Von Arx, E. (211) 262

Ashton, G.C., see Buteau, C. (212) 23

Atkinson, T., see Lowe, C.R. (215) 303

Atkinson, T., see Small, D.A.P. (216) 175

Atwood, J.G. and Golay, M.J.E.
Dispersion of peaks by short straight open tubes in LC systems (218) 97

Atwood, J.G., see DiCesare, J.L. (217) 369

Axelson, M., see Sahlberg, B.-L. (217) 453

Ayers, J.A., see Hearn, M.T.W. (218) 509

Ayers, J.S., see Bethell, G.S. (219) 353, (219) 361

Azanza, J.-L., see Raymond, J. (212) 199
Azogu, C.I.
TLC analysis of aromatic hydrocarbons in crude oil and in petroleum products as their η^6-arene-η^5-cyclopentadienyliron hexafluorophosphates (219) 349
Azzar, G., Degiuli, A., Benedetto, J.P. and Got, R.
Fractionation of microsomes of hepatocytes by isotachophoresis (in French) (213) 177

B

Baaske, D.M., Smith, M.D., Karnatz, N. and Carter, J.E.
HPLC determination of sodium nitroprusside (212) 339
Bächmann, K., see Vierkorn-Rudolph, B. (217) 311
Baerns, M., see Müller, B. (214) 217
Bailey, E., Peal, J.A. and Verschoyle, R.D.
The determination of O,S,S-trimethylphosphorodithioate in the plasma and various tissues of rats using high-resolution GC with nitrogen-phosphorus detection (219) 285
Baker, S.R., see McKay, S.W. (214) 249, (219) 325
Balogh, S., see Vigh, G. (214) 335
Barbaro, A.M., see Guerra, M.C. (216) 93
Barceló, D., Galcerán, M.T. and Eek, L.
Scanning electron microscopy as an aid in GC. Study of porous polymers coated with stationary phases (217) 109
Barkan, S., Weber, J.D. and Smith, E.
Determination of cross-contamination of the diastereomers ephedrine and pseudoephedrine by HPLC, TLC and carbon-13 nuclear magnetic resonance spectroscopy (219) 81
Barnes, J.A., see Kennedy, J.F. (212) 179
Barre, J., see Zini, R. (216) 191
Barrnett, R., see Janero, D.R. (216) 417
Barth, T., Tjessem, K. and Aaberg, A.
Fractionation of polar organic constituents in environmental samples using the lipophilic dextran gels Sephadex LH-20 and Sephasorb HP Ultrafine. Application to a weathered Ekofisk crude oil (214) 83
Bascoul, J., see Korahani, V. (211) 392
Becher, G.
Glass capillary columns in the GC separation of aromatic amines. II. Application to samples from workplace atmospheres using nitrogen-selective detection (211) 103
Becker, B. and Wojnowski, W.
Contributions to the chemistry of silicon-sulphur compounds. XXVI. Relationships between chromatographic behaviour and structure of triorganosilanes, triorganosilanethiols, hexaorganodisilthianes and triphenylthioalkylsilanes (219) 302
Beecher, G.R., see Vanderslice, J.T. (216) 338
Behl, H.M., see Proksch, P. (213) 345
Belenkii, B.G., see Janča, J. (211) 39
Bellew, L., see Katusz, R.M. (213) 331
Belov, Y.P., see Davankov, V.A. (218) 547
Benecke, I., see König, W.A. (217) 71
Benedetto, J.P., see Azzar, G. (213) 177
Beránek, J., see Ryba, M. (211) 337
Berezin, B.B., see Yamskov, I.A. (217) 539
Berg, J.H.M. van den, see Van den Berg, J.H.M. (218) 327
Bertani-Dziedzic, L., see Krstulović, A.M. (217) 523
Berthou, F. and Friocourt, M.P.
GC separation of diastereomeric isoprenoids as molecular markers of oil pollution (219) 393
Bethell, G.S., Ayers, J.S., Hearn, M.T.W. and Hancock, W.S.
Investigation of the activation of cross-linked agarose with carbonylating reagents and the preparation of matrices for affinity chromatography purifications (219) 353
Bethell, G.S., Ayers, J.S., Hearn, M.T.W. and Hancock, W.S.
Investigation of the activation of various insoluble polysaccharides with 1,1'-carbonyldiimidazole and of the properties of the activated matrices (219) 361
Bethell, G.S., see Hearn, M.T.W. (218) 509
Betts, T.J., Finucane, G.J. and Tweedie, H.A.
Practical system for polarity rating of packed GLC columns (213) 317
Beyer, T.A., see Sadler, J.E. (215) 181
Biagi, G.L., see Guerra, M.C. (216) 93
Bianchini, J.P. and Gaydou, E.M.
Role of water in qualitative and quantitative determination of polymethoxylated flavones by straight-phase HPLC: application to orange peel oils (211) 61
Bier, M., Mosher, R.A. and Palusinski, O.A.
Computer simulation and experimental validation of isoelectric focusing in ampholine-free systems (211) 313
Biftu, T.
Essential oil composition of *Aframomum korarima* (211) 280
Bilder, B.M., see Henion, J.D. (213) 475
Billiet, H.A.H., Schoenmakers, P.J. and De Galan, L.
Retention and selectivity characteristics of a non-polar perfluorinated stationary phase for LC (218) 443
Billiet, H.A.H., see Laurent, C. (218) 83
Billiet, H.A.H., see Schoenmakers, P.J. (218) 261
Bird, C.R. and Smith, T.A.
Separation of amines, guanidines and

hydroxycinnamic acid amides by ion-exchange chromatography (214) 263
Björkqvist, B.
Separation and determination of phenyl isocyanate-derivatized carbohydrates and sugar alcohols by HPLC with UV detection (218) 65
Bjørnson, I., see Jellum, E. (217) 231
Bjørseth, A., see Ugland, K. (213) 83
Bláha, K., see Turková, J. (215) 165
Bläker, F., see Weiland, E. (214) 156
Bleumer, J.P.A., see Tijssen, R. (218) 137
Bliss, G.W., see Anderson, A. (212) 347
Blomstrand, R., see Sisfontes, L. (216) 115
Boček, P., Gebauer, P. and Deml, M.
Migration behaviour of the hydrogen ion and its role in isotachophoresis of cations (217) 209
Boček, P., Gebauer, P. and Deml, M.
Concept of the effective mobility of the hydrogen ion and its use in cationic isotachophoresis (219) 21
Bochkov, A.S., see Davankov, V.A. (218) 547
Bock, U.E.G. and Waser, P.G.
GC determination of some biogenic amines as their pentafluorobenzoyl derivatives in the picogram range and its applicability to biological materials (213) 413
Bodor, G., see Samay, G. (218) 473
Boer, F. de, see De Boer, F. (214) 257
Boettner, E.A., see Sweetman, J.A. (212) 115
Boll, E., see Funk, W. (217) 349
Borea, P.A., see Guerra, M.C. (216) 93
Borvák, J., see Štrop, P. (214) 317
Borys, A.
HPLC determination of 4-nitroso- and 4-nitrophenols in the presence of phenol and alkylphenols (216) 361
Bos, P., see Lawrence, J.F. (212) 245
Boschetti, E., see Girot, P. (213) 389
Bossi, L., see Semerdjian-Rouquier, L. (218) 663
Boursier, B., see Fremy, J.-M. (219) 156
Bóveda, M.D., see Alonso-Fernández, J.R. (217) 357
Bradbury, A.G.W., Halliday, D.J. and Medcalf, D.G.
Separation of monosaccharides as trimethylsilylated alditols on fused-silica capillary columns (213) 146
Brassinne, C., Vanhaelen-Fastré, R. and Vanhaelen, M.
HPLC determination of Nocodazole in liposomes and L 1210 cells (213) 170
Braun, W.H., see Kastl, P.E. (213) 156
Bree, F., see Zini, R. (216) 191
Breitenbach, M., see Halperin, G. (215) 211
Brenner, K.S. and Huber, W.
The role of chromatography in BASF. Chromatographic techniques employed in BASF for investigatory studies and for problem solving (Review) (220) 95
Březina, M., Vodička, L., Tříska, J. and Kříž, J.
Separation of diamantane-3-oneoxime stereoisomers by preparative HPLC (219) 179
Brinkman, U.A.T., Welling, P.L.M., De Vries, G., Scholten, A.H.M.T. and Frei, R.W.
LC of demoxepam and phenothiazines using a post-column photochemical reactor and fluorescence detection (217) 463
Brinkman, U.A.T., see Scholten, A.H.M.T. (218) 3
Brobst, K.M., see Scobell, H.D. (212) 51
Broquaire, M., see Guinebault, P. (217) 509
Brown, J.F., see Vanderslice, J.T. (216) 338
Brown, N.D., Strickler, M.P., Sleeman, H.K. and Doctor, B.P.
Determination of N-methyl pyridinium-2-aldoxime chloride and its hydrolytic by-products by ion-pair HPLC (212) 361
Brown, P.R., see Cucco, J.A. (213) 253
Brownlee, S.G., see Vanderslice, J.T. (216) 338
Brugman, W.J.T., Heemstra, S. and Kraak, J.C.
HPLC of organic acids on bare silica (218) 285
Buchholz, K., see Kasche, V. (216) 169
Buchmann, M. and Kesselring, U.W.
Separation of the decomposition products of dimetindene on Sephadex G-10 and LH-20 (in French) (213) 515
Budahegyi, M.V., see Czerwiec, Z. (214) 47
Buhl, F., Hachuła, U. and Chwistek, M.
Application of the combined redox-complexation reaction to the detection of antipyrine and its derivatives in TLC (219) 189
Bull, R.V.A., Lim, C.K. and Gray, C.H.
HPLC of bile pigments: separation and characterization of the urobilinoids (218) 647
Burkman, A.M., see Wu, C.C. (216) 239
Bury, A.F.
Analysis of protein and peptide mixtures. Evaluation of three sodium dodecyl sulphate-polyacrylamide gel electrophoresis buffer systems (213) 491
Bushway, R.J.
HPLC determination of carbaryl and 1-naphthol at residue levels in various water sources by direct injection and trace enrichment (211) 135
Bushway, R.J., see Wilson, A.M. (214) 140
Buswell, J.A., Eriksson, K.-E. and Pettersson, B.
Purification and partial characterization of vanillate hydroxylase (decarboxylating) from *Sporotrichum pulverulentum* (215) 99
Buteau, C., Duitschaever, C.L. and Ashton, G.C.
Stability of the *o*-phthalaldehyde-histamine complex (212) 23
Butler, H., see Poole, C.F. (217) 39
Butte, W., Fooken, C., Klussmann, R. and Schuller, D.
Evaluation of lipophilic properties for a

series of phenols, using reversed-phase HPLC and HPTLC (214) 59
Buytenhuys, F.A.
Ion chromatography of inorganic and organic ionic species using refractive index detection (218) 57

C

Cadman, E., see Dreyer, R. (219) 273
Cairati, L., Cannizzaro, M., Comini, A., Gatti, G. and Vita-Finzi, P.
HPLC separation of the intermediate products in the synthesis of trimethylolpropane (216) 402
Calam, D.H. and Davidson, J.
Analysis of glycoprotein hormones and other medically important proteins by high-performance gel filtration chromatography (218) 581
Calam, D.H., see Pask-Hughes, R.A. (214) 307
Caldwell, K.D., Karaiskakis, G. and Giddings, J.C.
Characterization of T4D virus by sedimentation field-flow fractionation (215) 323
Calixto, F.S. and Raso, A.G.
Relationship between GC retention indices of homologous compounds and the number of carbon atoms and temperature (216) 326
Cannizzaro, M., see Cairati, L. (216) 402
Cantelli Forti, G., see Guerra, M.C. (216) 93
Capp, M.W., see Nice, E.C. (218) 569
Carlier, M., see Pauwels, J.-F. (211) 247
Carlson, R.M.K., see Lee, G.J.-L. (212) 108
Carter, J.E., see Baaske, D.M. (212) 339
Castello, G. and D'Amato, G.
Characterization of Chromosorb porous polymer bead columns by GC retention values of light hydrocarbons and carbon dioxide (212) 261
Certa, U. and Wessel, D.
Simple gel apparatus for horizontal polyacrylamide and agarose gel electrophoresis (211) 257
Chadwick, V.S., see Murphy, R. (211) 160
Chai, M.S.Y., see Vidal, C.J. (219) 71
Chang, G.-D., see Lo, T.-B. (215) 229
Chaplin, J.F., see Severson, R.F. (211) 111
Chapoteau, E., see Malin, M.J. (219) 117
Chase, T.N., see Knight, M. (212) 356
Chauhan, J. and Darbre, A.
Determination of acetyl and formyl groups by GLC (211) 347
Chaytor, J.P. and Saxby, M.J.
Determination of patulin and penicillic acid in unroasted cocoa beans (214) 135
Chen, E.C.M. and Wentworth, W.E.
Correlation and prediction of electron-capture response from molecular parameters (217) 151
Chen, T., see Fenselau, C. (218) 21
Chibata, I., Tosa, T. and Sato, T.
Application of carrageenan beads for chromatographic purification of proteins (215) 93
Chretien, J.R., see Haken, J.K. (217) 125
Churáček, J., see Jandera, P. (214) 35
Chwistek, M., see Buhl, F. (219) 189
Claes, P., see Decoster, W. (211) 223
Cochrane, W.P., Singh, J., Miles, W. and Wakeford, B.
Determination of chlorinated dibenzo-*p*-dioxin contaminants in 2,4-D products by GC-MS techniques (217) 289
Cole, S.C.J., Flanagan, R.J., Johnston, A. and Holt, D.W.
Rapid HPLC method for the measurement of verapamil and norverapamil in blood plasma or serum (218) 621
Colenutt, B.A., see Tyman, J.H.P. (213) 287
Colin, H., see Jandera, P. (214) 35
Collins, D.J., see Sahlberg, B.-L. (217) 453
Comini, A., see Cairati, L. (216) 402
Conte, A. and Cossi, G.
GC determination of free toluene diisocyanate in flexible urethane foams (213) 162
Cooke, M., see Roberts, D.J. (213) 73
Cooke, N., see Nice, E.C. (218) 569
Corfield, A.P., see Veh, R.W. (212) 313
Corran, P.H., see Pask-Hughes, R.A. (214) 307
Cosgrove, B.A. and Walkley, J.
Solubilities of gases in H_2O and ^2H_2O (216) 161
Cossi, G., see Conte, A. (213) 162
Cotter, R., see Fenselau, C. (218) 21
Cottrell, R.C., see Walters, D.G. (219) 152
Coulet, P.R. and Gautheron, D.C.
Enzymes immobilized on collagen membranes: a tool for fundamental research and enzyme engineering (215) 65
Čoupek, J., see Turková, J. (215) 165
Coustard, J.M. and Sudraud, G.
Separation of ascorbic and isoascorbic acids by ion-pair reversed-phase chromatography (in French) (219) 338
Coy, K., see Taylor, M.W. (219) 133
Crastes de Paulet, A., see Korahani, V. (211) 392
Cristalli, M., see Grassini-Strazza, G. (214) 209
Crosby, D.G., see Draper, W.M. (216) 413
Crowther, J.B., Jones, R. and Hartwick, R.A.
HPLC of the oligonucleotides (217) 479
Cser, F., see Samay, G. (218) 473
Cucco, J.A. and Brown, P.R.
Confirming the presence of N-nitrosamines in ambient air and cigarette smoke by converting to and photochemically altering their corresponding N-nitroamines (213) 253
Culbreth, P.H. and Sampson, E.J.
LC measurement of cortisol in methylene

chloride extracts of aqueous solutions (212) 221
Czerwiec, Z., Budahegyi, M.V. and Takács, J.M.
GC data for chlorobenzenes, chloroanilines and N-sulphinylanilines. Relationship between molecular structure and retention based on retention index increments calculated by computer (214) 47

D

Dam, H.C. van, see Van Dam, H.C. (218) 83
D'Amato, G., see Castello, G. (212) 261
Dammann, V., see Funk, W. (217) 349
Darbre, A., see Chauhan, J. (211) 347
Davankov, V.A., Bochkov, A.S. and Belov, Y.P.
Ligand-exchange chromatography of racemates. XV. Resolution of α-amino acids on reversed-phase silica gels coated with N-decyl-L-histidine (218) 547
Davankov, V.A., see Kurganov, A.A. (218) 559
Davankov, V.A., see Yamskov, I.A. (217) 539
Davidson, J., see Calam, D.H. (218) 581
Davis, P.J., Schoenl, M. and LaMantia, R.S.
Interaction of heme proteins and thyroid hormone. II. Localization of the site on thyroid hormone that binds to hemoglobin (213) 148
Davis, P.J., see Hughes, J.T. (219) 321
Davis, P.J., see Kerr, K.M. (219) 317
De Boer, F., see Huizing, H.J. (214) 257
De Decker, G., see Verzele, M. (214) 95
De Galan, L., see Billiet, H.A.H. (218) 443
De Galan, L., see Laurent, C. (218) 83
De Galan, L., see Schoenmakers, P.J. (218) 261
De Pauw, C., see Verzele, M. (214) 95
De Taeye, L., see Verzele, M. (214) 95
De Vries, G., see Brinkman, U.A.T. (217) 463
Dean, P.D.G., see Jessup, W. (219) 419
DeArman Lukacs, K., see Jorgenson, J.W. (218) 209
Dębowski, J., see Kutner, W. (218) 45
Decker, G. de, see De Decker, G. (214) 95
Decoster, W., Claes, P. and Vanderhaeghe, H.
Determination of the relative amounts of the B and C components of neomycin by ion-exclusion chromatography using refractometric detection (211) 223
Deelder, R.S. and Van den Berg, J.H.M.
Study on the retention of amines in reversed-phase ion-pair chromatography on bonded phases (218) 327
Degen, P.H., see Schneider, W. (217) 263
Degiuli, A., see Azzar, G. (213) 177
Deigin, V.I., see Ulyashin, V.V. (215) 263
Delitala, L. Falchi, see Falchi Delitala, L. (219) 332
Dellacherie, E., see Sacco, D. (211) 79
DellaMonica, E.S., see Egoville, M.J. (212) 121
Deming, S.N. and Kong, R.C.
Correlation of Hammet substituent constants with reversed-phase "ion-pair" LC adsorption parameters of charged organic compounds (217) 421
Deml, M., see Boček, P. (217) 209, (219) 21
Denkert, M., Hackzell, L., Schill, G. and Sjögren, E.
Reversed-phase ion-pair chromatography with UV-absorbing ions in the mobile phase (218) 31
Desideri, P.G., see Lepri, L. (211) 29
Desiderio, D.M., Yamada, S., Tanzer, F.S., Horton, J. and Trimble, J.
HPLC and field desorption MS measurement of picomole amounts of endogenous neuropeptides in biologic tissue (217) 437
Dewaele, C., see Verzele, M. (217) 399
DiCesare, J.L., Dong, M.W. and Atwood, J.G.
Very-high-speed LC. II. Some instrumental factors influencing performance (217) 369
Dilli, S., see Pillai, D.N. (220) 253
DiMari, S.J., Robinson, J.P. and Hash, J.H.
Use of a ternary gradient for the separation of phenylthiohydantoin-amino acids, including the methyl esters of aspartic and glutamic acids by HPLC (213) 91
Dimov, N., see Papazova, D. (216) 321
DiRico, D.E., see Krstulović, A.M. (217) 523
Dobashi, Y., see Oka, K. (217) 387
Dobberstein, R.H., see Ahmed, M.S. (213) 340
Doctor, B.P., see Brown, N.D. (212) 361
Doi, T., see Ôi, N. (211) 274, (213) 137
Donald, A.S.R., see Anderson, A. (211) 170
Dong, M.W., see DiCesare, J.L. (217) 369
Dose, K., see Nawroth, T. (214) 126
Dose, K., see Schmidt, R. (212) 370
Dostavalov, I.N., see Yamskov, I.A. (217) 539
Draffehn, J., Ponsold, K. and Schönecker, B.
R_M values of steroids as an expression of their hydrophobic character. III. Interpretation of the slopes of the R_M *versus* acetone concentration curves (216) 69
Draffehn, J., Schönecker, B. and Ponsold, K.
R_M values of steroids as an expression of their hydrophobic character. II. Effect of neighbouring groups on the hydrophobicity of substituents (216) 63
Draper, W.M. and Crosby, D.G.
Sensitive, enzyme-catalyzed, chromogenic reagent for hydrogen peroxide and other peroxygen compounds on TLC plates (216) 413
Drew, P.K.P., see Alder, J.F. (212) 167
Dreyer, R. and Cadman, E.
Use of periodate and methylamine for the quantitation of intracellular 5-fluoro-2'-deoxyuridine-5'-monophosphate by HPLC (219) 273
Dufek, P., Pacáková, V. and Žyvný, K.
GC behaviour of mono- and

dihydroxybiphenyls on various silicone phases (211) 150
Duitschaever, C.L., see Buteau, C. (212) 23
Durand, E.F., see Tulliez, J.E. (219) 411
Duval, M., see Lamarre, C. (213) 481
Dyck, J. van, see Van Dyck, J. (214) 95
Dziedzic, S.W., see Krstulović, A.M. (217) 523

E

Eck, W.S., Gray, R.R., Gegoux, T.A., Schoo, G.M. and Strickler, M.P.
Determination of aprophen in biological samples by normal-phase HPLC (219) 171
Eckert, T. and Knie, U.
Influence of reflecting supports of HPTLC silica gel 60 stationary phase on quantitative *in situ* measurements of thin-layer chromatograms (in German) (213) 453
Eek, L., see Barceló, D. (217) 109
Egly, J.-M.
Aromatic interaction chromatography. Acriflavin gel for separation of nucleotides, oligonucleotides and nucleic acids (215) 243
Egoville, M.J. and DellaMonica, E.S.
GC determination of water in acetone (212) 121
Ehhalt, D.H., see Rudolph, J. (217) 301
Einarsson, M., Forsberg, B., Larm, O., Riquelme, M.E. and Scholander, E.
Coupling of proteins and other amines to Sepharose by bromine oxidation and reductive amination (215) 45
Ek, K., see Righetti, P.G. (220) 115
Elliger, C.A. and Rabin, L.B.
Separation of plant polyphenolics by chromatography on a boronate resin (216) 261
Ellwood, P.A., see Goldberg, P.A. (212) 93
Elmi-Akhounie, E., see Vidal, C.J. (219) 71
Elverdam, I., Larsen, P. and Lund, E.
Isolation and characterization of three new polymyxins in polymyxins B and E by HPLC (218) 653
Engelhardt, H. and Müller, H.
Chromatographic characterization of silica surfaces (218) 395
Engewald, W., see Priboth, I. (219) 113
Eriksson, K.-E., see Buswell, J.A. (215) 99
Eszéniova, A., see Madajová, V. (216) 313
Ettre, L.S.
The nomenclature of chromatography. II. Liquid chromatography (Review) (220) 29
Ettre, L.S.
The nomenclature of chromatography. III. General rules for future revisions (Review) (220) 65
Eyem, J.
Properties of a capillary column injector with a controlled rate of injection (217) 99

F

Fagan, E.A., see Murphy, R. (211) 160
Falchi Delitala, L., see Solinas, V. (219) 332
Falkowski, P.G. and Sucher, J.
Rapid, quantitative separation of chlorophylls and their degradation products by HPLC (213) 349
Fanali, S. and Ossicini, L.
High-performance paper electrophoresis. II. Comparison of separations obtained by high-voltage paper electrophoresis and high-performance paper electrophoresis (212) 374
Farkaš, P., see Soják, L. (219) 225
Fauchère, J.-L., see Pliška, V. (216) 79
Faupel, M. and Von Arx, E.
An efficient chemically bonded reversed-phase thin-layer plate permitting the use of water-rich mobile phases (211) 262
Fellous, R., Lafaye de Micheaux, D., Lizzani-Cuvelier, L. and Luft, R.
Behaviour of stationary phases. I. Linear relationships between retention data of benzene derivatives (in French) (213) 223
Fenselau, C., Cotter, R., Hansen, G., Chen, T. and Heller, D.
Middle molecule mass spectrometry (a review) (218) 21
Fiala, E.S. and Kulakis, C.
Separation of hydrazine, monomethylhydrazine, 1,1-dimethylhydrazine and 1,2-dimethylhydrazine by HPLC with electrochemical detection (214) 229
Fielden, P.R., see Alder, J.F. (212) 167
Fini, A., see Guerra, M.C. (216) 93
Finucane, G.J., see Betts, T.J. (213) 317
Fischer, F.C., see Van der Sluis, W.G. (214) 349
Fisher, M., see Murphy, R. (211) 160
Flanagan, R.J., see Cole, S.C.J. (218) 621
Flodin, P. and Lagerkvist, P.
Polymer gels (215) 7
Foery, R.F., see Katusz, R.M. (213) 331
Foffani, M.T., see Guerra, M.C. (216) 93
Fong, H.H.S., see Ahmed, M.S. (213) 340
Fooken, C., see Butte, W. (214) 59
Förch, A.D., see Hulshoff, A. (220) 275
Forsberg, B., see Einarsson, M. (215) 45
Forsythe, R., see Huang, J.-C. (214) 269
Forti, G. Cantelli, see Cantelli Forti, G. (216) 93
Foster, P.M.D., see Walters, D.G. (219) 152
Foster, T.S. and Akhtar, M.H.
HPLC of fenvalerate (216) 303

Fotso, M., see Raymond, J. (212) 199
Fox, S.W.
Copolyamino acid fractionation and protobiochemistry (215) 115
Fraga, J.M., see Alonso-Fernández, J.R. (217) 357
Franc, J. and Sechovec, J.
The role of hydrogen bonds in chromatography. I. Amino- and hydroxyanthraquinones (212) 139
Francis, G.W. and Veland, K.
Alkylthiolation for the determination of double-bond positions in linear alkenes (219) 379
Frank, J., see Kreuzig, F. (218) 615
Frei, R.W., see Brinkman, U.A.T. (217) 463
Frei, R.W., see Lawrence, J.F. (212) 245
Frei, R.W., see Scholten, A.H.M.T. (218) 3
Fremy, J.-M. and Boursier, B.
Rapid determination of aflatoxin M_1 in dairy products by reversed-phase HPLC (219) 156
Frenz, J.H., see Horváth, C. (218) 365
Fričová, V., see Přistoupil, T.I. (213) 183, (219) 128
Friedman, M. and Noma, A.T.
Histamine analysis on a single column amino acid analyzer (219) 343
Friocourt, M.P., see Berthou, F. (219) 393
Frydrychová, A., see Turková, J. (215) 165
Funazo, K., Tanaka, M. and Shono, T.
Methylation of inorganic anions for GC determination (211) 361
Funazo, K., see Wu, H.-L. (219) 312
Funk, W., Kerler, R., Boll, E. and Dammann, V.
HPTLC determination of fluorescence-labelled cortisol (217) 349
Füzes, L., see Samay, G. (218) 473

G

Gabel, D., see Genieser, H.-G. (215) 235
Galan, L. de, see De Galan, L. (218) 83, (218) 261, (218) 443
Galcerán, M.T., see Barceló, D. (217) 109
Galeffi, C., see Marini-Bettòlo, G.B. (213) 113
Galunsky, B., see Kasche, V. (216) 169
Ganapathy, V., Ramachandramurty, B. and Radhakrishnan, A.N.
Distinctive test with copper(II)-ninhydrin reagent for small α-peptides separated PC (213) 307
Gandhe, B.R., Panday, P., Sharma, R.K., Vaidyanathaswamy, R. and Shinde, S.K.
Studies on the GC behaviour of organophosphorus esters (219) 297
Ganjian, I., Loher, W. and Kubo, I.
Determination of prostaglandin E_2 in the cricket, *Teleogryllus commodus*, by reversed-phase HPLC (216) 380
Gatti, G., see Cairati, L. (216) 402
Gautheron, D.C., see Coulet, P.R. (215) 65
Gauthier, J., see Lamarre, C. (213) 481
Gaydou, E.M., see Bianchini, J.P. (211) 61
Gazdag, M. and Szepesi, G.
Separation of large polypeptides by HPLC (218) 603
Gazdag, M., see Szepesi, G. (218) 597
Gebauer, P., see Boček, P. (217) 209, (219) 21
Gegoux, T.A., see Eck, W.S. (219) 171
Genieser, H.-G., Gabel, D. and Jastorff, B.
Quantitative ether cleavage of ligands in hydrophobic agaroses: precise determination of the degree of substitution (215) 235
Germershausen, J., see Karkas, J.D. (214) 267
Gessa, C., see Solinas, V. (219) 332
Ghaemi, Y. and Wall, R.A.
Hydrophobic chromatography with dynamically coated stationary phases. V. "Duplex soap chromatography" (212) 271
Ghias-Ud-Din, M., Smith, A.E. and Phillips, D.V.
Separation of pinitol and some other cyclitols by HPLC (211) 295
Giddings, J.C., see Caldwell, K.D. (215) 323
Gies, D., see Veh, R.W. (212) 313
Giese, R.W., see Yu, T.J. (218) 519
Gilbert, J., see Shepherd, M.J. (218) 703
Gill, R., Alexander, S.P. and Moffat, A.C.
Group-contribution approach to the behaviour of 2-phenylethylamines in reversed-phase HPLC (218) 639
Gill, W.N.
Doubly peaked chromatograms (213) 355
Girot, P. and Boschetti, E.
Physico-chemical and chromatographic properties of new ion exchangers. I. CM-Trisacryl M (213) 389
Glad, M., see Lowe, C.R. (215) 303
Glajch, J.L. and Snyder, L.R.
Solvent strength of multicomponent mobile phases in LSC. Mixtures of three or more solvents (214) 21
Glajch, J.L., see Snyder, L.R. (214) 1, (218) 299
Glasel, J.A. and Venn, R.F.
Fluorescence and UV detection of opiates separated by reversed-phase HPLC (213) 337
Goegelman, R.T., see Gullo, V.P. (212) 234
Golay, M.J.E.
The height equivalent to a theoretical plate of retentionless rectangular tubes (216) 1
Golay, M.J.E., see Atwood, J.G. (218) 97
Goldberg, P.A., Walker, R.F., Ellwood, P.A. and Hardy, H.L.
Determination of trace atmospheric isocyanate concentrations by reversed-phase HPLC using 1-(2-pyridyl)piperazine reagent (212) 93

Goldstein, L.
Polymeric supports bearing isonitrile functional groups for covalent fixation of biologically active molecules (a review) (215) 31
Golovnya, R.V. and Samusenko, A.L.
Analysis of polar compounds on polyethylene glycol glass capillary columns. The influence of KF and Na_3PO_4 on the change in partial molar free energy (217) 183
Goodall, E., see Anderson, A. (212) 347
Goretti, G., see Laganà, A. (219) 263
Got, R., see Azzar, G. (213) 177
Gotto, A.M., see Hancock, W.S. (216) 285
Gowda, T.V., see Kini, R.M. (216) 395
Gozzini, L. and Montecucchi, P.C.
Reversed-phase HPLC of dermorphins, opiate-like peptides from amphibian skins (216) 355
Grassini-Strazza, G. and Cristalli, M.
Comparison of C_n bonded silica gel TLC plates: conditions for use and separations of some barbiturates (214) 209
Gray, C.H., see Bull, R.V.A. (218) 647
Gray, D.O., see Mitchell, S.D. (216) 137
Gray, R.R., see Eck, W.S. (219) 171
Grego, B., see Hearn, M.T.W. (218) 497
Greibrokk, T., see Svendsen, H. (212) 153, (213) 429
Greibrokk, T., see Ugland, K. (213) 83
Grob, G., see Grob, K. (211) 243, (213) 211, (219) 13
Grob, K. and Grob, G.
Capillary columns with immobilized stationary phases. II. Practical advantages and details of procedures (213) 211
Grob, K., Grob, G. and Grob, K. Jr.
Capillary columns with immobilized stationary phases. I. A new simple preparation procedure (211) 243
Grob, K., Grob, G. and Grob, K. Jr.
Testing capillary GC columns (219) 13
Grob, K. Jr.
Peak broadening or splitting caused by solvent flooding after splitless or cold on-column injection in capillary GC (213) 3
Grob, K. Jr. and Romann, A.
Sample transfer in splitless injections in capillary GC (214) 118
Grob, K. Jr., see Grob, K. (211) 243, (219) 13
Gruenauer, M.H., see Nelsen, T.R. (212) 366
Gschwender, H.G., see Haller, W. (211) 53
Gübitz, G., Wintersteiger, R. and Hartinger, A.
Fluorescence derivatization of tertiary amines with 2-naphthyl chloroformate (218) 51
Guerra, M.C., Barbaro, A.M., Cantelli Forti, G., Foffani, M.T., Biagi, G.L., Borea, P.A. and Fini, A.
R_M and log P values of 5-nitroimidazoles (216) 93
Guha, O.K. and Mishra, K.P.
Effect of pH on the GC behaviour of silica gel (219) 101
Guillochon, D., see Remy, M.H. (215) 87
Guinebault, P. and Broquaire, M.
Large-volume injection of samples dissolved in a non-eluting solvent. Application to the determination of antipyrine using normal-phase HPLC (217) 509
Guiochon, G., see Krupčík, J. (213) 189
Guiochon, G., see Sellier, F. (219) 197, (219) 213
Guise, G.B. and Smith, G.C.
Chromatography of polycarbamoyl sulphonates (anionic polyurethane derivatives) (214) 69
Gullo, V.P., Goegelman, R.T., Putter, I. and Lam, Y.-K.
HPLC analysis of derivatized hypocholesteremic agents from fermentation broths (212) 234
Gwynn, G.R., see Severson, R.F. (211) 111

H

Hachuła, U., see Buhl, F. (219) 189
Hackzell, L., see Denkert, M. (218) 31
Haefelfinger, P.
Limits of the internal standard technique in chromatography (218) 73
Haeffner-Gormley, L., Poludniak, N.H. and Wetlaufer, D.B.
Separation of the tryptic peptides from reduced, alkylated hen egg white lysozyme by HPLC (214) 185
Hafkenscheid, T.L. and Tomlinson, E.
Estimation of aqueous solubilities of organic non-electrolytes using LC retention data (218) 409
Hafkenscheid, T.L., see Riley, C.M. (218) 427
Hairsine, P.W., see Westwood, R. (219) 140
Haken, J.K., Chretien, J.R. and Lion, C.
GLC retention behaviour of hindered aliphatic esters (217) 125
Haken, J.K. and Obita, J.A.
Analysis of resinous or fatty polyamides using GC and GPC (213) 55
Haken, J.K. and Srisukh, D.
GC of homologous esters. XV. Molecular retention indices of aliphatic esters (219) 45
Haky, J.E. and Muschik, G.M.
Evaluation of liquid crystal smectic mesophases for GLC separations (214) 161
Haller, W., Gschwender, H.G. and Peters, K.-R.
Application of a general equation for controlled pore glass permeation chromatography to an aggregating, spherical virus (211) 53
Halliday, D.J., see Bradbury, A.G.W. (213) 146

Halperin, G., Breitenbach, M., Tauber-Finkelstein, M. and Shaltiel, S. Hydrophobic chromatography on homologous series of alkylagaroses. A comparison of charged and electrically neutral column materials (215) 211
Hancock, W.S., Pownall, H.J., Gotto, A.M. and Sparrow, J.T. Separation of apolipoproteins A-I and A-II by ion-paired reversed-phase HPLC (216) 285
Hancock, W.S., see Bethell, G.S. (219) 353, (219) 361
Hancock, W.S., see Hearn, M.T.W. (218) 509
Hanekamp, H.B., see Lawrence, J.F. (212) 245
Hansen, G., see Fenselau, C. (218) 21
Hansen, S.H. Ion-exchange and straight-phase partition HPLC of the opiates on silica (212) 229
Hansson, H. and Kågedal, L. Adsorption and desorption of proteins in metal chelate affinity chromatography. Purification of albumin (215) 333
Hara, A., see Hirai, H. (215) 195
Hara, S., see Oka, K. (217) 387
Hardy, H.L., see Goldberg, P.A. (212) 93
Harris, E.L., see Hearn, M.T.W. (218) 509
Harris, J.M., see Leach, R.A. (218) 15
Hartinger, A., see Gübitz, G. (218) 51
Hartwick, R.A., see Crowther, J.B. (217) 479
Harvey, D.J. GC and MS studies of ginger constituents. Identification of gingerdiones and new hexahydrocurcumin analogues (212) 75
Hash, J.H., see DiMari, S.J. (213) 91
Hashimoto, T., see Kato, Y. (211) 383
Haskó, F., see Vasileva, R. (216) 279
Hazai, I., see Alexander, G. (217) 19
Hearn, M.T.W. and Grego, B. HPLC of amino acids, peptides and proteins. XXXVI. Organic solvent modifier effects in the separation of unprotected peptides by reversed-phase LC (218) 497
Hearn, M.T.W., Harris, E.L., Bethell, G.S., Hancock, W.S. and Ayers, J.A. Application of 1,1'-carbonyldiimidazole-activated matrices for the purification of proteins. III. The use of 1,1'-carbonyl-diimidazole-activated agaroses in the biospecific affinity chromatographic isolation of serum antibodies (218) 509
Hearn, M.T.W. and Lyttle, D.J. Buffer-focusing chromatography using multicomponent electrolyte elution systems (218) 483
Hearn, M.T.W., see Bethell, G.S. (219) 353, (219) 361
Heemstra, S., see Brugman, W.J.T. (218) 285
Heftmann, E., see Lin, J.-T. (212) 239, (213) 507
Heimler, D., see Lepri, L. (211) 29
Heller, D., see Fenselau, C. (218) 21
Helmer, U., Olausson, Å. and Stensiö, K.-E. Determination of distearylcarbamoyl chloride by HPLC (211) 369
Helmich, O., Špaček, P. and Hradec, J. Separation of [^{3}H]cholesteryl esters by reversed-phase HPLC (213) 105
Henion, J.D., Nosanchuk, J.S. and Bilder, B.M. Capillary GC-MS determination of histamine in tuna fish causing scombroid poisoning (213) 475
Hermann, E.A., see Kastl, P.E. (213) 156
Hershey, H.V., see Taylor, M.W. (219) 133
Hibi, N., see Hirai, H. (215) 195
Hikino, H., Konno, C., Watanabe, H. and Ishikawa, O. Determination of aconitine alkaloids by HPLC (211) 123
Hill, H.H. Jr., see Osman, M.A. (213) 397
Hill, R.L., see Sadler, J.E. (215) 181
Hirai, H., Tsukada, Y., Hara, A., Hibi, N., Nishi, S., Wepsic, H.T., Koji, T. and Ishii, N. Purification of specific antibody to α-foetoprotein and its immunological effect on cancer cells (215) 195
Hirose, K., see Kawase, S. (213) 265
Hjertén, S. and Kunquan, Y. HPLC of macromolecules on agarose and its derivatives (215) 317
Hjertén, S., Kunquan, Y. and Ogunlesi, M. Immobilization of enzymes on columns of brushite (215) 25
Hlavay, J., see Vigh, G. (214) 335
Hlebarova, N., see Lekova, K. (212) 85
Hodgeman, D.K.C. Analysis of 2-hydroxybenzophenone and 2'-hydroxyphenylbenzotriazole UV stabilizers by HPLC (214) 237
Holík, M. and Matějková, B. Determination of hydroxyl groups and water content in silica by nuclear magnetic resonance spectroscopy (213) 33
Holt, D.W., see Cole, S.C.J. (218) 621
Holzer, G., see Poole, C.F. (217) 39
Hoogmartens, J., Roets, E. and Vanderhaeghe, H. Determination of 5-ethyl-5-(1-ethylpropyl)-barbituric acid in pentobarbital by HPLC (219) 431
Horáček, J., see Turková, J. (215) 165
Hořejší, V. and Tichá, M. Theory of affinity electrophoresis. Evaluation of the effects of protein multivalency, immobilized ligand heterogeneity and microdistribution and determination of effective concentration of immobilized ligand (216) 43
Horgan, R., see Martin, G.C. (219) 167
Horton, J., see Desiderio, D.M. (217) 437
Horváth, C., Nahum, A. and Frenz, J.H. High-performance displacement chromatography (218) 365
Horváth, C., see Kalász, H. (215) 295
Houston, R., see Ullman, A.H. (211) 398
Hradec, J., see Helmich, O. (213) 105
Huang, F.-L., see Lo, T.-B. (215) 229

Huang, J.-C., Forsythe, R. and Madey, R.
GSC of ethane on activated carbon at 25°C (214) 269
Huber, R., see Voelter, W. (217) 491
Huber, W., see Brenner, K.S. (220) 95
Hughes, J.T. and Davis, P.J.
HPLC determination of N,N-dimethylcolchiceinamide and its metabolites, N-methylcolchiceinamide and colchiceinamide, in microbial culture (219) 321
Huizing, H.J., De Boer, F. and Malingré, T.M.
Preparative ion-pair HPLC and GC of pyrrolizidine alkaloids from comfrey (214) 257
Hulshoff, A. and Förch, A.D.
Alkylation with alkyl halides as a derivatization method for the GC determination of acidic pharmaceuticals (Review) (220) 275
Huser, H., see Jürgens, D. (216) 295

I

Imai, J., see Okamoto, M. (219) 175
Inda, Y., see Ôi, N. (213) 137
Ishii, D. and Takeuchi, T.
Study of the performance of cation-exchange columns in open-tubular microcapillary LC (218) 189
Ishii, D., see Takeuchi, T. (213) 25, (216) 153, (218) 199
Ishii, N., see Hirai, H. (215) 195
Ishikawa, O., see Hikino, H. (211) 123
Ishizaki, H., see Matsumoto, H. (211) 403
Ito, R., see Takada, K. (216) 385
Ito, Y.
Efficient preparative counter-current chromatography with a coil planet centrifuge (214) 122
Ito, Y., see Knight, M. (212) 356
Itokawa, Y., see Kimura, M. (211) 290
Ivanov, V.T., see Ulyashin, V.V. (215) 263
Iverson, F., see Lawrence, J.F. (212) 245
Iwaeda, T., see Kato, Y. (211) 383
Iwai, H., see Arakawa, Y. (216) 209

J

Jacob, K., see Meyer, H.D. (217) 473
Jacob, K., see Vogt, W. (217) 91
Jakab, M., see Vasileva, R. (216) 279
Jaksch, Y., see Westerlund, D. (211) 181
Jamieson, W.B., see McKay, S.W. (214) 249
Janča, J., Pokorný, S., Vilenchik, L.Z. and Belenkii, B.G.
Concentration effects in gel permeation chromatography. VII. Viscosity phenomena and column geometry (211) 39
Jandera, P., Churáček, J. and Colin, H.
Gradient elution in LC. XIV. Theory of ternary gradients in reversed-phase LC (214) 35
Jandik, P. and Schmidbaur, H.
Applications of HPLC in organometallic chemistry: ligand exchange in complexes of ylides (in German) (213) 47
Jane, I., Scott, A., Sharpe, R.W.L. and White, P.C.
Quantitation of cocaine in a variety of matrices by HPLC (214) 243
Janero, D.R. and Barrnett, R.
Analytical separation of green-plant and animal neutral lipids by TLC (216) 417
Jastorff, B., see Genieser, H.-G. (215) 235
Jebsen, C., see Rudolph, J. (217) 301
Jellum, E., Bjørnson, I., Nesbakken, R., Johansson, E. and Wold, S.
Classification of human cancer cells by means of capillary GC and pattern recognition analysis (217) 231
Jennissen, H.P.
Immobilization of residues on agarose gels: effects on protein adsorption isotherms and chromatographic parameters (215) 73
Jensen, E.C., see Lochmüller, C.H. (216) 333
Jessup, W. and Dean, P.D.G.
Purification of 6-phosphogluconate dehydrogenase from *Acer pseudoplatanus* L. using immobilised Procion Red HE-3B (219) 419
Johansen, I. and Wendelboe, J.F.
Dimethylformamide and carbon disulphide desorption efficiencies for organic vapours on gas-sampling charcoal tube analyses with a GC backflush technique (217) 317
Johansson, E., see Jellum, E. (217) 231
Johansson, E., see Lindberg, W. (211) 201
Johansson, K., see Lindberg, W. (211) 201
John, M., see Schmidt, J. (213) 151
Johnson, A.W., see Severson, R.F. (211) 111
Johnston, A., see Cole, S.C.J. (218) 621
Johnston, M.A.
General method for the assay of oral contraceptives by HPLC (216) 269
Jojková, K., see Přistoupil, T.I. (213) 183
Jones, P. and Wellington, C.A.
Optimisation in chromatography. Theory and application to the separation of aromatic acids in reversed-phase LC (213) 357
Jones, R., see Crowther, J.B. (217) 479
Jönsson, J.Å. and Mathiasson, L.
System for injection of continuously variable amounts of sample vapours for

physico-chemical measurements by GC (219) 427
Jorgenson, J.W. and DeArman Lukacs, K. High-resolution separations based on electrophoresis and electroosmosis (218) 209
Jurand, J., see Knox, J.H. (218) 341, (218) 355
Jürgens, D. and Huser, H. Large-scale purification of staphylococcal lipase by hydrophobic interaction chromatography (216) 295

K

Kagan, M.Z., Kraevskaya, M.A., Vasilieva, V.S. and Zinkevich, E.P. Analytical and preparative separation of the *cis*- and *trans*-isomers of 4-(4'-*tert*.-butylcyclohexyl)-4-methylpentan-2-one by reversed-phase HPLC (219) 183
Kågedal, L., see Hansson, H. (215) 333
Kainuma, K., Nakakuki, T. and Ogawa, T. HPLC of maltosaccharides (212) 126
Kalász, H. and Horváth, C. Preparative-scale separation of polymyxins with an analytical HPLC system by using displacement chromatography (215) 295
Kalász, H., see Tyihák, E. (211) 45
Kalinichev, A.I., see Kolotinskaya, E.V. (212) 133
Kaliszan, R. Chromatography in studies of quantitative structure-activity relationships (220) 71
Kaniansky, D., see Madajová, V. (216) 313
Kara-Murza, S.G. Dynamics of dissemination in the case of affinity chromatography (Review) (220) 85
Karaiskakis, G., see Caldwell, K.D. (215) 323
Karaiskakis, G., see Vattis, D. (214) 171
Kardjieva, L., see Lekova, K. (212) 85
Karger, B.L., see Yu, T.J. (218) 519
Karkas, J.D., Germershausen, J. and Liou, R. Purification of potassium phosphate for HPLC (214) 267
Karlsson, K.-E. Two-phase derivatization of amitriptyline and structurally related tertiary amines for GC with electron-capture detection (219) 373
Karlstam, B. Application of polyacrylamide gradient gel electrophoresis to the study of multiple components of concanavalin A and related lectins (211) 233
Karmen, A., Pritchett, T. and Lam, S. Therapeutic drug assays with GLC and optical detection (217) 247
Karnatz, N., see Baaske, D.M. (212) 339
Kasahara, Y., see Nimura, N. (213) 327
Kasche, V., Buchholz, K. and Galunsky, B. Resolution in high-performance liquid affinity chromatography. Dependence on eluite diffusion into the stationary phase (216) 169
Kašička, V., see Štrop, P. (214) 317
Kastl, P.E., Hermann, E.A. and Braun, W.H. Quantitative determination of 1,2-dibromo-3-chloropropane in whole rat blood and drinking water by GC (213) 156
Katchalski-Katzir, E., see Solomon, B. (215) 121
Kato, Y., Komiya, K., Iwaeda, T., Sasaki, H. and Hashimoto, T. Packing of Toyopearl columns for gel filtration. IV. Gravitational packing and influence of slurry reservoir size (211) 383
Katsanos, N.A. and Tsiatsios, A. Kinetics of *n*-butene isomerization studied by stopped-flow GC (213) 15
Katsanos, N.A., see Vattis, D. (214) 171
Katusz, R.M., Bellew, L., Mangravite, J.A. and Foery, R.F. Recovery of HPLC-grade acetonitrile by spinning-band distillation (213) 331
Kawaguchi, Y., see Uobe, K. (214) 177
Kawase, S., Hirose, K. and Ukai, S. GLC determination of N-alkylpyridinium salts (213) 265
Kemula, W., Sybilska, D. and Lipkowski, J. Selectivity and efficiency of separation of isomers of organic acids by clathrate chromatography (218) 465
Kemula, W., see Kutner, W. (218) 45
Kennedy, J.F. and Barnes, J.A. Use of $F(ab')_2$ antibody fragments in the synthesis of immunoadsorbents for preparing monospecific antigen (212) 179
Kennedy, J.M. and McNamara, P.E. HPLC analysis of pilocarpine hydrochloride, isopilocarpine, pilocarpic acid and isopilocarpic acid in eye-drop preparations (212) 331
Kerler, R., see Funk, W. (217) 349
Kerr, K.M., Smith, R.V. and Davis, P.J. HPLC determination of pergolide and its metabolite, pergolide sulfoxide, in microbial extracts (219) 317
Kesselring, U.W., see Buchmann, M. (213) 515
Khedim, A., see Rudolph, J. (217) 301
Kimura, M. and Itokawa, Y. Separation and determination of thiamine-binding proteins in rats by HPLC (211) 290
Kinberger, B. Sensitive method for the HPLC determination of amphetamine in urinary extracts (213) 166
Kini, R.M. and Gowda, T.V. Single-step fractionation of *Vipera russelli* venom. A sensitive fluorimetric method to study the elution profile (216) 395
Kinoshita, T., see Nimura, N. (213) 327
Kirkland, J.J., see Snyder, L.R. (218) 299
Kirkland, J.J., see Yau, W.W. (218) 217
Kishida, Y., see Takaya, T. (215) 279

Kishimoto, H., see Okamoto, M. (212) 251
Kitahara, H., see Ôi, N. (211) 274, (213) 137
Kizaki, H. and Sakurada, T.
Separation of cAMP from adenine nucleotides and nucleosides by electrophoresis on cellulose acetate membranes (211) 409
Kling, F., see Tóth, G. (213) 511
Klink, F., see Abbott, S. (218) 123
Klussmann, R., see Butte, W. (214) 59
Knedel, M., see Meyer, H.D. (217) 473
Knie, U., see Eckert, T. (213) 453
Knight, M., Ito, Y. and Chase, T.N.
Preparative purification of the peptide des-enkephalin γ-endorphin. Comparison of HPLC and counter-current chromatography (212) 356
Knox, J.H. and Jurand, J.
Mechanism of zwitterion-pair chromatography. I. Nucleotides (218) 341
Knox, J.H. and Jurand, J.
Mechanism of zwitterion-pair chromatography. II. Ampicilline, lysergic acid, tryptophan and other solutes (218) 355
Kobayashi, T., see Takada, K. (216) 385
Koch, K., see Zeman, A. (216) 199
Koenigsberger, R.U., see McKay, S.W. (219) 325
Kohashi, K., see Zaitsu, K. (211) 129
Koji, T., see Hirai, H. (215) 195
Kolotinskaya, E.V., Kalinichev, A.I. and Semenovskaya, T.D.
Computer simulation of the kinetics of ion exchange accompanied by complex formation in ionites (212) 133
Komiya, K., see Kato, Y. (211) 383
Kong, R.C., see Deming, S.N. (217) 421
König, W.A., Benecke, I. and Sievers, S.
New results in the GC separation of enantiomers of hydroxy acids and carbohydrates (217) 71
Konno, C., see Hikino, H. (211) 123
Korahani, V., Bascoul, J. and Crastes de Paulet, A.
Capillary column GLC analysis of cholesterol derivatives. Application to the autoxidation products of cholesterol (211) 392
Korhonen, I.O.O.
GLC analyses. II. Glass capillary GC of methyl monochloro esters of aliphatic C_2-C_{18} *n*-carboxylic acids (211) 267
Korhonen, I.O.O.
GLC analyses of chlorination products of propionyl chloride (213) 63
Korhonen, I.O.O.
GLC analyses. III. Glass capillary GC of chloromethyl monochloro esters of aliphatic C_3-C_{12} *n*-carboxylic acids (219) 306
Korte, W.D.
Alternate procedure for the preparation and separation of benzyl derivatives of organic acids (214) 131
Koshimura, T., see Uobe, K. (214) 177
Kotinopoulos, M., see Vattis, D. (214) 171
Kouřilová, D., see Krejčí, M. (219) 61
Kraak, J.C., see Brugman, W.J.T. (218) 285
Kraevskaya, M.A., see Kagan, M.Z. (219) 183
Královičová, E., see Soják, L. (219) 225
Kraml, J., see Kramlová, M. (211) 308
Kraml, J., see Přistoupil, T.I. (213) 183, (219) 128, (219) 436
Kramlová, M., Přistoupil, T.I. and Kraml, J.
Electrofocusing of stroma-free haemoglobin and its derivatives in agarose isoelectric focusing gels (211) 308
Kramlová, M., see Přistoupil, T.I. (213) 183, (219) 128, (219) 436
Kraus, G., see Soják, L. (219) 225
Kraus, L., see Nasner, A. (216) 389
Krejčí, M., Kouřilová, D. and Vespalec, R.
Electrokinetic detection at different points in a narrow-bore glass column in LC (219) 61
Krejčí, M., Tesařík, K., Rusek, M. and Pajurek, J.
Flow characteristics and technology of capillary columns with inner diameters less than 15 μm in LC (218) 167
Kreuzig, F. and Frank, J.
Rapid automated determination of D-penicillamine in plasma and urine by ion-exchange HPLC with electrochemical detection using a gold electrode (218) 615
Kreveld, M.E. van, see Van Kreveld, M.E. (218) 137
Kříž, J., Vodička, L., Punčochářová, J. and Kuraš, M.
HPLC of alkylbenzenes on silica (219) 53
Kříž, J., see Březina, M. (219) 179
Krstulović, A.M., Dziedzic, S.W., Bertani-Dziedzic, L. and DiRico, D.E.
Plasma catecholamines in hypertension and pheochromocytoma determined using ion-pair reversed-phase chromatography with amperometric detection. Investigation of the separation mechanism and clinical methodology (217) 523
Krupčík, J., Guiochon, G. and Schmitter, J.M.
Measurement of retention data on open tubular columns coupled in series (213) 189
Kubo, I., see Ganjian, I. (216) 380
Kučera, J.
Affinity chromatography of chymotrypsin (E.C. 3.4.21.1) on the potato trypsin inhibitor bound to bead cellulose by the benzoquinone method (213) 352
Kucera, P. and Manius, G.
High-resolution reversed-phase LC utilizing microbore column concatenation (216) 9
Kucera, P. and Manius, G.
Recycling LC using microbore columns (219) 1
Kühn, H., see Priboth, I. (219) 113
Kulakis, C., see Fiala, E.S. (214) 229
Kumanotani, J., see Oshima, R. (213) 142
Kumanotani, J., see Yamauchi, Y. (214) 343
Kunquan, Y., see Hjertén, S. (215) 25, (215) 317
Kura, G.
Chromatographic study of the acidic

hydrolysis of cyclic octametaphosphate (211) 87
Kura, G., Nakashima, T. and Oshima, F. Study of the acidic hydrolysis of cyclic trimetaphosphate by LC (219) 385
Kuraš, M., see Kříž, J. (219) 53
Kurganov, A.A. and Davankov, V.A. Ligand-exchange chromatography of racemates. XVI. Microbore column chromatography of amino acid racemates using N,N,N',N'-tetramethyl-(*R*)-propanediamine-1,2-copper(II) complexes as chiral additives to the eluent (218) 559
Kuroda, T., see Tanaka, M. (219) 108
Kuroki, M., see Nakazawa, T. (211) 388
Kushinsky, S., see Lee, G.J.-L. (212) 108
Kuster, B.F.M., see Verhaar, L.A.T. (220) 313
Kutner, W., Dębowski, J. and Kemula, W. Extra-column effects in polarographic *versus* UV detection in HPLC (218) 45
Kutney, J.P., Sindelar, R.D. and Stuart, K.L. Rapid TLC assay of tripdiolide using fluorimetric detection (214) 152
Kuwata, K., Uebori, M. and Yamazaki, Y. Rapid method for packing microparticulate columns packed with a chemically bonded stationary phase for HPLC (211) 378

L

Låås, T., see Olsson, I. (215) 373
Labadie, R.P., see Van der Sluis, W.G. (214) 349
Lafaye de Micheaux, D., see Fellous, R. (213) 223
Lafont, P. and Siriwardana, M.G. Assay method for aflatoxin in milk (219) 162
Laganà, A., Goretti, G., Petronio, B.M. and Rotatori, M. Concentration and isolation of organic acids on graphitized carbon black (219) 263
Lagerkvist, P., see Flodin, P. (215) 7
Lam, S., see Karmen, A. (217) 247
Lam, Y.-K., see Gullo, V.P. (212) 234
LaMantia, R.S., see Davis, P.J. (219) 148
Lamarre, C., Duval, M. and Gauthier, J. Determination of 2,6-di-(*tert.* -butyl)-4-methylphenol in new and aged transformer oils using HPLC (in French) (213) 481
Landen, W.O. Jr. Resolution of fat-soluble vitamins in HPLC with methanol-containing mobile phases (211) 155
Larm, O., see Einarsson, M. (215) 45
Larsen, P., see Elverdam, I. (218) 653
Larsson, P.-O., see Lowe, C.R. (215) 303
Laurent, C., Billiet, H.A.H., Van Dam, H.C. and De Galan, L. Computer-controlled single-pump solvent programmer for HPLC (218) 83
Lawrence, J.F. Use of 214-nm and 229-nm discrete-line sources for the UV absorbance detection of some pesticides separated by HPLC (211) 144
Lawrence, J.F., Iverson, F., Hanekamp, H.B., Bos, P. and Frei, R.W. LC with UV absorbance and polarographic detection of ethylenethiourea and related sulfur compounds. Application to rat urine analysis (212) 245
Leach, R.A. and Harris, J.M. Thermal lens calorimetry. Application to chromatographic detection (218) 15
Lee, G.J.-L., Carlson, R.M.K. and Kushinsky, S. Use of organic modifiers in the mobile phase for the reversed-phase HPLC separation of steroidal hormones (212) 108
Lee, G.J.-L., Liu, D.-W., Pav, J.W. and Tieckelmann, H. Separation of reduced disaccharides derived from glycosaminoglycans by HPLC (212) 65
Lee, T.H., see Morris, S.C. (219) 403
Legentil, J.-P., see Szymanowicz, A. (211) 175
Lekova, K., Kardjieva, L., Hlebarova, N. and Natan, V. GLC of *s*-triazines on a surface-bonded support in short columns (212) 85
Lepri, L., Desideri, P.G. and Heimler, D. TLC of closely related polypeptides on silanized silica gel (211) 29
Levine, R.A., see Taylor, M.W. (219) 133
Li, C.H., see Yamashiro, D. (215) 255
Liebes, L.F. Continuous-flow scanning of selected HPLC peak components by microprocessor control. Application to analysis of extracts from human lymphocytes (219) 255
Liebich, H.M., Pickert, A. and Wöll, J. Analysis of the oxocarboxylic acid fraction in serum and urine as O-methyloximes by thermionic specific detection (217) 255
Lim, C.K., see Bull, R.V.A. (218) 647
Lin, J.-T. and Heftmann, E. HPLC of naturally occurring estrogens (212) 239
Lin, J.-T. and Heftmann, E. Adsorption and reversed-phase partition HPLC of gibberellins (213) 507
Lindberg, W., Johansson, E. and Johansson, K. Application of statistical optimization methods to the separation of morphine, codeine, noscapine and papaverine in reversed-phase ion-pair chromatography (211) 201
Lindström, K., see Renberg, L. (214) 327
Lion, C., see Haken, J.K. (217) 125
Liou, R., see Karkas, J.D. (214) 267
Lipkowski, J., see Kemula, W. (218) 465

Lipsky, S.R. and McMurray, W.J.
Role of surface groups in affecting the chromatographic performance of certain types of fused-silica glass capillary columns. I (217) 3
Lis, H. and Sharon, N.
Affinity chromatography for the purification of lectins (a review) (215) 361
Liu, D.-W., see Lee, G.J.-L. (212) 65
Lizzani-Cuvelier, L., see Fellous, R. (213) 223
Lloyd, R.J.
Nitrogen-selective detector for HPLC (216) 127
Lo, T.-B., Huang, F.-L. and Chang, G.-D.
Separation of subunits of pike eel gonadotropin by hydrophobic interaction chromatography (215) 229
Lochmüller, C.H. and Jensen, E.C.
Non-ionic mobile phase dopants. I. Chiral charge-transfer acceptors and helicene resolution (216) 333
Lockhart, W.L., see Murray, D.A.J. (212) 305
Loher, W., see Ganjian, I. (216) 380
Lotan, N., see Solomon, B. (215) 121
Lövkvist, P., see Mathiasson, L. (217) 177
Lowe, C.R., Glad, M., Larsson, P.-O., Ohlson, S., Small, D.A.P., Atkinson, T. and Mosbach, K.
High-performance liquid affinity chromatography of proteins on Cibacron Blue F3G-A bonded silica (215) 303
Lowe, C.R., see Small, D.A.P. (216) 175
Löwentap, M.S.H., see Ševčík, J. (217) 139
Lu Peichang, see Wang Chinghai (217) 191
Luft, R., see Fellous, R. (213) 223
Lund, E., see Elverdam, I. (218) 653
Lundanes, E., see Ugland, K. (213) 83
Lycourghiotis, A., see Vattis, D. (214) 171
Lyttle, D.J., see Hearn, M.T.W. (218) 483

M

Mabuchi, H. and Nakahashi, H.
Systematic separation of medium-sized biologically active peptides by HPLC (213) 275
McCoy, B.J., see Alkharasani, M.A. (213) 203
MacDonald, E.M.S., Akiyoshi, D.E. and Morris, R.O.
Combined HPLC radioimmunoassay for cytokinins (214) 101
McDuffie, K.L., see Severson, R.F. (211) 111
McGuffin, V.L. and Novotný, M.
Micro-column HPLC and flame-based detection principles (218) 179
Machin, A.F., see Mundy, D.E. (216) 229
McKay, S.W., Mallen, D.N.B., Shrubsall, P.R., Smith, J.M., Baker, S.R., Jamieson, W.B., Ross, W.J., Morgan, S.E. and Rackham, D.M.
Semi-preparative HPLC and spectroscopic characterisation of eight geometric isomers of leukotriene A methyl ester (214) 249
McKay, S.W., Mallen, D.N.B., Shrubsall, P.R., Smith, J.M., Baker, S.R. and Koenigsberger, R.U.
Separation of the N-trifluoroacetyl dimethyl esters of leukotriene D and E isomers by semi-preparative HPLC (219) 325
McMurray, W.J., see Lipsky, S.R. (217) 3
McMurrough, I.
HPLC of flavonoids in barley and hops (218) 683
McNamara, P.E., see Kennedy, J.M. (212) 331
Madajová, V., Kaniansky, D., Raděj, Z. and Eszéniova, A.
Isotachophoretic analysis of organic acids produced by fermentation from *n*-alkanes (216) 313
Madey, R., see Huang, J.-C. (214) 269
Maire, C.E., see Vanderslice, J.T. (216) 338
Majer, J., see Van Quy, T. (211) 177
Malgras, A., see Szymanowicz, A. (211) 175
Malin, M.J. and Chapoteau, E.
Determination of carbon chain distribution in alkyl sulfates by *in situ* hydrolysis-GC (219) 117
Malingré, T.M., see Huizing, H.J. (214) 257
Mallen, D.N.B., see McKay, S.W. (214) 249, (219) 325
Malmström, B.M., Nyman, P.-O. and Strid, L.
Improved separation of basic peptides in anion-exchange chromatography (215) 109
Mancheva, I.N., Nikolov, R.N. and Pfletschinger, J.
HPLC of diphenylindenonylthiohydantoin derivatives of amino acids (213) 99
Manecke, G. and Polakowski, D.
Some carriers for the immobilization of enzymes based on derivatized poly(vinyl alcohol) and on copolymers of methacrylates with different spacer lengths (215) 13
Mangravite, J.A., see Katusz, R.M. (213) 331
Manius, G., see Kucera, P. (216) 9, (219) 1
Marini-Bettòlo, G.B., Nicoletti, M., Patamia, M., Galeffi, C. and Messana, I.
Plant screening by chemical and chromatographic procedures under field conditions (213) 113
Martin, G.C., Horgan, R. and Scott, I.M.
HPLC analysis of permethylated cytokinins (219) 167
Matějková, B., see Holík, M. (213) 33
Mathiasson, L. and Lövkvist, P.
Comparison of column packings for trace analysis of free amines by GLC (217) 177
Mathiasson, L., see Jönsson, J.Å. (219) 427
Matsumoto, H., Takata, R.H. and Ishizaki, H.
Determination of the carcinogen methylazoxymethyl-β-D-glucosiduronic acid in rat bile and urine (211) 403

Matsuoka, K., see Takeuchi, T. (216) 153
Matsutera, E., Nobuhara, Y. and Nakanishi, Y.
Separation of nadolol diastereoisomers by reversed-phase HPLC (216) 374
Matsuzaki, K., see Ogamo, A. (213) 439
Medcalf, D.G., see Bradbury, A.G.W. (213) 146
Meek, J.L. and Rossetti, Z.L.
Factors affecting retention and resolution of peptides in HPLC (211) 15
Messana, I., see Marini-Bettòlo, G.B. (213) 113
Meyer, H.D., Jacob, K., Vogt, W. and Knedel, M.
Analysis of free stool porphyrins by HPLC (217) 473
Michalski, J.-C., see Veh, R.W. (212) 313
Micheaux, D. Lafaye de, see Lafaye de Micheaux, D. (213) 223
Miheeva, L.M., see Zaslavsky, B.Y. (212) 13, (216) 103
Miles, W., see Cochrane, W.P. (217) 289
Mincsovics, E., see Tyihák, E. (211) 45
Miron, T. and Wilchek, M.
Polyacrylhydrazido-agarose: preparation via periodate oxidation and use for enzyme immobilization and affinity chromatography (215) 55
Mishra, K.P., see Guha, O.K. (219) 101
Mitchell, S.D. and Gray, D.O.
Advantages of a new ion-exchange resin for the LC of biogenic amines: a methacrylic acid polymer cross-linked with butanedioldiacrylate (216) 137
Mitra, G.D.
Conversion of linear retention indices into logarithmic retention indices (211) 239
Mizobuchi, Y., see Tanaka, M. (219) 108
Mizunuma, H., see Nakagawa, T. (211) 1
Moffat, A.C., see Ardrey, R.E. (220) 195
Moffat, A.C., see Gill, R. (218) 639
Molnár-Perl, I. and Szakács-Pintér, M.
GLC separation and determination of the components of maltitol syrups (216) 219
Montecucchi, P.C., see Gozzini, L. (216) 355
Morávek, L., see Štrop, P. (214) 317
Moree-Testa, P. and Saint-Jalm, Y.
Determination of α-dicarbonyl compounds in cigarette smoke (217) 197
Morel, D. and Serpinet, J.
Influence of the LC mobile phase on the phase transitions of alkyl-bonded silicas studied by GC (214) 202
Moreno-Azorero, R., see Ahmed, M.S. (213) 340
Morgan, E.D., see Wilson, I.D. (212) 211
Morgan, S.E., see McKay, S.W. (214) 249
Mori, K.
Automated measurement of catecholamines in urine, plasma and tissue homogenates by HPLC with fluorometric detection (218) 631
Morris, R.O., see MacDonald, E.M.S. (214) 101
Morris, S.C. and Lee, T.H.
Analysis of potato glycoalkaloids with radially compressed HPLC cartridges and ethanolamine in the mobile phase (219) 403
Mosbach, K., see Lowe, C.R. (215) 303
Mosher, R.A., see Bier, M. (211) 313
Mühlstädt, M., see Priboth, I. (219) 113
Müller, B. and Baerns, M.
Direct GC determination of the products of catalytic air oxidation of n-butene-1 to maleic anhydride in the gaseous reaction mixture (214) 217
Müller, H., see Engelhardt, H. (218) 395
Mundy, D.E. and Machin, A.F.
Determination of pentachlorophenol and related compounds in animal materials by HPLC and GC (216) 229
Murakami, T., see Yamauchi, Y. (214) 343
Murphy, P.A.
Separation of genistin, daidzin and their aglucones, and coumesterol by gradient HPLC (211) 166
Murphy, R., Selden, A.C., Fisher, M., Fagan, E.A. and Chadwick, V.S.
HPLC analysis of polyethylene glycols (211) 160
Murray, D.A.J. and Lockhart, W.L.
Microextraction and GC analysis of selected petroleum hydrocarbons in water and fish tissue (212) 305
Muschik, G.M., see Haky, J.E. (214) 161
Myasoedov, N.F., see Yamskov, I.A. (217) 539

N

Nagasawa, K., see Ogamo, A. (213) 439
Nagy, J., see Tyihák, E. (211) 45
Nahum, A., see Horváth, C. (218) 365
Nakagawa, G., see Tsuda, T. (214) 283
Nakagawa, T., Mizunuma, H., Shibukawa, A. and Uno, T.
LC with crown ether-containing mobile phases. I. Capacity factors of amino compounds on a hydrophobic stationary phase (211) 1
Nakahashi, H., see Mabuchi, H. (213) 275
Nakakuki, T., see Kainuma, K. (212) 126
Nakanishi, Y., see Matsutera, E. (216) 374
Nakashima, T., see Kura, G. (219) 385
Nakatsu, K.
Fraction collector controller for collecting LC peaks (212) 105
Nakazawa, H., Sakaguchi, U. and Yoneda, H.
Chromatographic study of optical resolution. VII. Directional ion-association model for the optical resolution of *cis*-$[Co(O)_2(N)_4]^+$ type of complexes by the antimony d-tartrate ion (213) 323
Nakazawa, T., Kuroki, M. and Tsunashima, Y.
The chemistry of carbazoles. IX. Substituent effect in the GLC of methylcarbazoles (211) 388

Nasner, A. and Kraus, L.
Separation of some lecithin components by HPLC (in German) (216) 389
Natan, V., see Lekova, K. (212) 85
Navarro, X., see Segura, R. (217) 329
Nawrocki, J. and Szczepaniak, W.
Coordination polymers as adsorbents and stationary phases in GC. VII. Alkaline earth metal di-*n*-hexylphosphinates: specific interactions (212) 29
Nawroth, T. and Dose, K.
Application of a three-dimensional drawing procedure to the evaluation of series of protein samples after analysis by gel electrophoresis and other methods (214) 126
Nelsen, T.R. and Gruenauer, M.H.
GC-MS method for the quantitative analysis of carbofuran in water (212) 366
Nesbakken, R., see Jellum, E. (217) 231
Newland, P., see Thomas, A.H. (216) 367
Nice, E.C., Capp, M.W., Cooke, N. and O'Hare, M.J.
Comparison of short and ultrashort-chain alkylsilane-bonded silicas for the HPLC of proteins by hydrophobic interaction methods (218) 569
Nickless, G., see Roberts, D.J. (213) 73
Nicoletti, M., see Marini-Bettòlo, G.B. (213) 113
Niki, E., see Watanabe, N. (216) 406
Nikolov, R.N., see Mancheva, I.N. (213) 99
Nilsson, B.F. and Samuelson, O.
Chromatographic behaviour of aromatic compounds on non-polar phases with adsorbed quaternary ammonium salts (212) 1
Nilsson, L.B., see Westerlund, D. (211) 181
Nimura, N., Kasahara, Y. and Kinoshita, T.
Resolution of enantiomers of norepinephrine and epinephrine by reversed-phase HPLC (213) 327
Nishi, H., see Terabe, S. (212) 295
Nishi, S., see Hirai, H. (215) 195
Nishikawa, T., see Uobe, K. (214) 177
Nobuhara, Y., see Matsutera, E. (216) 374
Nohta, H., see Zaitsu, K. (211) 129
Noma, A.T., see Friedman, M. (219) 343
Norder, H., see Rydén, L. (215) 341
Nosanchuk, J.S., see Henion, J.D. (213) 475
Novák, J., see Vejrosta, J. (217) 167, (219) 37
Novotný, M., see McGuffin, V.L. (218) 179
Nowicki, H.G.
Detection of benzidines on thin-layer chromatograms with fluorescamine (211) 304
Nyborg, G., see Sisfontes, L. (216) 115
Nyman, P.-O., see Malmström, B.M. (215) 109

O

Obita, J.A., see Haken, J.K. (213) 55
Ochoa, J.-L.
Consideration of the nature of the lectin-carbohydrate interaction (215) 351
Oda, T., see Shinozawa, S. (212) 323
Ogamo, A., Matsuzaki, K., Uchiyama, H. and Nagasawa, K.
Hydrophobic interaction chromatography of mucopolysaccharides. Examination of fundamental conditions for fractionation of heparin on hydrophobic gels (213) 439
Ogawa, T., see Kainuma, K. (212) 126
Ogunlesi, M., see Hjertén, S. (215) 25
O'Hare, M.J., see Nice, E.C. (218) 569
Ohkuma, T., see Oka, K. (217) 387
Ohkura, Y., see Zaitsu, K. (211) 129
Ohlson, S., see Lowe, C.R. (215) 303
Ohnesorge, A.-B., see Vogt, W. (217) 91
Ohtsuka, K., see Okamoto, M. (219) 175
Ohzeki, H., see Watanabe, N. (216) 406
Ôi, N., Kitahara, H., Inda, Y. and Doi, T.
N-(1*R*, 3*R*)-*trans*-Chrysanthemoyl (*R*)-1-(*α*-naphthyl)ethylamine as a stationary phase for the separation of optical isomers by GC (213) 137
Ôi, N., Shiba, K., Tani, T., Kitahara, H. and Doi, T.
GC separation of some enantiomers on optically active copper(II) complexes (211) 274
Oka, K., Dobashi, Y., Ohkuma, T. and Hara, S.
Liquid column switching extraction and chromatography for programmed flow preparation (217) 387
Okada, Y., see Zaitsu, K. (211) 129
Okamoto, M. and Kishimoto, H.
Preparation and evaluation of N-(2-aminoethyl)-*γ*-aminopropyltrimethoxysilane-treated silica columns for the HPLC analysis of some aromatic amines (212) 251
Okamoto, M., Ohtsuka, K., Imai, J. and Yamada, F.
HPLC determination of acetaldehyde in wine as its lutidine derivative (219) 175
Olausson, Å., see Helmer, U. (211) 369
Olivelle, S., see Taylor, M.W. (219) 133
Olsson, I. and Låås, T.
Isoelectric focusing in agarose under denaturating conditions (215) 373
Oshima, F., see Kura, G. (219) 385
Oshima, R., Yoshikawa, A. and Kumanotani, J.
High-resolution GC separation of alditol acetates on fused-silica wall-coated open-tubular columns (213) 142
Osman, M.A. and Hill, H.H. Jr.
Response enhancement by metal doping of a silicon-selective flame ionization detector for GC (213) 397
Ossicini, L., see Fanali, S. (212) 374
Ostrovský, I., see Soják, L. (219) 225

Otson, R. and Williams, D.T.
Evaluation of a liquid-liquid extraction technique for water pollutants (212) 187
Ovchinnikov, Y.A., see Ulyashin, V.V. (215) 263

P

Pacáková, V., see Dufek, P. (211) 150
Pacáková, V., see Štulík, K. (213) 41
Pajurek, J., see Krejčí, M. (218) 167
Palusinski, O.A., see Bier, M. (211) 313
Panday, P., see Gandhe, B.R. (219) 297
Panfil, I., see Picman, A.K. (212) 379
Papazova, D. and Dimov, N.
Prediction of retention indices of aromatic hydrocarbons on phases of different polarity (216) 321
Papp-Hites, E., see Vigh, G. (214) 335
Parrado, C., see Alonso-Fernández, J.R. (217) 357
Pask-Hughes, R.A., Corran, P.H. and Calam, D.H.
Assay of the combined formulation of ergometrine and oxytocin by HPLC (214) 307
Patamia, M., see Marini-Bettòlo, G.B. (213) 113
Pauw, C. de, see De Pauw, C. (214) 95
Pauwels, J.-F., Carlier, M. and Sochet, L.-R.
Coupling of a low-pressure sampling technique to a gas chromatography (in French) (211) 247
Pav, J.W., see Lee, G.J.-L. (212) 65
Peal, J.A., see Bailey, E. (219) 285
Peña, J., see Alonso-Fernández, J.R. (217) 357
Peters, K.-R., see Haller, W. (211) 53
Petronio, B.M., see Laganà, A. (219) 263
Pettersson, B., see Buswell, J.A. (215) 99
Pfletschinger, J., see Mancheva, I.N. (213) 99
Phillips, D.V., see Ghias-Ud-Din, M. (211) 295
Picker, J.E. and Sievers, R.E.
Applications of selective complexation by a europium(III) coordination polymer sorbent for the pre-fractionation of volatile compounds (217) 275
Pickert, A., see Liebich, H.M. (217) 255
Picman, A.K., Panfil, I. and Towers, G.H.N.
Separation and identification of stereoisomers of sesquiterpene lactones by multiple development of thin-layer chromatograms (212) 379
Pillai, D.N. and Dilli, S.
Analysis of barbiturates by GC (Review) (220) 253
Plešek, J., see Plzák, Z. (212) 283
Pliška, V., Schmidt, M. and Fauchère, J.-L.
Partition coefficients of amino acids and hydrophobic parameters π of their side-chains as measured by TLC (216) 79
Plummer, D.T., see Vidal, C.J. (219) 71
Plzák, Z., Plešek, J. and Štíbr, B.
Reversed-phase ion-pair LC separations of some heteroborane anions (212) 283
Pokorný, S., see Janča, J. (211) 39
Polakowski, D., see Manecke, G. (215) 13
Poludniak, N.H., see Haeffner-Gormley, L. (214) 185
Ponsold, K., see Draffehn, J. (216) 63, (216) 69
Poole, C.F., Butler, H., Agnello, S.A., Sye, W.-F., Zlatkis, A. and Holzer, G.
Evaluation of substituted polyphenyl ethers as polar phases in GC (217) 39
Poole, C.F., Sye, W.-F., Zlatkis, A. and Spencer, P.S.
Determination of acrylamide in nerve tissue homogenates by electron-capture GC (217) 239
Popl, M., Vít, I. and Šmejkal, F.
Prediction of retention behaviour of some phenolic antioxidants (213) 363
Porath, J.
Development of modern bioaffinity chromatography (a review) (218) 241
Pospíšil, J., see Rotschová, J. (211) 299, (216) 251
Pownall, H.J., see Hancock, W.S. (216) 285
Poy, F., Visani, S. and Terrosi, F.
Automatic injection in high-resolution GC: a programmed temperature vaporizer as a general purpose injection system (217) 81
Preston, N.W. and Timberlake, C.F.
Separation of anthocyanin chalcones by HPLC (214) 222
Priboth, I., Engewald, W., Kühn, H. and Mühlstädt, M.
Influence of intramolecular interactions on the GC separation of epimers of dicyclopentadiene derivatives (in German) (219) 113
Přistoupil, T.I., Kramlová, M., Fričová, V., Ulrych, S., Kraml, J. and Jojková, K.
Characterization of polymerized stroma-free haemoglobin by chromatography, electrophoresis and electrofocusing in gels (213) 183
Přistoupil, T.I., Kramlová, M., Kraml, J. and Ulrych, S.
Chromatofocusing of pyridoxalated and polymerized human haemoglobin (219) 436
Přistoupil, T.I., Kramlová, M., Ulrych, S., Fričová, V. and Kraml, J.
Isoelectric focusing, gel chromatography and electrophoresis of pyridoxalated and polymerized stroma-free haemoglobin (219) 128
Přistoupil, T.I., see Kramlová, M. (211) 308
Pritchett, T., see Karmen, A. (217) 247
Prohaska, P.G., see Ramos, L.S. (211) 284
Proksch, P., Behl, H.M. and Rodriguez, E.
Detection and quantification of guayulins A and B in *Parthenium argentatum* (guayule) and F_1 hybrids by HPLC (213) 345
Prusík, Z., see Štrop, P. (214) 317

Punčochářová, J., see Kříž, J. (219) 53
Putter, I., see Gullo, V.P. (212) 234

Q

Quinlan, G.J., see Thomas, A.H. (216) 367
Quy, T. van, see Van Quy, T. (211) 177

R

Rabin, L.B., see Elliger, C.A. (216) 261
Rackham, D.M., see McKay, S.W. (214) 249
Raděj, Z., see Madajová, V. (216) 313
Radhakrishnan, A.N., see Ganapathy, V. (213) 307
Rajakylä, E.
Separation and determination of some organic acids and their sodium salts by HPLC (218) 695
Ramachandramurty, B., see Ganapathy, V. (213) 307
Ramos, L.S. and Prohaska, P.G.
Sephadex LH-20 chromatography of extracts of marine sediment and biological samples for the isolation of polynuclear aromatic hydrocarbons (211) 284
Raso, A.G., see Calixto, F.S. (216) 326
Raymond, J., Azanza, J.-L. and Fotso, M.
Hydrophobic interaction chromatography: a new method for sunflower protein fractionation (212) 199
Rayss, J. and Waksmundzki, A.
Temperature dependence of HETP for columns with small amounts of liquid stationary phase (217) 225
Remy, M.H., Guillochon, D. and Thomas, D.
Comparative study of native and chemically modified chymotrypsin as monomers, soluble polymers and membranes (215) 87
Renberg, L. and Lindström, K.
C_{18} reversed-phase trace enrichment of chlorinated phenols, guaiacols and catechols in water (214) 327
Repta, A.J., see Riley, C.M. (217) 405, (219) 235
Righetti, P.G., Tudor, G. and Ek, K.
Isoelectric points and molecular weights of proteins. A new table (Review) (220) 115
Riley, C.M., Sternson, L.A. and Repta, A.J.
HPLC of *cis*-dichlorodiammineplatinum(II) using chemically-bonded and solvent-generated ion exchangers (217) 405
Riley, C.M., Sternson, L.A. and Repta, A.J.
HPLC of inorganic platinum(II) complexes using solvent-generated anion exchangers. II. The effect of electrolytes on solute retention (219) 235
Riley, C.M., Tomlinson, E. and Hafkenscheid, T.L.
Structural effects in enthalpy/entropy compensated and non-compensated behaviour in ion-pair reversed-phase HPLSC (218) 427
Riquelme, M.E., see Einarsson, M. (215) 45
Roberts, D.J., Cooke, M. and Nickless, G.
Determination of polychlorinated alkanes via carbon skeleton capillary GC (213) 73
Robinson, J.P., see DiMari, S.J. (213) 91
Rodriguez, E., see Proksch, P. (213) 345
Roets, E., see Hoogmartens, J. (219) 431
Rogozhin, S.V., see Ryashentsev, V.Y. (216) 346
Rogozhin, S.V., see Zaslavsky, B.Y. (212) 13, (216) 103
Rogstad, A. and Yndestad, M.
Analysis of xylazine in biological material by GC using packed and capillary columns (216) 350
Romann, A., see Grob, K. Jr. (214) 118
Rosenbauer, D., see Steudel, R. (216) 399
Ross, W.J., see McKay, S.W. (214) 249
Rossetti, Z.L., see Meek, J.L. (211) 15
Rotatori, M., see Laganà, A. (219) 263
Roth, M., see Vejrosta, J. (217) 167, (219) 37
Rotschová, J. and Pospíšil, J.
Antioxidants and stabilizers. LXXXVI. Use of chromatography in the investigation of the mechanism of action of amine antidegradants (211) 299
Rotschová, J., Taimr, L. and Pospíšil, J.
Antioxidants and stabilizers. LXXXVII. The chromatographic behaviour of transformation products of an antioxidant, N,N'-diphenyl-1,4-phenylenediamine (216) 251
Roumeliotis, P. and Unger, K.K.
Assessment and optimization of system parameters in size exclusion separation of proteins on diol-modified silica columns (218) 535
Rudolph, J., Ehhalt, D.H., Khedim, A. and Jebsen, C.
Determination of C_2-C_5 hydrocarbons in the atmosphere at low parts per 10^9 to high parts per 10^{12} levels (217) 301
Rusek, M., see Krejčí, M. (218) 167
Ryashentsev, V.Y., Voskoboinikov, M.A., Vainerman, E.S. and Rogozhin, S.V.
Behaviour of biomolecules in water-organic solvent-inorganic salt two-phase ternary systems (216) 346
Ryba, M.
Reversed-phase LCC of pyrimidine and purine derivatives. I. Unbuffered binary aqueous organic mobile phases on octadecylsilica (219) 245
Ryba, M. and Beránek, J.
LC separation of purines, pyrimidines and

their nucleosides on silica gel columns (211) 337
Rydén, L. and Norder, H.
Covalent chromatography as a means of isolating thiol peptides from large proteins. Application to human ceruloplasmin (215) 341

S

Sacco, D. and Dellacherie, E.
LC of cephalosporin C and α-amino acid mixtures on polyfunctional polystyrene resins (211) 79
Sadler, J.E., Beyer, T.A. and Hill, R.L.
Affinity chromatography of glycosyltransferases (215) 181
Sahlberg, B.-L., Axelson, M., Collins, D.J. and Sjövall, J.
Analysis of isomeric ethynylestradiol glucuronides in urine (217) 453
Saint-Jalm, Y., see Moree-Testa, P. (217) 197
Sairam, M.R.
Isolation of hormonal proteins and antibodies by affinity chromatography (215) 143
Sakaguchi, U., see Nakazawa, H. (213) 323
Sakakibara, S., see Takaya, T. (215) 279
Sakodynskii, K.I.
New data on M.S. Tswett's life and work (Review) (220) 1
Sakurada, T., see Kizaki, H. (211) 409
Samay, G., Füzes, L., Cser, F. and Bodor, G.
Comparison of dimensions obtained by size-exclusion chromatography and X-ray diffraction of rigid molecules (218) 473
Sampson, E.J., see Culbreth, P.H. (212) 221
Samuelson, O., see Nilsson, B.F. (212) 1
Samusenko, A.L., see Golovnya, R.V. (217) 183
Sander-Wewer, M., see Veh, R.W. (212) 313
Sasaki, H., see Kato, Y. (211) 383
Sass, S. and Stutz, M.H.
TLC of some sulfur and nitrogen mustards (213) 173
Sato, T., see Chibata, I. (215) 93
Saxby, M.J., see Chaytor, J.P. (214) 135
Scalia, S., see Wilson, I.D. (212) 211
Scatton, B., see Semerdjian-Rouquier, L. (218) 663
Schauer, R., see Veh, R.W. (212) 313
Schill, G., see Denkert, M. (218) 31
Schlett, R., see Von Stetten, O. (218) 591
Schmidbaur, H., see Jandik, P. (213) 47
Schmidt, J., John, M. and Wandrey, C.
Rapid separation of malto-, xylo- and cello-oligosaccharides (DP 2-9) on cation-exchange resin using water as eluent (213) 151
Schmidt, M., see Pliška, V. (216) 79
Schmidt, R., Ziegenhagen, E. and Dose, K.
HPLC of trichothecenes. I. Detection of T-2 toxin and HT-2 toxin (212) 370
Schmitter, J.M., see Krupčík, J. (213) 189
Schneider, W. and Degen, P.H.
Simultaneous determination of diclofenac sodium and its hydroxy metabolites by capillary column GC with electron-capture detection (217) 263
Schoenl, M., see Davis, P.J. (219) 148
Schoenmakers, P.J., Billiet, H.A.H. and De Galan, L.
Systematic study of ternary solvent behaviour in reversed-phase LC (218) 261
Schoenmakers, P.J., see Billiet, H.A.H. (218) 443
Scholander, E., see Einarsson, M. (215) 45
Scholten, A.H.M.T., Brinkman, U.A.T. and Frei, R.W.
Fluorescence detection of chloroanilines in LC using a post-column reaction with fluorescamine. Comparison of reactor types and mixing tees (218) 3
Scholten, A.H.M.T., see Brinkman, U.A.T. (217) 463
Schönecker, B., see Draffehn, J. (216) 63, (216) 69
Schoo, G.M., see Eck, W.S. (219) 171
Schrenker, H.
Effect of mobile phase pre-heating on HPLC column performance: a new type of column thermostat (213) 243
Schuller, D., see Butte, W. (214) 59
Schulten, H.-R. and Soldati, F.
Identification of ginsenosides from *Panax ginseng* in fractions obtained by HPLC by field desorption MS, multiple internal reflection IR spectroscopy and TLC (212) 37
Schurig, V. and Weber, R.
Manganese(II)-bis(3-heptafluorobutyryl-1*R*-camphorate): a versatile agent for the resolution of racemic cyclic ethers by complexation GC (217) 51
Schwartz, H., see Yu, T.J. (218) 519
Schwertfeger, G., see Vogt, W. (217) 91
Scobell, H.D. and Brobst, K.M.
Rapid high-resolution separation of oligosaccharides on silver form cation-exchange resins (212) 51
Scott, A., see Jane, I. (214) 243
Scott, I.M., see Martin, G.C. (219) 167
Sebille, B., see Zini, R. (216) 191
Sechovec, J., see Franc, J. (212) 139
Segura, R. and Navarro, X.
Use of metal salts as fluorescence-inducing reagents in TLC (217) 329
Selden, A.C., see Murphy, R. (211) 160
Sellier, F., Tersac, G. and Guiochon, G.
Thermal stability of a poly(oxy aryl sulfonyl arylene) used as stationary phase in GLC (in French) (219) 197
Sellier, F., Tersac, G. and Guiochon, G.
Polarity of a poly(oxy aryl sulfonyl arylene)

used as stationary phase in GLC (in French) (219) 213
Semenovskaya, T.D., see Kolotinskaya, E.V. (212) 133
Semerdjian-Rouquier, L., Bossi, L. and Scatton, B.
Determination of 5-hydroxytryptophan, serotonin and 5-hydroxyindoleacetic acid in rat and human brain and biological fluids by reversed-phase HPLC with electrochemical detection (218) 663
Sepaniak, M.J. and Yeung, E.S.
HPLC studies of coal liquids by laser-based detectors (211) 95
Serpinet, J., see Morel, D. (214) 202
Ševčík, J. and Löwentap, M.S.H.
New criterion for polarity of stationary phases in GC (217) 139
Severson, R.F., McDuffie, K.L., Arrendale, R.F., Gwynn, G.R., Chaplin, J.F. and Johnson, A.W.
Rapid method for the analysis of tobacco nicotine alkaloids (211) 111
Shaltiel, S., see Halperin, G. (215) 211
Shanfield, H., see Zhou, L. (217) 341
Sharma, R.K., see Gandhe, B.R. (219) 297
Sharon, N., see Lis, H. (215) 361
Sharpe, R.W.L., see Jane, I. (214) 243
Shepherd, M.J. and Gilbert, J.
Analysis of additives in plastics by high-performance size-exclusion chromatography (218) 703
Shiba, K., see Ôi, N. (211) 274
Shibukawa, A., see Nakagawa, T. (211) 1
Shinde, S.K., see Gandhe, B.R. (219) 297
Shinozawa, S. and Oda, T.
Determination of adriamycin (doxorubicin) and related fluorescent compounds in rat lymph and gall by HPLC (212) 323
Shono, T., see Funazo, K. (211) 361
Shono, T., see Tanaka, M. (219) 108
Shono, T., see Wu, H.-L. (219) 312
Shrubsall, P.R., see McKay, S.W. (214) 249, (219) 325
Sievers, R.E., see Picker, J.E. (217) 275
Sievers, S., see König, W.A. (217) 71
Simpson, R., see Abbott, S. (218) 123
Sindelar, R.D., see Kutney, J.P. (214) 152
Singh, J., see Cochrane, W.P. (217) 289
Siriwardana, M.G., see Lafont, P. (219) 162
Sisfontes, L., Nyborg, G., Svensson, L. and Blomstrand, R.
Separation of complex long chain fatty acid mixtures by high-performance glass capillary GC (216) 115
Sjögren, E., see Denkert, M. (218) 31
Sjövall, J., see Sahlberg, B.-L. (217) 453
Sleeman, H.K., see Brown, N.D. (212) 361
Sluis, W.G. van der, see Van der Sluis, W.G. (214) 349
Small, D.A.P., Atkinson, T. and Lowe, C.R.
High-performance liquid affinity chromatography of enzymes on silica-immobilised triazine dyes (216) 175
Small, D.A.P., see Lowe, C.R. (215) 303
Šmejkal, F., see Popl, M. (213) 363
Smit, A.L.C., see Tijssen, R. (218) 137
Smith, A.E., see Ghias-Ud-Din, M. (211) 295
Smith, B.E.F., see Truedsson, L.-Å. (214) 291
Smith, E., see Barkan, S. (219) 81
Smith, G.C., see Guise, G.B. (214) 69
Smith, J.M., see McKay, S.W. (214) 249, (219) 325
Smith, M.D., see Baaske, D.M. (212) 339
Smith, R.V., see Kerr, K.M. (219) 317
Smith, T.A., see Bird, C.R. (214) 263
Snyder, L.R. and Glajch, J.L.
Solvent strength of multicomponent mobile phases in LSC. Binary-solvent mixtures and solvent localization (214) 1
Snyder, L.R., Glajch, J.L. and Kirkland, J.J.
Theoretical basis for systematic optimization of mobile phase selectivity in LSC. Solvent-solute localization effects (218) 299
Snyder, L.R., see Glajch, J.L. (214) 21
Sochet, L.-R., see Pauwels, J.-F. (211) 247
Soczewiński, E. and Wawrzynowicz, T.
TLC as a pilot technique for the optimization of preparative column chromatography (218) 729
Soejarto, D.D., see Ahmed, M.S. (213) 340
Soják, L., Kraus, G., Ostrovský, I., Královičová, E. and Farkaš, P.
High-resolution GC with liquid crystal glass capillaries. III. Separation of hydrocarbon isomers on cholesteric mesophase (in German) (219) 225
Sokoloski, T.D., see Wu, C.C. (216) 239
Soldati, F., see Schulten, H.-R. (212) 37
Solinas, V., Gessa, C. and Falchi Delitala, L.
HPLC analysis of carvacrol and thymol in the essential oil of *Thymus capitatus* (219) 332
Solomon, B., Lotan, N. and Katchalski-Katzir, E.
Interaction of glucose oxidase with Blue Dextran (215) 121
Špaček, P., see Helmich, O. (213) 105
Sparrow, J.T., see Hancock, W.S. (216) 285
Spencer, P.S., see Poole, C.F. (217) 239
Srisukh, D., see Haken, J.K. (219) 45
Stárková, B., see Štulík, K. (213) 41
Stein, R.L.
Separation of choline and acetylcholine by cation-exchange chromatography (214) 148
Stensiö, K.-E., see Helmer, U. (211) 369
Sternson, L.A., see Riley, C.M. (217) 405, (219) 235
Stetten, O. von, see Von Stetten, O. (218) 591
Steudel, R. and Rosenbauer, D.
Separation of cyclic sulphur-nitrogen compounds by HPLC. LXXVIII (216) 399
Štíbr, B., see Plzák, Z. (212) 283
Strickler, M.P., see Brown, N.D. (212) 361
Strickler, M.P., see Eck, W.S. (219) 171
Strid, L., see Malmström, B.M. (215) 109
Štrop, P., Borvák, J., Kašička, V., Prusík, Z. and Morávek, L.
Isolation of human haemopexin by

bioaffinity chromatography on haeme-Sepharose (214) 317
Stuart, K.L., see Kutney, J.P. (214) 152
Štulík, K., Pacáková, V. and Stárková, B.
Carbon pastes for voltammetric detectors in HPLC (213) 41
Stutz, M.H., see Sass, S. (213) 173
Stuurman, H.W. and Wahlund, K.-G.
Separation of proton-donating solutes by LC with a strong proton acceptor, tri-*n*-octylphosphine oxide, in the liquid stationary phase (218) 455
Sucher, J., see Falkowski, P.G. (213) 349
Sudraud, G., see Coustard, J.M. (219) 338
Sutherland, I.W.
Affinity column for separation of pyruvylated and non-pyruvylated polysaccharides (213) 301
Svendsen, H. and Greibrokk, T.
HPLC determination of biogenic amines. I. Use of aqueous acidic mobile phases with silica columns (212) 153
Svendsen, H. and Greibrokk, T.
HPLC determination of biogenic amines. II. Comparison of detection methods (213) 429
Svensson, L., see Sisfontes, L. (216) 115
Sweetman, J.A. and Boettner, E.A.
Detection of brominated and chlorinated organics by a GC microcoulometric detector. Effect of pyrolysis tube conditions (212) 115
Sybilska, D., see Kemula, W. (218) 465
Sye, W.-F., see Poole, C.F. (217) 39, (217) 239
Synge, R.L.M.
The Faraday Society's discussion at Reading in 1949 and the exploitation of molecular-sieve effects for chemical separations (215) 1
Szakács-Pintér, M., see Molnár-Perl, I. (216) 219
Szczepaniak, W., see Nawrocki, J. (212) 29
Szepesi, G. and Gazdag, M.
Improved HPLC method for the analysis of insulins and related compounds (218) 597
Szepesi, G., see Gazdag, M. (218) 603
Szymanowicz, A., Legentil, J.-P. and Malgras, A.
Improved single-column procedure for the amino acid analysis of collagen-like proteins (211) 175
Szyszko, E. and Wejman, W.
GC determination of noxyptyline in substance, tablets and in biological material (219) 291

T

Taeye, L. de, see De Taeye, L. (214) 95
Taimr, L., see Rotschová, J. (216) 251
Takács, J.M.
Contribution to the theory of open-tubular (capillary) columns in GLC (213) 371
Takács, J.M., see Czerwiec, Z. (214) 47
Takada, K., Ito, R., Tokunaga, K., Kobayashi, T. and Takao, T.
Determination of 7-dehydrocholesterol in rat skin and liver by HPLC (216) 385
Takagi, T.
HPLC of protein polypeptides on porous silica gel columns (TSK-GEL SW) in the presence of sodium dodecyl sulphate: comparison with SDS-polyacrylamide gel electrophoresis (219) 123
Takao, T., see Takada, K. (216) 385
Takata, R.H., see Matsumoto, H. (211) 403
Takaya, T., Kishida, Y. and Sakakibara, S.
Determination of the optical purity of amino acids by HPLC. Modification of the Manning and Moore procedure (215) 279
Takeda, R., see Uobe, K. (214) 177
Takeuchi, T. and Ishii, D.
High-performance micro packed flexible columns in LC (213) 25
Takeuchi, T. and Ishii, D.
Application of ultra-micro HPLC to trace analysis (218) 199
Takeuchi, T., Watanabe, Y., Matsuoka, K. and Ishii, D.
High-back-pressure LC. I. Development of micro-HPLC using liquefied alkanes as the mobile phase (216) 153
Takeuchi, T., see Ishii, D. (218) 189
Tanaka, M., Mizobuchi, Y., Kuroda, T. and Shono, T.
Sorption behaviour of several organic compounds on 6-deoxycyclodextrin polyurethane resins (219) 108
Tanaka, M., see Funazo, K. (211) 361
Tanaka, M., see Wu, H.-L. (219) 312
Tani, T., see Ôi, N. (211) 274
Tanzer, F.S., see Desiderio, D.M. (217) 437
Tauber-Finkelstein, M., see Halperin, G. (215) 211
Taylor, M.W., Hershey, H.V., Levine, R.A., Coy, K. and Olivelle, S.
Improved method of resolving nucleotides by reversed-phase HPLC (219) 133
Terabe, S., Nishi, H. and Ando, T.
Separation of cytochromes *c* by reversed-phase HPLC (212) 295
Terrosi, F., see Poy, F. (217) 81
Tersac, G., see Sellier, F. (219) 197, (219) 213
Tesařík, K., see Krejčí, M. (218) 167
Thomas, A.H., Newland, P. and Quinlan, G.J.
Identification and determination of the qualitative composition of nystatin using TLC and HPLC (216) 367
Thomas, D., see Remy, M.H. (215) 87
Thomsen, M. and Willumsen, D.
Quantitative ion-pair extraction of 4(5)-methylimidazole from caramel colour and its determination by reversed-phase ion-pair LC (211) 213

Thorn, W., see Weiland, E. (214) 156
Tichá, M., see Hořejší, V. (216) 43
Tieckelmann, H., see Lee, G.J.-L. (212) 65
Tijssen, R., Bleumer, J.P.A., Smit, A.L.C. and Van Kreveld, M.E.
Microcapillary LC in open tubular columns with diameters of 10-50 μm. Potential application to chemical ionization MS detection (218) 137
Tillement, J.-P., see Zini, R. (216) 191
Timberlake, C.F., see Preston, N.W. (214) 222
Tishbee, A., see Weiner, S. (213) 501
Tjessem, K., see Barth, T. (214) 83
Tokunaga, K., see Takada, K. (216) 385
Tomlinson, E., see Hafkenscheid, T.L. (218) 409
Tomlinson, E., see Riley, C.M. (218) 427
Tosa, T., see Chibata, I. (215) 93
Tóth, G., Wéber, M. and Kling, F.
Adsorption chromatographic separation of [125I]progesterone-succinyl-tyrosine methyl ester (213) 511
Towers, G.H.N., see Picman, A.K. (212) 379
Treiber, L.R.
Quantitative analysis of Cephamycin C in fermentation broths by means of thin-layer spectrodensitometry (213) 129
Trimble, J., see Desiderio, D.M. (217) 437
Tříska, J., see Březina, M. (219) 179
Truedsson, L.-Å. and Smith, B.E.F.
Study of retention behaviour of primary, secondary and tertiary anilines in normal- and reversed-phase LC (214) 291
Tsiatsios, A., see Katsanos, N.A. (213) 15
Tsuboi, K., see Tsuda, T. (214) 283
Tsuda, T., Tsuboi, K. and Nakagawa, G.
Open-tubular microcapillary LC with 20-μm I.D. columns (214) 283
Tsukada, Y., see Hirai, H. (215) 195
Tsunashima, Y., see Nakazawa, T. (211) 388
Tsutsui, M., see Uobe, K. (214) 177
Tudor, G., see Righetti, P.G. (220) 115
Tulliez, J.E. and Durand, E.F.
A fast and simple HPLC assay for aryl hydrocarbon hydroxylase (219) 411
Turková, J., Bláha, K., Horáček, J., Vajčner, J., Frydrychová, A. and Čoupek, J.
Hydroxyalkyl methacrylate gels derivatized with epichlorohydrin as supports for large-scale and high-performance affinity chromatography (215) 165
Tweedie, H.A., see Betts, T.J. (213) 317
Tychopoulos, V., see Tyman, J.H.P. (213) 287
Tyihák, E., Mincsovics, E., Kalász, H. and Nagy, J.
Optimization of operating parameters in overpressured TLC (211) 45
Tyman, J.H.P., Tychopoulos, V. and Colenutt, B.A.
Long-chain phenols. XXI. Quantitative analysis of the phenolic lipids in technical cashew nut-shell liquid, from *Anacardium occidentale*, by HPLC (213) 287

U

Uchiyama, H., see Ogamo, A. (213) 439
Uebori, M., see Kuwata, K. (211) 378
Ugland, K., Lundanes, E., Greibrokk, T. and Bjørseth, A.
Determination of chlorinated phenols by HPLC (213) 83
Ui, N.
High-speed gel filtration of glycopolypeptides in 6 *M* guanidine hydrochloride (215) 289
Ukai, S., see Kawase, S. (213) 265
Ullman, A.H. and Houston, R.
Determination of residual epichlorohydrin in middle cut alkylglycidyl ethers by headspace GC (211) 398
Ulrych, S., see Přistoupil, T.I. (213) 183, (219) 128, (219) 436
Ulyashin, V.V., Deigin, V.I., Ivanov, V.T. and Ovchinnikov, Y.A.
Size-exclusion HPLC of protected peptides (215) 263
Unger, K.K., see Roumeliotis, P. (218) 535
Uno, T., see Nakagawa, T. (211) 1
Uobe, K., Takeda, R., Wato, M., Nishikawa, T., Yamaguchi, S., Koshimura, T., Kawaguchi, Y. and Tsutsui, M.
Trimethylsilylation reaction of prostaglandin-E methyl ester with various trimethylsilylating reagents (214) 177

V

Vaidyanathaswamy, R., see Gandhe, B.R. (219) 297
Vainerman, E.S., see Ryashentsev, V.Y. (216) 346
Vajčner, J., see Turková, J. (215) 165
Valášková, I., see Van Quy, T. (211) 177
Van Arkel, J., see Van der Sluis, W.G. (214) 349
Van Dam, H.C., see Laurent, C. (218) 83
Van den Berg, J.H.M., see Deelder, R.S. (218) 327
Van der Sluis, W.G., Van Arkel, J., Fischer, F.C. and Labadie, R.P.
TLC assay of photoactive compounds (furocoumarins) using the fungus *Penicillium expansum* as a test organism (214) 349
Van Dyck, J., see Verzele, M. (214) 95
Van Kreveld, M.E., see Tijssen, R. (218) 137
Van Quy, T., Valášková, I. and Majer, J.
Separation of metal ions by paper

electrophoresis in N-(phosphono)iminodisulphuric acid and glycine-N,N-bis(methylphosphonic acid) (in German) (211) 177
Vanderhaeghe, H., see Decoster, W. (211) 223
Vanderhaeghe, H., see Hoogmartens, J. (219) 431
Vanderslice, J.T., Brown, J.F., Beecher, G.R., Maire, C.E. and Brownlee, S.G. Automation of a complex HPLC system. Procedures and hardware for a vitamin B_6 model system (216) 338
Vanhaelen, M., see Brassinne, C. (213) 170
Vanhaelen-Fastré, R., see Brassinne, C. (213) 170
Varga-Puchony, Z., see Vigh, G. (214) 335
Vasileva, R., Jakab, M. and Haskó, F. Application of ion-exchange chromatography for the production of human albumin (216) 279
Vasilieva, V.S., see Kagan, M.Z. (219) 183
Vattis, D., Katsanos, N.A., Karaiskakis, G., Lycourghiotis, A. and Kotinopoulos, M. Conversions in catalytic deamination calculated by stopped-flow GC (214) 171
Veh, R.W., Michalski, J.-C., Corfield, A.P., Sander-Wewer, M., Gies, D. and Schauer, R. New chromatographic system for the rapid analysis and preparation of colostrum sialyloligosaccharides (212) 313
Vejrosta, J., Roth, M. and Novák, J. Comparison of the distribution constants of benzene in different chromatographic sorbent-gas systems, determined by direct measurement of sorption equilibria and calculated from GC retention data (217) 167
Vejrosta, J., Roth, M. and Novák, J. Comparison of the distribution constants of acetone in different chromatographic sorbent-gas systems, determined by direct measurement of sorption equilibria and calculated from GC retention data (219) 37
Veland, K., see Francis, G.W. (219) 379
Venn, R.F., see Glasel, J.A. (213) 337
Verhaar, L.A.T. and Kuster, B.F.M. LC of sugars on silica-based stationary phases (Review) (220) 313
Verschoyle, R.D., see Bailey, E. (219) 285
Verzele, M., De Taeye, L., Van Dyck, J., De Decker, G. and De Pauw, C. HPLC of *Vinca rosea* alkaloids and the correlation of plate height and molecular weight (214) 95
Verzele, M. and Dewaele, C. Stationary phase characterization in HPLC. A test for trace metal activity in octadecyl bonded silica gel (217) 399
Vespalec, R., see Krejčí, M. (219) 61
Vidal, C.J., Elmi-Akhounie, E., Chai, M.S.Y. and Plummer, D.T. Affinity chromatography of acetylcholinesterase. The use of Amberlite CG-120 for dissociating the enzyme-inhibitor complex (219) 71
Vierkorn-Rudolph, B. and Bächmann, K. Determination of trace amounts of hydrogen halides in the atmosphere by collection in sampling tubes and subsequent derivatization and GC (217) 311
Vigh, G., Varga-Puchony, Z., Papp-Hites, E., Hlavay, J. and Balogh, S. Determination of chlorophacinone in formulations by reversed-phase ion-pair chromatography (214) 335
Vilenchik, L.Z., see Janča, J. (211) 39
Visani, S., see Poy, F. (217) 81
Vít, I., see Popl, M. (213) 363
Vita-Finzi, P., see Cairati, L. (216) 402
Vodička, L., see Březina, M. (219) 179
Vodička, L., see Kříž, J. (219) 53
Voelter, W., Huber, R. and Zech, K. Fluorescence labelling in trace analysis of biological samples. Simultaneous determination of free fatty acids and related carboxylic compounds (217) 491
Vogt, W., Jacob, K., Ohnesorge, A.-B. and Schwertfeger, G. Application of the split-splitless injector to environmental analysis (217) 91
Vogt, W., see Meyer, H.D. (217) 473
Von Arx, E., see Faupel, M. (211) 262
Von Stetten, O. and Schlett, R. HPLC of ^{125}I-labelled proteins with on-line detection (218) 591
Voskoboinikov, M.A., see Ryashentsev, V.Y. (216) 346
Vouros, P., see Yu, T.J. (218) 519
Vries, G. de, see De Vries, G. (217) 463

W

Wada, O., see Arakawa, Y. (216) 209
Wahlund, K.-G. Separation of acidic drugs in the µg/ml range in untreated blood plasma by direct injection on LC columns (218) 671
Wahlund, K.-G., see Stuurman, H.W. (218) 455
Wakeford, B., see Cochrane, W.P. (217) 289
Waksmundzki, A., see Rayss, J. (217) 225
Walendziak, L., see Wroński, M. (211) 252
Walker, R.F., see Goldberg, P.A. (212) 93
Walkley, J., see Cosgrove, B.A. (216) 161
Wall, R.A., see Ghaemi, Y. (212) 271
Waller, D.P., see Ahmed, M.S. (213) 340
Walters, D.G., Foster, P.M.D. and Cottrell, R.C. HPLC of progesterone and its metabolites (219) 152
Wandrey, C., see Schmidt, J. (213) 151
Wang Chinghai, Xia Aizhu, Wang Guanghua and Lu Peichang GC behaviour of chemically bonded microparticulate silica (217) 191

Wang, F.-S., see Zhou, L. (217) 341
Wang Guanghua, see Wang Chinghai (217) 191
Waser, P.G., see Bock, U.E.G. (213) 413
Watanabe, H., see Hikino, H. (211) 123
Watanabe, N., Ohzeki, H. and Niki, E.
Enantiomeric resolution of amino acids by high-performance ligand-exchange chromatography using a chemically modified hydrophilic porous polymer gel (216) 406
Watanabe, Y., see Takeuchi, T. (216) 153
Wato, M., see Uobe, K. (214) 177
Wawrzynowicz, T., see Soczewiński, E. (218) 729
Weber, J.D., see Barkan, S. (219) 81
Wéber, M., see Tóth, G. (213) 511
Weber, R., see Schurig, V. (217) 51
Weiland, E., Thorn, W. and Bläker, F.
Sialic acid quantitation by analytical isotachophoresis (214) 156
Weiner, S. and Tishbee, A.
Separation of Dns-amino acids using reversed-phase HPLC: a sensitive method for determining N-termini of peptides and proteins (213) 501
Wejman, W., see Szyszko, E. (219) 291
Welling, P.L.M., see Brinkman, U.A.T. (217) 463
Wellington, C.A., see Jones, P. (213) 357
Wendelboe, J.F., see Johansen, I. (217) 317
Wentworth, W.E., see Chen, E.C.M. (217) 151
Wepsic, H.T., see Hirai, H. (215) 195
Wessel, D., see Certa, U. (211) 257
Westerlund, D., Nilsson, L.B. and Jaksch, Y.
Straight-phase ion-pair chromatography of zimelidine and similar divalent amines. II. The chromatographic system (211) 181
Westwood, R. and Hairsine, P.W.
Use of routine preparative HPLC in the separation of isomers (219) 140
Wetlaufer, D.B., see Haeffner-Gormley, L. (214) 185
Whall, T.J.
Determination of streptomycin sulfate and dihydrostreptomycin sulfate by HPLC (219) 89
White, P.C., see Jane, I. (214) 243
Wilchek, M., see Miron, T. (215) 55
Williams, D.T., see Otson, R. (212) 187
Willumsen, D., see Thomsen, M. (211) 213
Wilson, A.M. and Bushway, R.J.
HPLC determination of azinphos methyl and azinphos methyl oxon in fruits and vegetables (214) 140
Wilson, I.D., Scalia, S. and Morgan, E.D.
Reversed-phase TLC for the separation and analysis of ecdysteroids (212) 211
Wintersteiger, R., see Gübitz, G. (218) 51
Wojnowski, W., see Becker, B. (219) 302
Wold, S., see Jellum, E. (217) 231
Wöll, J., see Liebich, H.M. (217) 255
Woodhouse, R.N., see Anderson, A. (212) 347
Wroński, M. and Walendziak, L.
Photometric monitoring of thiols by means of a thiomercurimetric detector (211) 252
Wu, C.C., Sokoloski, T.D., Burkman, A.M. and Wu, L.S.
Separation, identification and quantitation of nitroglycerin and its metabolic or hydrolysis products (216) 239
Wu, C.-M., see Wu, J.L.-P. (214) 234
Wu, H.-L., Funazo, K., Tanaka, M. and Shono, T.
Electron-capture GC determination of sulphide as a new pentafluorobenzyl derivative (219) 312
Wu, J.L.-P. and Wu, C.-M.
HPLC separation of shallot volatile oil (214) 234
Wu, L.S., see Wu, C.C. (216) 239

X

Xia Aizhu, see Wang Chinghai (217) 191

Y

Yamada, F., see Okamoto, M. (219) 175
Yamada, S., see Desiderio, D.M. (217) 437
Yamaguchi, S., see Uobe, K. (214) 177
Yamashiro, D. and Li, C.H.
Partition chromatography and HPLC of β-lipotropin and synthetic β-endorphin analogues (215) 255
Yamauchi, Y., Murakami, T. and Kumanotani, J.
Separation of urushiol by HPLC on an 8% octa,decylsilane chemically bonded silica gel column with electrochemical detection. Analysis of urushiol in the sap of lac trees (*Rhus vernicifera*) and that in the Japanese lac-making process (214) 343
Yamazaki, S. and Yoneda, H.
Chromatographic study of optical resolution. VIII. Theoretical study of the chromatographic behaviour of the enantiomers of racemic complex cations on a cation-exchange column (219) 29
Yamazaki, Y., see Kuwata, K. (211) 378
Yamskov, I.A., Berezin, B.B., Davankov, V.A., Zolotarev, Y.A., Dostavalov, I.N. and Myasoedov, N.F.
Ligand-exchange chromatography of amino acid racemates on Separon gels containing L-proline or L-hydroxyproline groupings (217) 539

Yau, W.W. and Kirkland, J.J.
Comparison of sedimentation field-flow fractionation with chromatographic methods for particulate and high-molecular-weight macromolecular characterizations (218) 217
Yeung, E.S., see Sepaniak, M.J. (211) 95
Yndestad, M., see Rogstad, A. (216) 350
Yoneda, H., see Nakazawa, H. (213) 323
Yoneda, H., see Yamazaki, S. (219) 29
Yoshikawa, A., see Oshima, R. (213) 142
Young, H.
Direct desorption of traps for capillary column GC (214) 197
Yu, T.H., see Arakawa, Y. (216) 209
Yu, T.J., Schwartz, H., Giese, R.W., Karger, B.L. and Vouros, P.
Analysis of N-acetyl-N,O,S-permethylated peptides by combined LC-MS (218) 519

Z

Zaitsu, K., Okada, Y., Nohta, H., Kohashi, K. and Ohkura, Y.
Assay for catechol-O-methyltransferase by HPLC with fluorescence detection (211) 129
Zaslavsky, B.Y., Miheeva, L.M. and Rogozhin, S.V.
Parameterization of hydrophobic properties of aqueous polymeric biphasic systems and water-organic solvent systems (212) 13
Zaslavsky, B.Y., Miheeva, L.M. and Rogozhin, S.V.
Comparison of conventional partitioning systems used for studying the hydrophobicity of polar organic compounds (216) 103
Zech, K., see Voelter, W. (217) 491
Zeman, A. and Koch, K.
Determination of odorous volatiles in air using chromatographic profiles (216) 199
Zhou, L., Shanfield, H., Wang, F.-S. and Zlatkis, A.
Fluorescence-inducing procedures for use in HPTLC (217) 341
Ziegenhagen, E., see Schmidt, R. (212) 370
Zini, R., Barre, J., Bree, F., Tillement, J.-P. and Sebille, B.
Evidence for a concentration-dependent polymerization of a commercial human serum albumin (216) 191
Zinkevich, E.P., see Kagan, M.Z. (219) 183
Zlatkis, A., see Poole, C.F. (217) 39, (217) 239
Zlatkis, A., see Zhou, L. (217) 341
Zolotarev, Y.A., see Yamskov, I.A. (217) 539
Žyvný, K., see Dufek, P. (211) 150

SUBJECT INDEX

VOLUMES 211—220

The volume number is placed in parentheses before the page number

A

Acetaldehyde
HPLC determination of acetaldehyde in wine as its lutidine derivative (219) 175

Acetone
GC determination of water in acetone (212) 121

Comparison of the distribution constants of acetone in different chromatographic sorbent-gas systems, determined by direct measurement of sorption equilibria and calculated from GC retention data (219) 37

Acetonitrile
Recovery of HPLC-grade acetonitrile by spinning-band distillation (213) 331

Acetylamino acid amides
Partition chromatography and HPLC of β-lipotropin and synthetic β-endorphin analogues (215) 255

Acetylcholine
Separation of choline and acetylcholine by cation-exchange chromatography (214) 148

Acetylcholinesterase
Affinity chromatography of acetylcholinesterase. The use of Amberlite CG-120 for dissociating the enzyme-inhibitor complex (219) 71

Acetyl groups
Determination of acetyl and formyl groups by GLC (211) 347

Acidic pharmaceuticals
Alkylation with alkyl halides as a derivatization method for the GC determination of acidic pharmaceuticals (Review) (220) 275

Aconitine alkaloids
Determination of aconitine alkaloids by HPLC (211) 123

Acrylamide
Determination of acrylamide in nerve tissue homogenates by electron-capture GC (217) 239

Actinomycins
TLC of closely related polypeptides on silanized silica gel (211) 29

Adenine nucleotides
Separation of cAMP from adenine nucleotides and nucleosides by electrophoresis on cellulose acetate membranes (211) 409

Adenosine
High-performance displacement chromatography (218) 365

Adriamycin
Determination of adriamycin (doxorubicin) and related fluorescent compounds in rat lymph and gall by HPLC (212) 323

Adsorbents
see also sorbents

Coordination polymers as adsorbents and stationary phases in GC. VII. Alkaline earth metal di-*n*-hexylphosphinates: specific interactions (212) 29

Use of $F(ab')_2$ antibody fragments in the synthesis of immunoadsorbents for preparing monospecific antigen (212) 179

Immobilization of residues on agarose gels: effects on protein adsorption isotherms and chromatographic parameters (215) 73

Application of carrageenan beads for chromatographic purification of proteins (215) 93

Hydrophobic chromatography on homologous series of alkylagaroses. A comparison of charged and electrically neutral column materials (215) 211

Quantitative ether cleavage of ligands in hydrophobic agaroses: precise determination of the degree of substitution (215) 235

High-performance liquid affinity chromatography of proteins on Cibacron Blue F3G-A bonded silica (215) 303

High-performance liquid affinity chromatography of enzymes on silica-immobilised triazine dyes (216) 175

HPLC of organic acids on bare silica (218) 285

Application of 1,1'-carbonyldiimidazole-activated matrices for the purification of proteins. III. The use of 1,1'-carbonyldiimidazole-activated agaroses in the biospecific affinity chromatographic isolation of serum antibodies (218) 509

Effect of pH on the GC behaviour of silica gel (219) 101

Concentration and isolation of organic acids on graphitized carbon black (219) 263

Investigation of the activation of cross-linked agarose with carbonylating reagents and the preparation of matrices for affinity chromatography purifications (219) 353

Investigation of the activation of various insoluble polysaccharides with 1,1'-carbonyldiimidazole and of the properties of the activated matrices (219) 361

Adsorption effects
GLSC: effects of adsorption and partitioning (213) 203

Adsorption parameters
Correlation of Hammet substituent constants with reversed-phase "ion-pair" LC adsorption parameters of charged organic compounds (217) 421

Affinity chromatography
Dynamics of dissemination in the case of affinity chromatography (Review) (220) 85

Affinity electrophoresis
Theory of affinity electrophoresis. Evaluation of the effects of protein multivalency, immobilized ligand heterogeneity and microdistribution and determination of effective concentration of immobilized ligand (216) 43

Aflatoxins
Rapid determination of aflatoxin M_1 in dairy products by reversed-phase HPLC (219) 156

Assay method for aflatoxin in milk (219) 162

Agarose gels
Simple gel apparatus for horizontal polyacrylamide and agarose gel electrophoresis (211) 257

Agaroses
Immobilization of residues on agarose gels: effects on protein adsorption isotherms and chromatographic parameters (215) 73

Hydrophobic chromatography on homologous series of alkylagaroses. A comparison of charged and electrically neutral column materials (215) 211

Quantitative ether cleavage of ligands in hydrophobic agaroses: precise determination of the degree of substitution (215) 235

HPLC of macromolecules on agarose and its derivatives (215) 317

Isoelectric focusing in agarose under denaturating conditions (215) 373

Application of 1,1'-carbonyldiimidazole-activated matrices for the purification of proteins. III. The use of 1,1'-carbonyldiimidazole-activated agaroses in the biospecific affinity chromatographic isolation of serum antibodies (218) 509

Investigation of the activation of cross-linked agarose with carbonylating reagents and the preparation of matrices for affinity chromatography purifications (219) 353

Albumin
Adsorption and desorption of proteins in metal chelate affinity chromatography. Purification of albumin (215) 333

Evidence for a concentration-dependent polymerization of a commercial human serum albumin (216) 191

Application of ion-exchange chromatography for the production of human albumin (216) 279

Alcohols
GC separation of some enantiomers on optically active copper(II) complexes (211) 274

Improved high-frequency permittivity detector for the chromatography of alcohols and other polar species (212) 167

Comparison of conventional partitioning systems used for studying the hydrophobicity of polar organic compounds (216) 103

Alditol acetates
High-resolution GC separation of alditol acetates on fused-silica wall-coated open-tubular columns (213) 142

Alditols
Separation of monosaccharides as trimethylsilylated alditols on fused-silica capillary columns (213) 146

Alkaline earth metal complexes
Coordination polymers as adsorbents and stationary phases in GC. VII. Alkaline earth metal di-*n*-hexylphosphinates: specific interactions (212) 29

Alkaloids
Rapid method for the analysis of tobacco nicotine alkaloids (211) 111

Determination of aconitine alkaloids by HPLC (211) 123

Application of statistical optimization methods to the separation of morphine, codeine, noscapine and papaverine in reversed-phase ion-pair chromatography (211) 201

Ion-exchange and straight-phase partition HPLC of the opiates on silica (212) 229

HPLC analysis of pilocarpine hydrochloride, isopilocarpine, pilocarpic acid and isopilocarpic acid in eye-drop preparations (212) 331

Fluorescence and UV detection of opiates separated by reversed-phase HPLC (213) 337

HPLC of *Vinca rosea* alkaloids and the correlation of plate height and molecular weight (214) 95

Preparative ion-pair HPLC and GC of pyrrolizidine alkaloids from comfrey (214) 257

Assay of the combined formulation of ergometrine and oxytocin by HPLC (214) 307

HPLC determination of pergolide and its metabolite, pergolide sulfoxide, in microbial extracts (219) 317

Analysis of potato glycoalkaloids with radially compressed HPLC cartridges and ethanolamine in the mobile phase (219) 403

Alkanals
Relationship between GC retention indices of homologous compounds and the number of carbon atoms and temperature (216) 326

Alkanes
High-back-pressure LC. I. Development of micro-HPLC using liquefied alkanes as the mobile phase (216) 153

Alkanols
Relationship between GC retention indices of homologous compounds and the number of carbon atoms and temperature (216) 326

Alkenes
High-resolution GC with liquid crystal glass capillaries. III. Separation of hydrocarbon isomers on cholesteric mesophase (in German) (219) 225

Alkylthiolation for the determination of double-bond positions in linear alkenes (219) 379

Alkylagaroses
Hydrophobic chromatography on homologous series of alkylagaroses. A comparison of charged and electrically neutral column materials (215) 211

Alkylation
Alkylation with alkyl halides as a derivatization method for the GC determination of acidic pharmaceuticals (Review) (220) 275

Alkylbenzenes
High-resolution reversed-phase LC utilizing microbore column concatenation (216) 9

HPLC of alkylbenzenes on silica (219) 53

High-resolution GC with liquid crystal glass capillaries. III. Separation of hydrocarbon isomers on cholesteric mesophase (in German) (219) 225

Alkyl-bonded silica
Stationary phase characterization in HPLC. A test for trace metal activity in octadecyl bonded silica gel (217) 399

Alkyl-bonded silicas
Influence of the LC mobile phase on the phase transitions of alkyl-bonded silicas studied by GC (214) 202

Comparison of C_n bonded silica gel TLC plates: conditions for use and separations of some barbiturates (214) 209

Alkylglycidyl ethers
Determination of residual epichlorohydrin in middle cut alkylglycidyl ethers by headspace GC (211) 398

Alkylphenols
HPLC determination of 4-nitroso- and 4-nitrophenols in the presence of phenol and alkylphenols (216) 361

Alkyl phosphates
Studies on the GC behaviour of organophosphorus esters (219) 297

Alkylpyridinium salts
GLC determination of N-alkylpyridinium salts (213) 265

Alkylsilane-bonded silica
Comparison of short and ultrashort-chain alkylsilane-bonded silicas for the HPLC of proteins by hydrophobic interaction methods (218) 569

Alkyl sulphates
Determination of carbon chain distribution in alkyl sulfates by *in situ* hydrolysis-GC (219) 117

Amine antidegradants
Antioxidants and stabilizers. LXXXVI. Use of chromatography in the investigation of the mechanism of action of amine antidegradants (211) 299

Amines
Glass capillary columns in the GC separation of aromatic amines. II. Application to samples from workplace atmospheres using nitrogen-selective detection (211) 103

Straight-phase ion-pair chromatography of zimelidine and similar divalent amines. II. The chromatographic system (211) 181

GC separation of some enantiomers on optically active copper(II) complexes (211) 274

Detection of benzidines on thin-layer chromatograms with fluorescamine (211) 304

Stability of the *o*-phthalaldehyde-histamine complex (212) 23

HPLC determination of biogenic amines. I. Use of aqueous acidic mobile phases with silica columns (212) 153

Preparation and evaluation of N-(2-amino-ethyl)-γ-aminopropyltrimethoxysilane-treated silica columns for the HPLC analysis of some aromatic amines (212) 251

Sensitive method for the HPLC determination of amphetamine in urinary extracts (213) 166

Resolution of enantiomers of norepinephrine and epinephrine by reversed-phase HPLC (213) 327

GC determination of some biogenic amines as their pentafluorobenzoyl derivatives in the picogram range and its applicability to biological materials (213) 413

HPLC determination of biogenic amines. II. Comparison of detection methods (213) 429

Separation of amines, guanidines and hydroxy-cinnamic acid amides by ion-exchange chromatography (214) 263

Coupling of proteins and other amines to Sepharose by bromine oxidation and reductive amination (215) 45

Comparison of conventional partitioning systems used for studying the hydrophobicity of polar organic compounds (216) 103

Advantages of a new ion-exchange resin for the LC of biogenic amines: a methacrylic acid polymer cross-linked with butanediol-diacrylate (216) 137

New results in the GC separation of enantiomers of hydroxy acids and carbohydrates (217) 71

Comparison of column packings for trace analysis of free amines by GLC (217) 177

Plasma catecholamines in hypertension and pheochromocytoma determined using ion-pair reversed-phase chromatography with amperometric detection. Investigation of the separation mechanism and clinical methodology (217) 523

Fluorescence detection of chloroanilines in LC using a post-column reaction with fluorescamine. Comparison of reactor types and mixing tees (218) 3

Fluorescence derivatization of tertiary amines with 2-naphthyl chloroformate (218) 51

Study on the retention of amines in reversed-phase ion-pair chromatography on bonded phases (218) 327

Group-contribution approach to the behaviour of 2-phenylethylamines in reversed-phase HPLC (218) 639

Two-phase derivatization of amitriptyline and structurally related tertiary amines for GC with electron-capture detection (219) 373

Amino acids

Optimization of operating parameters in over-pressured TLC (211) 45

LC of cephalosporin C and α-amino acid mixtures on polyfunctional polystyrene resins (211) 79

Improved single-column procedure for the amino acid analysis of collagen-like proteins (211) 175

GC separation of some enantiomers on optically active copper(II) complexes (211) 274

Determination of acetyl and formyl groups by GLC (211) 347

Parameterization of hydrophobic properties of aqueous polymeric biphasic systems and water-organic solvent systems (212) 13

Use of a ternary gradient for the separation of phenylthiohydantoin-amino acids, including the methyl esters of aspartic and glutamic acids by HPLC (213) 91

HPLC of diphenylindenonylthiohydantoin derivatives of amino acids (213) 99

Distinctive test with copper(II)-ninhydrin reagent for small α-peptides separated PC (213) 307

Separation of Dns-amino acids using reversed-phase HPLC: a sensitive method for determining N-termini of peptides and proteins (213) 501

PC separation of phosphate esters, tricarboxylic cycle acids and amino acids in extracts from malaria parasites (214) 111

Efficient preparative counter-current chromatography with a coil planet centrifuge (214) 122

Copolyamino acid fractionation and protobiochemistry (215) 115

Determination of the optical purity of amino acids by HPLC. Modification of the Manning and Moore procedure (215) 279

Partition coefficients of amino acids and hydrophobic parameters π of their side-chains as measured by TLC (216) 79

Behaviour of biomolecules in water-organic solvent-inorganic salt two-phase ternary systems (216) 346

Enantiomeric resolution of amino acids by high-performance ligand-exchange chromatography using a chemically modified hydrophilic porous polymer gel (216) 406

Ligand-exchange chromatography of amino acid racemates on Separon gels containing L-

proline or L-hydroxyproline groupings (217) 539

Ligand-exchange chromatography of racemates. XV. Resolution of α-amino acids on reversed-phase silica gels coated with N-decyl-L-histidine (218) 547

Ligand-exchange chromatography of racemates. XVI. Microbore column chromatography of amino acid racemates using N,N,N',N'-tetramethyl-(*R*)-propane-diamine-1,2-copper(II) complexes as chiral additives to the eluent (218) 559

Histamine analysis on a single column amino acid analyzer (219) 343

Amino alcohols
GC separation of some enantiomers on optically active copper(II) complexes (211) 274

New results in the GC separation of enantiomers of hydroxy acids and carbohydrates (217) 71

Aminoanthraquinones
The role of hydrogen bonds in chromatography. I. Amino- and hydroxyanthraquinones (212) 139

Amino compounds
LC with crown ether-containing mobile phases. I. Capacity factors of amino compounds on a hydrophobic stationary phase (211) 1

Aminocyclohexane
Conversions in catalytic deamination calculated by stopped-flow GC (214) 171

Aminoglycoside antibiotics
Determination of streptomycin sulfate and dihydrostreptomycin sulfate by HPLC (219) 89

Amitriptyline
Two-phase derivatization of amitriptyline and structurally related tertiary amines for GC with electron-capture detection (219) 373

cAMP
Separation of cAMP from adenine nucleotides and nucleosides by electrophoresis on cellulose acetate membranes (211) 409

Amphetamine
Sensitive method for the HPLC determination of amphetamine in urinary extracts (213) 166

Ampholine-free systems
Computer simulation and experimental validation of isoelectric focusing in ampholine-free systems (211) 313

Ampicilline
Mechanism of zwitterion-pair chromatography. II. Ampicilline, lysergic acid, tryptophan and other solutes (218) 355

Analgesics
Large-volume injection of samples dissolved in a non-eluting solvent. Application to the determination of antipyrine using normal-phase HPLC (217) 509

Rapid automated determination of D-penicillamine in plasma and urine by ion-exchange HPLC with electrochemical detection using a gold electrode (218) 615

Application of the combined redox-complexation reaction to the detection of antipyrine and its derivatives in TLC (219) 189

Androstenedione
HPLC of progesterone and its metabolites (219) 152

Angiotensins
TLC of closely related polypeptides on silanized silica gel (211) 29

Anilines
Study of retention behaviour of primary, secondary and tertiary anilines in normal- and reversed-phase LC (214) 291

Anions
Methylation of inorganic anions for GC determination (211) 361

Anthocyanins
Separation of anthocyanin chalcones by HPLC (214) 222

Anthracycline antibiotics
Determination of adriamycin (doxorubicin) and related fluorescent compounds in rat lymph and gall by HPLC (212) 323

Anthraquinones
The role of hydrogen bonds in chromatography. I. Amino- and hydroxyanthraquinones (212) 139

Antibiotics
LC of cephalosporin C and α-amino acid mixtures on polyfunctional polystyrene resins (211) 79

Determination of the relative amounts of the B and C components of neomycin by ion-exclusion chromatography using refractometric detection (211) 223

Determination of adriamycin (doxorubicin) and related fluorescent compounds in rat lymph and gall by HPLC (212) 323

Quantitative analysis of Cephamycin C in fermentation broths by means of thin-layer spectrodensitometry (213) 129

Preparative-scale separation of polymyxins with an analytical HPLC system by using displacement chromatography (215) 295

Identification and determination of the qualitative composition of nystatin using TLC and HPLC (216) 367

Isolation and characterization of three new polymyxins in polymyxins B and E by HPLC (218) 653

Determination of streptomycin sulfate and dihydrostreptomycin sulfate by HPLC (219) 89

Antibodies
Use of $F(ab')_2$ antibody fragments in the synthesis of immunoadsorbents for preparing monospecific antigen (212) 179

Polyacrylhydrazido-agarose: preparation via periodate oxidation and use for enzyme immobilization and affinity chromatography (215) 55

Isolation of hormonal proteins and antibodies by affinity chromatography (215) 143

Purification of specific antibody to α-foetoprotein and its immunological effect on cancer cells (215) 195

Application of 1,1'-carbonyldiimidazole-activated matrices for the purification of proteins. III. The use of 1,1'-carbonyl-diimidazole-activated agaroses in the biospecific affinity chromatographic isolation of serum antibodies (218) 509

Anticholinergic drugs
Determination of aprophen in biological samples by normal-phase HPLC (219) 171

Anticoccidial drugs
Analysis of the anti-coccidial drug, halofuginone, in chicken tissue and chicken feed using HPLC (212) 347

Antidepressant drugs
GC determination of noxyptyline in substance, tablets and in biological material (219) 291

Two-phase derivatization of amitriptyline and structurally related tertiary amines for GC with electron-capture detection (219) 373

Antigens
Use of $F(ab')_2$ antibody fragments in the synthesis of immunoadsorbents for preparing monospecific antigen (212) 179

Antihistamine drugs
Separation of the decomposition products of dimetindene on Sephadex G-10 and LH-20 (in French) (213) 515

Fluorescence derivatization of tertiary amines with 2-naphthyl chloroformate (213) 51

Antihypertensive drugs
HPLC determination of sodium nitroprusside (212) 339

Rapid HPLC method for the measurement of verapamil and norverapamil in blood plasma or serum (218) 621

Anti-inflammatory drugs
Simultaneous determination of diclofenac sodium and its hydroxy metabolites by capillary column GC with electron-capture detection (217) 263

Separation of acidic drugs in the µg/ml range in untreated blood plasma by direct injection on LC columns (218) 671

Antilipemic drugs
Therapeutic drug assays with GLC and optical detection (217) 247

Antimitotic drugs
HPLC determination of Nocodazole in liposomes and L 1210 cells (213) 170

Antimony tartrate
Chromatographic study of optical resolution. VII. Directional ion-association model for the optical resolution of *cis*-$[Co(O)_2(N)_4]^+$ type of complexes by the antimony *d*-tartrate ion (213) 323

Antineoplastic drugs
Determination of adriamycin (doxorubicin) and related fluorescent compounds in rat lymph and gall by HPLC (212) 323

HPLC of *cis*-dichlorodiammineplatinum(II) using chemically-bonded and solvent-generated ion exchangers (217) 405

HPLC of inorganic platinum(II) complexes using solvent-generated anion exchangers. II. The effect of electrolytes on solute retention (219) 235

Use of periodate and methylamine for the quantitation of intracellular 5-fluoro-2'-deoxy-uridine-5'-monophosphate by HPLC (219) 273

HPLC determination of N,N-dimethylcolchiceinamide and its metabolites, N-methylcolchiceinamide and colchiceinamide, in microbial culture (219) 321

Antioxidants
Antioxidants and stabilizers. LXXXVI. Use of chromatography in the investigation of the mechanism of action of amine antidegradants (211) 299

Prediction of retention behaviour of some phenolic antioxidants (213) 363

Determination of 2,6-di-(*tert.*-butyl)-4-methylphenol in new and aged transformer oils using HPLC (in French) (213) 481

Analysis of 2-hydroxybenzophenone and 2'-hydroxyphenylbenzotriazole UV stabilizers by HPLC (214) 237

Antioxidants and stabilizers. LXXXVII. The chromatographic behaviour of transformation products of an antioxidant, N,N'-diphenyl-1,4-phenylenediamine (216) 251

Antipyrine
Large-volume injection of samples dissolved in a non-eluting solvent. Application to the determination of antipyrine using normal-phase HPLC (217) 509

Application of the combined redox-complexation reaction to the detection of antipyrine and its derivatives in TLC (219) 189

Antirheumatic drugs
Rapid automated determination of D-penicillamine in plasma and urine by ion-exchange HPLC with electrochemical detection using a gold electrode (218) 615

Apolipoproteins
Separation of apolipoproteins A-I and A-II by ion-paired reversed-phase HPLC (216) 285

Aprophen
Determination of aprophen in biological samples by normal-phase HPLC (219) 171

Aqueous solubilities
Estimation of aqueous solubilities of organic non-electrolytes using LC retention data (218) 409

Aromatic acids
Optimisation in chromatography. Theory and application to the separation of aromatic acids in reversed-phase LC (213) 357

Aromatic compounds
Chromatographic behaviour of aromatic compounds on non-polar phases with adsorbed quaternary ammonium salts (212) 1

Aromatic hydrocarbons
Prediction of retention indices of aromatic hydrocarbons on phases of different polarity (216) 321

Aryl hydrocarbon hydroxylase
A fast and simple HPLC assay for aryl hydrocarbon hydroxylase (219) 411

Aryloxyethoxyethyl benzyl dimethyl ammonium salts
Speciation of aryloxyethoxyethyl benzyl dimethyl ammonium salts by glass capillary GC and HPLC (213) 463

Ascorbic acid
Separation of ascorbic and isoascorbic acids by ion-pair reversed-phase chromatography (in French) (219) 338

Aspartic acid
Use of a ternary gradient for the separation of phenylthiohydantoin-amino acids, including the methyl esters of aspartic and glutamic acids by HPLC (213) 91

Automatization
Automation of a complex HPLC system. Procedures and hardware for a vitamin B_6 model system (216) 338

Continuous-flow scanning of selected HPLC peak components by microprocessor control. Application to analysis of extracts from human lymphocytes (219) 255

Aza-arenes
Polarity of a poly(oxy aryl sulfonyl arylene) used as stationary phase in GLC (in French) (219) 213

Azinphos methyl
HPLC determination of azinphos methyl and azinphos methyl oxon in fruits and vegetables (214) 140

B

Barbiturates
Comparison of C_n bonded silica gel TLC plates: conditions for use and separations of some barbiturates (214) 209

Determination of 5-ethyl-5-(1-ethylpropyl)-barbituric acid in pentobarbital by HPLC (219) 431

Analysis of barbiturates by GC (Review) (220) 253

BASF
The role of chromatography in BASF. Chromatographic techniques employed in BASF for investigatory studies and for problem solving (Review) (220) 95

Beer bitter acids
Stationary phase characterization in HPLC. A test for trace metal activity in octadecyl bonded silica gel (217) 399

Benzene
Comparison of the distribution constants of benzene in different chromatographic sorbent-gas systems, determined by direct measurement of sorption equilibria and calculated from GC retention data (217) 167

Benzene derivatives
Behaviour of stationary phases. I. Linear relationships between retention data of benzene derivatives (in French) (213) 223

Benzidines
Detection of benzidines on thin-layer chromatograms with fluorescamine (211) 304

Benzoates
Chromatographic behaviour of aromatic compounds on non-polar phases with adsorbed quaternary ammonium salts (212) 1

Benzoic acids
Correlation of Hammet substituent constants with reversed-phase "ion-pair" LC adsorption parameters of charged organic compounds (217) 421

Benzopyrene
A fast and simple HPLC assay for aryl hydrocarbon hydroxylase (219) 411

Benzyl diisobutylphenoxyethoxyethyl dimethyl ammonium chloride
Speciation of aryloxyethoxyethyl benzyl dimethyl ammonium salts by glass capillary GC and HPLC (213) 463

Bergapten
TLC assay of photoactive compounds (furocoumarins) using the fungus *Penicillium expansum* as a test organism (214) 349

Beta-adrenergic receptor blocking drugs
Separation of nadolol diastereoisomers by reversed-phase HPLC (216) 374

Bile pigments
HPLC of bile pigments: separation and characterization of the urobilinoids (218) 647

Bioaffinity chromatography
Development of modern bioaffinity chromatography (a review) (218) 241

Biphenyls
GC behaviour of mono- and dihydroxybiphenyls on various silicone phases (211) 150

Blood coagulation system
Purification and studies of components of the haemostatic system by affinity chromatography (215) 153

Borane anions
Reversed-phase ion-pair LC separations of some heteroborane anions (212) 283

Boronate resin
Separation of plant polyphenolics by chromatography on a boronate resin (216) 261

Bromide
Methylation of inorganic anions for GC determination (211) 361

Brushite
Immobilization of enzymes on columns of brushite (215) 25

Buffer-focusing
Buffer-focusing chromatography using multi-component electrolyte elution systems (218) 483

Butene
Direct GC determination of the products of catalytic air oxidation of *n*-butene-1 to maleic anhydride in the gaseous reaction mixture (214) 217

Butenes
Kinetics of *n*-butene isomerization studied by stopped-flow GC (213) 15

(Butylcyclohexyl)methylpentanone
Analytical and preparative separation of the *cis*- and *trans*-isomers of 4-(4'-*tert.*-butylcyclohexyl)-4-methylpentan-2-one by reversed-phase HPLC (219) 183

C

Calibration methods
Non-linear calibration by cubic splines in gel permeation chromatography (216) 23

Limits of the internal standard technique in chromatography (218) 73

Calorimetric detection
Thermal lens calorimetry. Application to chromatographic detection (218) 15

Camphorates
Manganese(II)-bis(3-heptafluorobutyryl-1*R*-camphorate): a versatile agent for the resolution of racemic cyclic ethers by complexation GC (217) 51

Cancer cells
Classification of human cancer cells by means of capillary GC and pattern recognition analysis (217) 231

Capillary column injector
Properties of a capillary column injector with a controlled rate of injection (217) 99

Capillary columns
Capillary columns with immobilized stationary phases. I. A new simple preparation procedure (211) 243

High-performance micro packed flexible columns in LC (213) 25

Measurement of retention data on open tubular columns coupled in series (213) 189

Capillary columns with immobilized stationary phases. II. Practical advantages and details of procedures (213) 211

Contribution to the theory of open-tubular (capillary) columns in GLC (213) 371

Open-tubular microcapillary LC with 20-μm I.D. columns (214) 283

Separation of complex long chain fatty acid mixtures by high-performance glass capillary GC (216) 115

Role of surface groups in affecting the chromatographic performance of certain types of fused-silica glass capillary columns. I (217) 3

Analysis of polar compounds on polyethylene glycol glass capillary columns. The influence of KF and Na_3PO_4 on the change in partial molar free energy (217) 183

Microcapillary LC in open tubular columns with diameters of 10-50 μm. Potential application to chemical ionization MS detection (218) 137

Flow characteristics and technology of capillary columns with inner diameters less than 15 μm in LC (218) 167

Study of the performance of cation-exchange columns in open-tubular microcapillary LC (218) 189

Testing capillary GC columns (219) 13

Caramel colour
Quantitative ion-pair extraction of 4(5)-methylimidazole from caramel colour and its determination by reversed-phase ion-pair LC (211) 213

Carbamates
HPLC determination of carbaryl and 1-naphthol at residue levels in various water sources by direct injection and trace enrichment (211) 135

GC-MS method for the quantitative analysis of carbofuran in water (212) 366

Carbamoyl chlorides
Determination of distearylcarbamoyl chloride by HPLC (211) 369

Carbaryl
HPLC determination of carbaryl and 1-naphthol at residue levels in various water sources by direct injection and trace enrichment (211) 135

Carbazoles
The chemistry of carbazoles. IX. Substituent effect in the GLC of methylcarbazoles (211) 388

Carbofuran
GC-MS method for the quantitative analysis of carbofuran in water (212) 366

Carbohydrates
see also sugars and saccharides

Improved method for the isolation of 2'-fucosyllactose from human milk (211) 170

Separation of pinitol and some other cyclitols by HPLC (211) 295

Determination of acetyl and formyl groups by GLC (211) 347

High-resolution GC separation of alditol acetates on fused-silica wall-coated open-tubular columns (213) 142

Separation of monosaccharides as trimethylsilylated alditols on fused-silica capillary columns (213) 146

Rapid separation of malto-, xylo- and cello-oligosaccharides (DP 2-9) on cation-exchange resin using water as eluent (213) 151

Affinity column for separation of pyruvylated and non-pyruvylated polysaccharides (213) 301

Consideration of the nature of the lectin-carbohydrate interaction (215) 351

GLC separation and determination of the components of maltitol syrups (216) 219

New results in the GC separation of enantiomers of hydroxy acids and carbohydrates (217) 71

Separation and determination of phenyl isocyanate-derivatized carbohydrates and sugar alcohols by HPLC with UV detection (218) 65

Carbon black
Concentration and isolation of organic acids on graphitized carbon black (219) 263

Carbon dioxide
Characterization of Chromosorb porous polymer bead columns by GC retention values of light hydrocarbons and carbon dioxide (212) 261

Carbon disulphide
Dimethylformamide and carbon disulphide desorption efficiencies for organic vapours on gas-sampling charcoal tube analyses with a GC backflush technique (217) 317

Carbon paste electrodes
Carbon pastes for voltammetric detectors in HPLC (213) 41

Carboxylic acids
GLC analyses. II. Glass capillary GC of methyl monochloro esters of aliphatic C_2-C_{18} *n*-carboxylic acids (211) 267

PC separation of phosphate esters, tricarboxylic cycle acids and amino acids in extracts from malaria parasites (214) 111

Alternate procedure for the preparation and separation of benzyl derivatives of organic acids (214) 131

Isotachophoretic analysis of organic acids produced by fermentation from *n*-alkanes (216) 313

Analysis of the oxocarboxylic acid fraction in serum and urine as O-methyloximes by thermionic specific detection (217) 255

Fluorescence labelling in trace analysis of biological samples. Simultaneous determination of free fatty acids and related carboxylic compounds (217) 491

HPLC of organic acids on bare silica (218) 285

Separation of proton-donating solutes by LC with a strong proton acceptor, tri-*n*-octylphosphine oxide, in the liquid stationary phase (218) 455

Separation and determination of some organic acids and their sodium salts by HPLC (218) 695

Concentration and isolation of organic acids on graphitized carbon black (219) 263

GLC analyses. III. Glass capillary GC of chloromethyl monochloro esters of aliphatic C_3-C_{12} *n*-carboxylic acids (219) 306

Carboxylic proteinases
Hydroxyalkyl methacrylate gels derivatized with epichlorohydrin as supports for large-scale and high-performance affinity chromatography (215) 165

Carcinogens
Glass capillary columns in the GC separation of aromatic amines. II. Application to samples from workplace atmospheres using nitrogen-selective detection (211) 103

Determination of the carcinogen methylazoxy-methyl-β-D-glucosiduronic acid in rat bile and urine (211) 403

Preparation and evaluation of N-(2-amino-ethyl)-γ-aminopropyltrimethoxysilane-treated silica columns for the HPLC analysis of some aromatic amines (212) 251

Confirming the presence of N-nitrosamines in ambient air and cigarette smoke by converting to and photochemically altering their corresponding N-nitroamines (213) 253

HPLC determination of 4-nitroso- and 4-nitrophenols in the presence of phenol and alkylphenols (216) 361

Cardanol
Long-chain phenols. XXI. Quantitative analysis of the phenolic lipids in technical cashew nut-shell liquid, from *Anacardium occidentale,* by HPLC (213) 287

Carrageenan beads
Application of carrageenan beads for chromatographic purification of proteins (215) 93

Carvacrol
HPLC analysis of carvacrol and thymol in the essential oil of *Thymus capitatus* (219) 332

Catalytic air oxidation
Direct GC determination of the products of catalytic air oxidation of *n*-butene-1 to maleic anhydride in the gaseous reaction mixture (214) 217

Catalytic deamination
Conversions in catalytic deamination calculated by stopped-flow GC (214) 171

Catalytic isomerization
Kinetics of *n*-butene isomerization studied by stopped-flow GC (213) 15

Catecholamines
HPLC determination of biogenic amines. I. Use of aqueous acidic mobile phases with silica columns (212) 153

Resolution of enantiomers of norepinephrine and epinephrine by reversed-phase HPLC (213) 327

GC determination of some biogenic amines as their pentafluorobenzoyl derivatives in the picogram range and its applicability to biological materials (213) 413

HPLC determination of biogenic amines. II. Comparison of detection methods (213) 429

Plasma catecholamines in hypertension and pheochromocytoma determined using ion-pair reversed-phase chromatography with amperometric detection. Investigation of the separation mechanism and clinical methodology (217) 523

Automated measurement of catecholamines in urine, plasma and tissue homogenates by HPLC with fluorometric detection (218) 631

Determination of cross-contamination of the diastereomers ephedrine and pseudoephedrine by HPLC, TLC and carbon-13 nuclear magnetic resonance spectroscopy (219) 81

Catechol-O-methyltransferase
Assay for catechol-O-methyltransferase by HPLC with fluorescence detection (211) 129

Catechols
C_{18} reversed-phase trace enrichment of chlorinated phenols, guaiacols and catechols in water (214) 327

Cations
Migration behaviour of the hydrogen ion and its role in isotachophoresis of cations (217) 209

Cephalosporin
LC of cephalosporin C and α-amino acid mixtures on polyfunctional polystyrene resins (211) 79

Cephamycin C
Quantitative analysis of Cephamycin C in fermentation broths by means of thin-layer spectrodensitometry (213) 129

Ceruloplasmin
Covalent chromatography as a means of isolating thiol peptides from large proteins. Application to human ceruloplasmin (215) 341

Charge-transfer acceptors
Non-ionic mobile phase dopants. I. Chiral charge-transfer acceptors and helicene resolution (216) 333

Chelating gels
Adsorption and desorption of proteins in metal chelate affinity chromatography. Purification of albumin (215) 333

Chemotherapeutic drugs
R_M and log *P* values of 5-nitroimidazoles (216) 93

Chiral additives
Manganese(II)-bis(3-heptafluorobutyryl-1*R*-camphorate): a versatile agent for the resolution of racemic cyclic ethers by complexation GC (217) 51

Ligand-exchange chromatography of racemates. XVI. Microbore column chromatography of amino acid racemates using N,N,N',N'-tetramethyl-(*R*)-propane-diamine-1,2-copper(II) complexes as chiral additives to the eluent (218) 559

Chiral stationary phases
GC separation of some enantiomers on optically active copper(II) complexes (211) 274

N-(1*R*, 3*R*)-*trans*-Chrysanthemoyl (*R*)-1-(α-naphthyl)ethylamine as a stationary phase for the separation of optical isomers by GC (213) 137

New results in the GC separation of enantiomers of hydroxy acids and carbohydrates (217) 71

Chlorinated phenols
Determination of chlorinated phenols by HPLC (213) 83

Chloroanilines
GC data for chlorobenzenes, chloroanilines and N-sulphinylanilines. Relationship between molecular structure and retention based on retention index increments calculated by computer (214) 47

Fluorescence detection of chloroanilines in LC using a post-column reaction with fluorescamine. Comparison of reactor types and mixing tees (218) 3

Chlorobenzenes
GC data for chlorobenzenes, chloroanilines and N-sulphinylanilines. Relationship between molecular structure and retention based on retention index increments calculated by computer (214) 47

Chloroepoxypropane
Determination of residual epichlorohydrin in middle cut alkylglycidyl ethers by headspace GC (211) 398

Chlorophacinone
Determination of chlorophacinone in formulations by reversed-phase ion-pair chromatography (214) 335

Chlorophenols
C_{18} reversed-phase trace enrichment of chlorinated phenols, guaiacols and catechols in water (214) 327

Determination of pentachlorophenol and related compounds in animal materials by HPLC and GC (216) 229

Chlorophenoxyisobutyric acid
Therapeutic drug assays with GLC and optical detection (217) 247

Chlorophyll protein
Separation of membrane components by partition in detergent-containing polymer phase systems. Isolation of the light harvesting chlorophyll *a/b* protein (215) 131

Chlorophylls
Rapid, quantitative separation of chlorophylls and their degradation products by HPLC (213) 349

Chloropropionyl chlorides
GLC analyses of chlorination products of propionyl chloride (213) 63

Cholesterol
Capillary column GLC analysis of cholesterol derivatives. Application to the autoxidation products of cholesterol (211) 392

Cholesteryl esters
Separation of [^{3}H]cholesteryl esters by reversed-phase HPLC (213) 105

Choline
Separation of choline and acetylcholine by cation-exchange chromatography (214) 148

Chondroitin sulphates
Separation of reduced disaccharides derived from glycosaminoglycans by HPLC (212) 65

Chromogenic reagents
Sensitive, enzyme-catalyzed, chromogenic reagent for hydrogen peroxide and other peroxygen compounds on TLC plates (216) 413

Chromosorb
Characterization of Chromosorb porous polymer bead columns by GC retention values of light hydrocarbons and carbon dioxide (212) 261

Chymotrypsin
Affinity chromatography of chymotrypsin (E.C. 3.4.21.1) on the potato trypsin inhibitor bound to bead cellulose by the benzoquinone method (213) 352

Comparative study of native and chemically modified chymotrypsin as monomers, soluble polymers and membranes (215) 87

Cigarette smoke
Confirming the presence of N-nitrosamines in ambient air and cigarette smoke by converting to and photochemically altering their corresponding N-nitroamines (213) 253

Determination of α-dicarbonyl compounds in cigarette smoke (217) 197

Cinnamic acids
Correlation of Hammet substituent constants with reversed-phase "ion-pair" LC adsorption parameters of charged organic compounds (217) 421

Cinnamoyl cocaine
Quantitation of cocaine in a variety of matrices by HPLC (214) 243

Cisplatin
HPLC of *cis*-dichlorodiammineplatinum(II) using chemically-bonded and solvent-generated ion exchangers (217) 405

HPLC of inorganic platinum(II) complexes using solvent-generated anion exchangers. II. The effect of electrolytes on solute retention (219) 235

Citric acid
Isotachophoretic analysis of organic acids produced by fermentation from *n*-alkanes (216) 313

Clathrates
Selectivity and efficiency of separation of isomers of organic acids by clathrate chromatography (218) 465

Clofibric acid
Therapeutic drug assays with GLC and optical detection (217) 247

CM-Trisacryl M
Physico-chemical and chromatographic properties of new ion exchangers. I. CM-Trisacryl M (213) 389

Coal extracts
Chromatographic fingerprinting of coal extracts (217) 19

Coal liquids
HPLC studies of coal liquids by laser-based detectors (211) 95

Cobalt complexes
High-performance paper electrophoresis. II. Comparison of separations obtained by high-voltage paper electrophoresis and high-performance paper electrophoresis (212) 374

Chromatographic study of optical resolution. VII. Directional ion-association model for the optical resolution of *cis*-$[Co(O)_2(N)_4]^+$ type of complexes by the antimony *d*-tartrate ion (213) 323

Chromatographic study of optical resolution. VIII. Theoretical study of the chromatographic behaviour of the enantiomers of racemic complex cations on a cation-exchange column (219) 29

Cocaine
Quantitation of cocaine in a variety of matrices by HPLC (214) 243

Codeine
Application of statistical optimization methods to the separation of morphine, codeine, noscapine and papaverine in reversed-phase ion-pair chromatography (211) 201

Coil planet centrifuge
Efficient preparative counter-current chromatography with a coil planet centrifuge (214) 122

Colchiceinamide
HPLC determination of N,N-dimethylcolchiceinamide and its metabolites, N-methylcolchiceinamide and colchiceinamide, in microbial culture (219) 321

Colistin
Isolation and characterization of three new polymyxins in polymyxins B and E by HPLC (218) 653

Collagen membranes
Enzymes immobilized on collagen membranes: a tool for fundamental research and enzyme engineering (215) 65

Column coating
Capillary columns with immobilized stationary phases. I. A new simple preparation procedure (211) 243

Capillary columns with immobilized stationary phases. II. Practical advantages and details of procedures (213) 211

Role of surface groups in affecting the chromatographic performance of certain types of fused-silica glass capillary columns. I (217) 3

Column geometry
Concentration effects in gel permeation chromatography. VII. Viscosity phenomena and column geometry (211) 39

Column performance
Effect of mobile phase pre-heating on HPLC column performance: a new type of column thermostat (213) 243

Study of the performance of cation-exchange columns in open-tubular microcapillary LC (218) 189

Column preparation
Open-tubular microcapillary LC with 20-μm I.D. columns (214) 283

Separation of complex long chain fatty acid mixtures by high-performance glass capillary GC (216) 115

Flow characteristics and technology of capillary columns with inner diameters less than 15 μm in LC (218) 167

Columns
High-performance micro packed flexible columns in LC (213) 25

Measurement of retention data on open tubular columns coupled in series (213) 189

Contribution to the theory of open-tubular (capillary) columns in GLC (213) 371

High-resolution reversed-phase LC utilizing microbore column concatenation (216) 9

Microcapillary LC in open tubular columns with diameters of 10-50 μm. Potential application to chemical ionization MS detection (218) 137

Application of ultra-micro HPLC to trace analysis (218) 199

Recycling LC using microbore columns (219) 1

Testing capillary GC columns (219) 13

Column switching
Liquid column switching extraction and chromatography for programmed flow preparation (217) 387

HPLC column switching techniques for rapid hydrocarbon group-type separations (218) 715

Column thermostat
Effect of mobile phase pre-heating on HPLC column performance: a new type of column thermostat (213) 243

Column treatment
Analysis of polar compounds on polyethylene glycol glass capillary columns. The influence of KF and Na_3PO_4 on the change in partial molar free energy (217) 183

Computer simulation
Computer simulation and experimental validation of isoelectric focusing in ampholine-free systems (211) 313

Computer simulation of the kinetics of ion exchange accompanied by complex formation in ionites (212) 133

Concanavalin A
Application of polyacrylamide gradient gel electrophoresis to the study of multiple components of concanavalin A and related lectins (211) 233

Continuous-flow scanning
Continuous-flow scanning of selected HPLC peak components by microprocessor control. Application to analysis of extracts from human lymphocytes (219) 255

Contraceptives
General method for the assay of oral contraceptives by HPLC (216) 269

Analysis of isomeric ethynylestradiol glucuronides in urine (217) 453

Copolyamino acids
Copolyamino acid fractionation and protobiochemistry (215) 115

Copper-ninhydrin reagent
Distinctive test with copper(II)-ninhydrin reagent for small α-peptides separated PC (213) 307

Corticosteroids
Application of ultra-micro HPLC to trace analysis (218) 199

Cortisol
LC measurement of cortisol in methylene chloride extracts of aqueous solutions (212) 221

HPTLC determination of fluorescence-labelled cortisol (217) 349

Coumarylagmatine
Separation of amines, guanidines and hydroxycinnamic acid amides by ion-exchange chromatography (214) 263

Coumesterol
Separation of genistin, daidzin and their aglucones, and coumesterol by gradient HPLC (211) 166

Crown ethers
LC with crown ether-containing mobile phases. I. Capacity factors of amino compounds on a hydrophobic stationary phase (211) 1

Crude oils
Microextraction and GC analysis of selected petroleum hydrocarbons in water and fish tissue (212) 305

Fractionation of polar organic constituents in environmental samples using the lipophilic dextran gels Sephadex LH-20 and Sephasorb HP Ultrafine. Application to a weathered Ekofisk crude oil (214) 83

TLC analysis of aromatic hydrocarbons in crude oil and in petroleum products as their η^6-arene-η^5-cyclopentadienyliron hexafluorophosphates (219) 349

GC separation of diastereomeric isoprenoids as molecular markers of oil pollution (219) 393

Cyanide
Methylation of inorganic anions for GC determination (211) 361

Cycasin
Determination of the carcinogen methylazoxymethyl-β-D-glucosiduronic acid in rat bile and urine (211) 403

Cyclic AMP
Separation of cAMP from adenine nucleotides and nucleosides by electrophoresis on cellulose acetate membranes (211) 409

Cyclic ethers
Manganese(II)-bis(3-heptafluorobutyryl-1*R*-camphorate): a versatile agent for the resolution of racemic cyclic ethers by complexation GC (217) 51

Cyclic octametaphosphate
Chromatographic study of the acidic hydrolysis of cyclic octametaphosphate (211) 87

Cyclic trimetaphosphate
Study of the acidic hydrolysis of cyclic trimetaphosphate by LC (219) 385

Cyclitols
Separation of pinitol and some other cyclitols by HPLC (211) 295

Cyclodextrin resins
Sorption behaviour of several organic compounds on 6-deoxycyclodextrin polyurethane resins (219) 108

Cytochromes
Separation of cytochromes *c* by reversed-phase HPLC (212) 295

Cytokinins
Combined HPLC radioimmunoassay for cytokinins (214) 101

HPLC analysis of permethylated cytokinins (219) 167

D

Daidzin
Separation of genistin, daidzin and their aglucones, and coumesterol by gradient HPLC (211) 166

Deamination
Conversions in catalytic deamination calculated by stopped-flow GC (214) 171

Dehydrocholesterol
Determination of 7-dehydrocholesterol in rat skin and liver by HPLC (216) 385

Demoxepam
LC of demoxepam and phenothiazines using a post-column photochemical reactor and fluorescence detection (217) 463

Deoxyribonucleic acids
see DNA

Derivatization
Methylation of inorganic anions for GC determination (211) 361

Stability of the *o*-phthalaldehyde-histamine complex (212) 23

Trimethylsilylation reaction of prostaglandin-E methyl ester with various trimethylsilylating reagents (214) 177

Fluorescence derivatization of tertiary amines with 2-naphthyl chloroformate (218) 51

Separation and determination of phenyl isocyanate-derivatized carbohydrates and sugar alcohols by HPLC with UV detection (218) 65

Two-phase derivatization of amitriptyline and structurally related tertiary amines for GC with electron-capture detection (219) 373

Alkylthiolation for the determination of double-bond positions in linear alkenes (219) 379

Analysis of barbiturates by GC (Review) (220) 253

Alkylation with alkyl halides as a derivatization method for the GC determination of acidic pharmaceuticals (Review) (220) 275

Dermorphins
Reversed-phase HPLC of dermorphins, opiate-like peptides from amphibian skins (216) 355

Des-enkephalin γ-endorphin
Preparative purification of the peptide des-enkephalin γ-endorphin. Comparison of HPLC and counter-current chromatography (212) 356

Desorption solvents
Dimethylformamide and carbon disulphide desorption efficiencies for organic vapours on gas-sampling charcoal tube analyses with a GC backflush technique (217) 317

Detectors
HPLC studies of coal liquids by laser-based detectors (211) 95

Use of 214-nm and 229-nm discrete-line sources for the UV absorbance detection of some pesticides separated by HPLC (211) 144

Photometric monitoring of thiols by means of a thiomercurimetric detector (211) 252

Detection of brominated and chlorinated organics by a GC microcoulometric detector. Effect of pyrolysis tube conditions (212) 115

Improved high-frequency permittivity detector for the chromatography of alcohols and other polar species (212) 167

Carbon pastes for voltammetric detectors in HPLC (213) 41

Response enhancement by metal doping of a silicon-selective flame ionization detector for GC (213) 397

Nitrogen-selective detector for HPLC (216) 127

Correlation and prediction of electron-capture response from molecular parameters (217) 151

Thermal lens calorimetry. Application to chromatographic detection (218) 15

Extra-column effects in polarographic *versus* UV detection in HPLC (218) 45

Micro-column HPLC and flame-based detection principles (218) 179

Electrokinetic detection at different points in a narrow-bore glass column in LC (219) 61

Detergents
GLC determination of N-alkylpyridinium salts (213) 265

Diacetoxyscirpenol
HPLC of trichothecenes. I. Detection of T-2 toxin and HT-2 toxin (212) 370

Diamantane oximes
Separation of diamantane-3-oneoxime stereo-isomers by preparative HPLC (219) 179

Diaminotoluenes
Preparation and evaluation of N-(2-aminoethyl)-γ-aminopropyltrimethoxysilane-treated silica columns for the HPLC analysis of some aromatic amines (212) 251

Dibenzodioxins
Determination of chlorinated dibenzo-*p*-dioxin contaminants in 2,4-D products by GC-MS techniques (217) 289

Dibromochloropropane
Quantitative determination of 1,2-dibromo-3-chloropropane in whole rat blood and drinking water by GC (213) 156

Dibutylmethylphenol
Determination of 2,6-di-(*tert.*-butyl)-4-methylphenol in new and aged transformer oils using HPLC (in French) (213) 481

Dicarbonyl compounds
Determination of α-dicarbonyl compounds in cigarette smoke (217) 197

Dichlorodiammineplatinum
HPLC of *cis*-dichlorodiammineplatinum(II) using chemically-bonded and solvent-generated ion exchangers (217) 405

HPLC of inorganic platinum(II) complexes using solvent-generated anion exchangers. II. The effect of electrolytes on solute retention (219) 235

Diclofenac sodium
Simultaneous determination of diclofenac sodium and its hydroxy metabolites by capillary column GC with electron-capture detection (217) 263

Dicyclopentadienes
Influence of intramolecular interactions on the GC separation of epimers of dicyclopentadiene derivatives (in German) (219) 113

Dihexylphosphinates
Coordination polymers as adsorbents and stationary phases in GC. VII. Alkaline earth metal di-*n*-hexylphosphinates: specific interactions (212) 29

Dihydromevinolin
HPLC analysis of derivatized hypocholesteremic agents from fermentation broths (212) 234

Dihydrostreptomyin
Determination of streptomycin sulfate and dihydrostreptomycin sulfate by HPLC (219) 89

Dihydroxybiphenyls
GC behaviour of mono- and dihydroxybiphenyls on various silicone phases (211) 150

Dimethylcolchiceinamide
HPLC determination of N,N-dimethylcolchiceinamide and its metabolites, N-methylcolchiceinamide and colchiceinamide, in microbial culture (219) 321

Dimethylformamide
Dimethylformamide and carbon disulphide desorption efficiencies for organic vapours on gas-sampling charcoal tube analyses with a GC backflush technique (217) 317

Dimethylhydrazine
Separation of hydrazine, monomethylhydrazine, 1,1-dimethylhydrazine and 1,2-dimethylhydrazine by HPLC with electrochemical detection (214) 229

Dimetindene
Separation of the decomposition products of dimetindene on Sephadex G-10 and LH-20 (in French) (213) 515

Diphenhydramine
Fluorescence derivatization of tertiary amines with 2-naphthyl chloroformate (218) 51

Diphenylphenylenediamine
Antioxidants and stabilizers. LXXXVII. The chromatographic behaviour of transformation products of an antioxidant, N,N'-diphenyl-1,4-phenylenediamine (216) 251

Disaccharides
Separation of reduced disaccharides derived from glycosaminoglycans by HPLC (212) 65

Disinfectants
GLC determination of N-alkylpyridinium salts (213) 265

Displacement chromatography
High-performance displacement chromatography (218) 365

Dissemination
Dynamics of dissemination in the case of affinity chromatography (Review) (220) 85

Distearylcarbamoyl chloride
Determination of distearylcarbamoyl chloride by HPLC (211) 369

Distribution constants
Comparison of the distribution constants of benzene in different chromatographic sorbent-gas systems, determined by direct measurement of sorption equilibria and calculated from GC retention data (217) 167

Comparison of the distribution constants of acetone in different chromatographic sorbent-gas systems, determined by direct measurement of sorption equilibria and calculated from GC retention data (219) 37

DNA
Simple gel apparatus for horizontal polyacrylamide and agarose gel electrophoresis (211) 257

Dopamine
GC determination of some biogenic amines as their pentafluorobenzoyl derivatives in the picogram range and its applicability to biological materials (213) 413

HPLC determination of biogenic amines. II. Comparison of detection methods (213) 429

Dopants
Non-ionic mobile phase dopants. I. Chiral charge-transfer acceptors and helicene resolution (216) 333

Doping
Response enhancement by metal doping of a silicon-selective flame ionization detector for GC (213) 397

Doubly peaked chromatograms
Doubly peaked chromatograms (213) 355

Doxorubicin
Determination of adriamycin (doxorubicin) and related fluorescent compounds in rat lymph and gall by HPLC (212) 323

Drugs of abuse
Quantitation of cocaine in a variety of matrices by HPLC (214) 243

E

Ecdysteroids
Reversed-phase TLC for the separation and analysis of ecdysteroids (212) 211

Effective mobility
Concept of the effective mobility of the hydrogen ion and its use in cationic isotachophoresis (219) 21

Electrokinetic detection
Electrokinetic detection at different points in a narrow-bore glass column in LC (219) 61

Electron capture detector
Correlation and prediction of electron-capture response from molecular parameters (217) 151

Enantiomer separation
GC separation of some enantiomers on optically active copper(II) complexes (211) 274

N-(1*R*, 3*R*)-*trans*-Chrysanthemoyl (*R*)-1-(α-naphthyl)ethylamine as a stationary phase for the separation of optical isomers by GC (213) 137

Chromatographic study of optical resolution. VII. Directional ion-association model for the optical resolution of *cis*-$[Co(O)_2(N)_4]^+$ type of complexes by the antimony *d*-tartrate ion (213) 323

Resolution of enantiomers of norepinephrine and epinephrine by reversed-phase HPLC (213) 327

Determination of the optical purity of amino acids by HPLC. Modification of the Manning and Moore procedure (215) 279

Non-ionic mobile phase dopants. I. Chiral charge-transfer acceptors and helicene resolution (216) 333

Separation of nadolol diastereoisomers by reversed-phase HPLC (216) 374

Enantiomeric resolution of amino acids by high-performance ligand-exchange chromatography using a chemically modified hydrophilic porous polymer gel (216) 406

Manganese(II)-bis(3-heptafluorobutyryl-1*R*-camphorate): a versatile agent for the resolution of racemic cyclic ethers by complexation GC (217) 51

New results in the GC separation of enantiomers of hydroxy acids and carbohydrates (217) 71

Ligand-exchange chromatography of amino acid racemates on Separon gels containing L-proline or L-hydroxyproline groupings (217) 539

Ligand-exchange chromatography of racemates. XV. Resolution of α-amino acids on reversed-phase silica gels coated with N-decyl-L-histidine (218) 547

Ligand-exchange chromatography of racemates. XVI. Microbore column chromatography of amino acid racemates using N,N,N',N'-tetramethyl-(*R*)-propane-diamine-1,2-copper(II) complexes as chiral additives to the eluent (218) 559

Chromatographic study of optical resolution. VIII. Theoretical study of the chromatographic behaviour of the enantiomers of racemic complex cations on a cation-exchange column (219) 29

Endorphin
Preparative purification of the peptide des-enkephalin γ-endorphin. Comparison of HPLC and counter-current chromatography (212) 356

Partition chromatography and HPLC of β-lipotropin and synthetic β-endorphin analogues (215) 255

Enthalpy/entropy compensations
Structural effects in enthalpy/entropy compensated and non-compensated behaviour in ion-pair reversed-phase HPLSC (218) 427

Enzyme immobilization
Some carriers for the immobilization of enzymes based on derivatized poly(vinyl alcohol) and on copolymers of methacrylates with different spacer lengths (215) 13

Immobilization of enzymes on columns of brushite (215) 25

Polymeric supports bearing isonitrile functional groups for covalent fixation of biologically active molecules (a review) (215) 31

Coupling of proteins and other amines to Sepharose by bromine oxidation and reductive amination (215) 45

Polyacrylhydrazido-agarose: preparation via periodate oxidation and use for enzyme immobilization and affinity chromatography (215) 55

Enzymes immobilized on collagen membranes: a tool for fundamental research and enzyme engineering (215) 65

Immobilization of residues on agarose gels: effects on protein adsorption isotherms and chromatographic parameters (215) 73

Enzymes
see also proteins

Assay for catechol-O-methyltransferase by HPLC with fluorescence detection (211) 129

Affinity chromatography of chymotrypsin (E.C. 3.4.21.1) on the potato trypsin inhibitor bound to bead cellulose by the benzoquinone method (213) 352

Separation of the tryptic peptides from reduced, alkylated hen egg white lysozyme by HPLC (214) 185

Comparative study of native and chemically modified chymotrypsin as monomers, soluble polymers and membranes (215) 87

Application of carrageenan beads for chromatographic purification of proteins (215) 93

Purification and partial characterization of vanillate hydroxylase (decarboxylating) from *Sporotrichum pulverulentum* (215) 99

Interaction of glucose oxidase with Blue Dextran (215) 121

Hydroxyalkyl methacrylate gels derivatized with epichlorohydrin as supports for large-scale and high-performance affinity chromatography (215) 165

Affinity chromatography of glycosyltransferases (215) 181

High-performance liquid affinity chromatography of proteins on Cibacron Blue F3G-A bonded silica (215) 303

Resolution in high-performance liquid affinity chromatography. Dependence on eluite diffusion into the stationary phase (216) 169

High-performance liquid affinity chromatography of enzymes on silica-immobilised triazine dyes (216) 175

Large-scale purification of staphylococcal lipase by hydrophobic interaction chromatography (216) 295

Affinity chromatography of acetylcholinesterase. The use of Amberlite CG-120 for dissociating the enzyme-inhibitor complex (219) 71

A fast and simple HPLC assay for aryl hydrocarbon hydroxylase (219) 411

Purification of 6-phosphogluconate dehydrogenase from *Acer pseudoplatanus* L. using immobilised Procion Red HE-3B (219) 419

Ephedrine
Determination of cross-contamination of the diastereomers ephedrine and pseudoephedrine by HPLC, TLC and carbon-13 nuclear magnetic resonance spectroscopy (219) 81

Epichlorohydrin
Determination of residual epichlorohydrin in middle cut alkylglycidyl ethers by headspace GC (211) 398

Epinephrine
Resolution of enantiomers of norepinephrine and epinephrine by reversed-phase HPLC (213) 327

Plasma catecholamines in hypertension and pheochromocytoma determined using ion-pair reversed-phase chromatography with amperometric detection. Investigation of the separation mechanism and clinical methodology (217) 523

Ergometrine
Assay of the combined formulation of ergometrine and oxytocin by HPLC (214) 307

Ergot alkaloids
HPLC determination of pergolide and its metabolite, pergolide sulfoxide, in microbial extracts (219) 317

Essential oils
Essential oil composition of *Aframomum korarima* (211) 280

HPLC separation of shallot volatile oil (214) 234

HPLC analysis of carvacrol and thymol in the essential oil of *Thymus capitatus* (219) 332

Esters
Relationship between GC retention indices of homologous compounds and the number of carbon atoms and temperature (216) 326

GLC retention behaviour of hindered aliphatic esters (217) 125

GC of homologous esters. XV. Molecular retention indices of aliphatic esters (219) 45

Use of routine preparative HPLC in the separation of isomers (219) 140

Estrogens
see oestrogens

Ethane
GSC of ethane on activated carbon at 25°C (214) 269

Ethers
Manganese(II)-bis(3-heptafluorobutyryl-1*R*-camphorate): a versatile agent for the resolution of racemic cyclic ethers by complexation GC (217) 51

Ethylenethiourea
LC with UV absorbance and polarographic detection of ethylenethiourea and related sulfur compounds. Application to rat urine analysis (212) 245

Ethyl(ethylpropyl)barbituric acid
Determination of 5-ethyl-5-(1-ethylpropyl)-barbituric acid in pentobarbital by HPLC (219) 431

Ethynylestradiol
Analysis of isomeric ethynylestradiol glucuronides in urine (217) 453

Europium complexes
Applications of selective complexation by a europium(III) coordination polymer sorbent for the pre-fractionation of volatile compounds (217) 275

Extra-column effects
Extra-column effects in polarographic *versus* UV detection in HPLC (218) 45

Extraction procedures
Liquid column switching extraction and chromatography for programmed flow preparation (217) 387

F

Fatty acids
Comparison of conventional partitioning systems used for studying the hydrophobicity of polar organic compounds (216) 103

Separation of complex long chain fatty acid mixtures by high-performance glass capillary GC (216) 115

Fluorescence labelling in trace analysis of biological samples. Simultaneous determination of free fatty acids and related carboxylic compounds (217) 491

Fenvalerate
HPLC of fenvalerate (216) 303

Flame ionization detector
Response enhancement by metal doping of a silicon-selective flame ionization detector for GC (213) 397

Flame photometric detector
Micro-column HPLC and flame-based detection principles (218) 179

Flavones
Role of water in qualitative and quantitative determination of polymethoxylated flavones by straight-phase HPLC: application to orange peel oils (211) 61

Flavonoids
HPLC of flavonoids in barley and hops (218) 683

Flow preparation
Liquid column switching extraction and chromatography for programmed flow preparation (217) 387

Fluorescence-inducing procedures
Fluorescence-inducing procedures for use in HPTLC (217) 341

Fluorescence-inducing reagents
Use of metal salts as fluorescence-inducing reagents in TLC (217) 329

Fluorescence reagents
Fluorescence derivatization of tertiary amines with 2-naphthyl chloroformate (218) 51

Fluorodeoxyuridine
Use of periodate and methylamine for the quantitation of intracellular 5-fluoro-2'-deoxyuridine-5'-monophosphate by HPLC (219) 273

Fluorouracil
Use of periodate and methylamine for the quantitation of intracellular 5-fluoro-2'-deoxyuridine-5'-monophosphate by HPLC (219) 273

Foetoprotein
Purification of specific antibody to α-foetoprotein and its immunological effect on cancer cells (215) 195

Food packings
Analysis of additives in plastics by high-performance size-exclusion chromatography (218) 703

Formyl groups
Determination of acetyl and formyl groups by GLC (211) 347

Fraction collector
Fraction collector controller for collecting LC peaks (212) 105

Fucosyllactose
Improved method for the isolation of 2'-fucosyllactose from human milk (211) 170

Fungicides
Simultaneous determination of trialkyltin homologues in biological materials (216) 209

Furocoumarins
TLC assay of photoactive compounds (furocoumarins) using the fungus *Penicillium expansum* as a test organism (214) 349

Fused-silica columns
High-performance micro packed flexible columns in LC (213) 25

Role of surface groups in affecting the chromatographic performance of certain types of fused-silica glass capillary columns. I (217) 3

G

Gas solubility
Solubilities of gases in H_2O and ^2H_2O (216) 161

Gel apparatus
Simple gel apparatus for horizontal polyacrylamide and agarose gel electrophoresis (211) 257

Genistin
Separation of genistin, daidzin and their aglucones, and coumesterol by gradient HPLC (211) 166

Gibberellins
Adsorption and reversed-phase partition HPLC of gibberellins (213) 507

Gingerdiones
GC and MS studies of ginger constituents. Identification of gingerdiones and new hexahydrocurcumin analogues (212) 75

Ginsenosides
Identification of ginsenosides from *Panax ginseng* in fractions obtained by HPLC by field desorption MS, multiple internal reflection IR spectroscopy and TLC (212) 37

Gluconic acid
Separation and determination of some organic acids and their sodium salts by HPLC (218) 695

Glucose
GLC separation and determination of the components of maltitol syrups (216) 219

Glucose oxidase
Interaction of glucose oxidase with Blue Dextran (215) 121

Glutamic acid
Use of a ternary gradient for the separation of phenylthiohydantoin-amino acids, including the methyl esters of aspartic and glutamic acids by HPLC (213) 91

Glycoalkaloids
Analysis of potato glycoalkaloids with radially compressed HPLC cartridges and ethanolamine in the mobile phase (219) 403

Glycopolypeptides
High-speed gel filtration of glycopolypeptides in 6 *M* guanidine hydrochloride (215) 289

Glycoproteins
Isolation of human haemopexin by bioaffinity chromatography on haeme-Sepharose (214) 317

Analysis of glycoprotein hormones and other medically important proteins by high-performance gel filtration chromatography (218) 581

Glycosaminoglycans
Separation of reduced disaccharides derived from glycosaminoglycans by HPLC (212) 65

Glycosyltransferases
Affinity chromatography of glycosyltransferases (215) 181

Gold complexes
Applications of HPLC in organometallic chemistry: ligand exchange in complexes of ylides (in German) (213) 47

Gonadotropin
Separation of subunits of pike eel gonadotropin by hydrophobic interaction chromatography (215) 229

Gradient elution
Gradient elution in LC. XIV. Theory of ternary gradients in reversed-phase LC (214) 35

Graphical methods
Application of a three-dimensional drawing procedure to the evaluation of series of protein samples after analysis by gel electrophoresis and other methods (214) 126

Graphitized carbon black
Concentration and isolation of organic acids on graphitized carbon black (219) 263

Guaiacols
C_{18} reversed-phase trace enrichment of chlorinated phenols, guaiacols and catechols in water (214) 327

Guanidines
Separation of amines, guanidines and hydroxycinnamic acid amides by ion-exchange chromatography (214) 263

Guayulins
Detection and quantification of guayulins A and B in *Parthenium argentatum* (guayule) and F_1 hybrids by HPLC (213) 345

Guthion
HPLC determination of azinphos methyl and azinphos methyl oxon in fruits and vegetables (214) 140

H

Haeme proteins
Interaction of heme proteins and thyroid hormone. II. Localization of the site on thyroid hormone that binds to hemoglobin (219) 148

Haemoglobin
Electrofocusing of stroma-free haemoglobin and its derivatives in agarose isoelectric focusing gels (211) 308

Characterization of polymerized stroma-free haemoglobin by chromatography, electrophoresis and electrofocusing in gels (213) 183

Isoelectric focusing, gel chromatography and electrophoresis of pyridoxalated and polymerized stroma-free haemoglobin (219) 128

Chromatofocusing of pyridoxalated and polymerized human haemoglobin (219) 436

Haemopexin
Isolation of human haemopexin by bioaffinity chromatography on haeme-Sepharose (214) 317

Haemostatic system
Purification and studies of components of the haemostatic system by affinity chromatography (215) 153

Halofuginone
Analysis of the anti-coccidial drug, halofuginone, in chicken tissue and chicken feed using HPLC (212) 347

Halogenated organics
Detection of brominated and chlorinated organics by a GC microcoulometric detector. Effect of pyrolysis tube conditions (212) 115

Height equivalent to a theoretical plate
The height equivalent to a theoretical plate of retentionless rectangular tubes (216) 1

Temperature dependence of HETP for columns with small amounts of liquid stationary phase (217) 225

Helicenes
Non-ionic mobile phase dopants. I. Chiral charge-transfer acceptors and helicene resolution (216) 333

Heparin
Separation of reduced disaccharides derived from glycosaminoglycans by HPLC (212) 65

Hydrophobic interaction chromatography of mucopolysaccharides. Examination of fundamental conditions for fractionation of heparin on hydrophobic gels (213) 439

Hepatocytes
Fractionation of microsomes of hepatocytes by isotachophoresis (in French) (213) 177

Herbicides
GLC of *s*-triazines on a surface-bonded support in short columns (212) 85

Heteroborane anions
Reversed-phase ion-pair LC separations of some heteroborane anions (212) 283

HETP
The height equivalent to a theoretical plate of retentionless rectangular tubes (216) 1

Temperature dependence of HETP for columns with small amounts of liquid stationary phase (217) 225

Hexahydrocurcumines
GC and MS studies of ginger constituents. Identification of gingerdiones and new hexahydrocurcumin analogues (212) 75

High-back-pressure LC
High-back-pressure LC. I. Development of micro-HPLC using liquefied alkanes as the mobile phase (216) 153

High-performance displacement chromatography
High-performance displacement chromatography (218) 365

Histamine
Stability of the *o*-phthalaldehyde-histamine complex (212) 23

Capillary GC-MS determination of histamine in tuna fish causing scombroid poisoning (213) 475

Histamine analysis on a single column amino acid analyzer (219) 343

Hordatines
Separation of amines, guanidines and hydroxycinnamic acid amides by ion-exchange chromatography (214) 263

Hormones
Use of organic modifiers in the mobile phase for the reversed-phase HPLC separation of steroidal hormones (212) 108

Reversed-phase TLC for the separation and analysis of ecdysteroids (212) 211

LC measurement of cortisol in methylene chloride extracts of aqueous solutions (212) 221

Adsorption and reversed-phase partition HPLC of gibberellins (213) 507

Adsorption chromatographic separation of [^{125}I]progesterone-succinyl-tyrosine methyl ester (213) 511

Combined HPLC radioimmunoassay for cytokinins (214) 101

Isolation of hormonal proteins and antibodies by affinity chromatography (215) 143

Separation of subunits of pike eel gonadotropin by hydrophobic interaction chromatography (215) 229

General method for the assay of oral contraceptives by HPLC (216) 269

HPTLC determination of fluorescence-labelled cortisol (217) 349

Analysis of glycoprotein hormones and other medically important proteins by high-performance gel filtration chromatography (218) 581

Interaction of heme proteins and thyroid hormone. II. Localization of the site on thyroid hormone that binds to hemoglobin (219) 148

HPLC of progesterone and its metabolites (219) 152

HPLC analysis of permethylated cytokinins (219) 167

HT-2 toxin
HPLC of trichothecenes. I. Detection of T-2 toxin and HT-2 toxin (212) 370

Human cancer cell
Classification of human cancer cells by means of capillary GC and pattern recognition analysis (217) 231

Hydrazine
Separation of hydrazine, monomethylhydrazine, 1,1-dimethylhydrazine and 1,2-dimethylhydrazine by HPLC with electrochemical detection (214) 229

Hydrocarbons
Coordination polymers as adsorbents and stationary phases in GC. VII. Alkaline earth metal di-*n*-hexylphosphinates: specific interactions (212) 29

Characterization of Chromosorb porous polymer bead columns by GC retention values of light hydrocarbons and carbon dioxide (212) 261

Microextraction and GC analysis of selected petroleum hydrocarbons in water and fish tissue (212) 305

Determination of polychlorinated alkanes via carbon skeleton capillary GC (213) 73

Prediction of retention indices of aromatic hydrocarbons on phases of different polarity (216) 321

Chromatographic fingerprinting of coal extracts (217) 19

Determination of C_2-C_5 hydrocarbons in the atmosphere at low parts per 10^9 to high parts per 10^{12} levels (217) 301

Thermal stability of a poly(oxy aryl sulfonyl arylene) used as stationary phase in GLC (in French) (219) 197

Polarity of a poly(oxy aryl sulfonyl arylene) used as stationary phase in GLC (in French) (219) 213

High-resolution GC with liquid crystal glass capillaries. III. Separation of hydrocarbon isomers on cholesteric mesophase (in German) (219) 225

GC separation of diastereomeric isoprenoids as molecular markers of oil pollution (219) 393

Hydrocarbons, gasoline range
HPLC column switching techniques for rapid hydrocarbon group-type separations (218) 715

Hydrocinnamic acids
Correlation of Hammet substituent constants with reversed-phase "ion-pair" LC adsorption parameters of charged organic compounds (217) 421

Hydrogen bonds
The role of hydrogen bonds in chromatography. I. Amino- and hydroxyanthraquinones (212) 139

Hydrogen chloride
Determination of trace amounts of hydrogen halides in the atmosphere by collection in sampling tubes and subsequent derivatization and GC (217) 311

Hydrogen halides
Determination of trace amounts of hydrogen halides in the atmosphere by collection in sampling tubes and subsequent derivatization and GC (217) 311

Hydrogen ions
Migration behaviour of the hydrogen ion and its role in isotachophoresis of cations (217) 209

Concept of the effective mobility of the hydrogen ion and its use in cationic isotachophoresis (219) 21

Hydrogen peroxide
Sensitive, enzyme-catalyzed, chromogenic reagent for hydrogen peroxide and other peroxygen compounds on TLC plates (216) 413

Hydrophobicity
Comparison of conventional partitioning systems used for studying the hydrophobicity of polar organic compounds (216) 103

Chromatography in studies of quantitative structure-activity relationships (220) 71

Hydrophobic parameters
R_M values of steroids as an expression of their hydrophobic character. II. Effect of neighbouring groups on the hydrophobicity of substituents (216) 63

R_M values of steroids as an expression of their hydrophobic character. III.

Interpretation of the slopes of the R_M *versus* acetone concentration curves (216) 69

Partition coefficients of amino acids and hydrophobic parameters π of their side-chains as measured by TLC (216) 79

Correlation of Hammet substituent constants with reversed-phase "ion-pair" LC adsorption parameters of charged organic compounds (217) 421

Hydrophobic properties
Parameterization of hydrophobic properties of aqueous polymeric biphasic systems and water-organic solvent systems (212) 13

Hydroxy acids
New results in the GC separation of enantiomers of hydroxy acids and carbohydrates (217) 71

Hydroxyalkyl methacrylate gels
Hydroxyalkyl methacrylate gels derivatized with epichlorohydrin as supports for large-scale and high-performance affinity chromatography (215) 165

Hydroxyanthraquinones
The role of hydrogen bonds in chromatography. I. Amino- and hydroxyanthraquinones (212) 139

Hydroxybenzophenones
Analysis of 2-hydroxybenzophenone and 2'-hydroxyphenylbenzotriazole UV stabilizers by HPLC (214) 237

Hydroxycinnamic acid amides
Separation of amines, guanidines and hydroxycinnamic acid amides by ion-exchange chromatography (214) 263

Hydroxyindoleacetic acid
Determination of 5-hydroxytryptophan, serotonin and 5-hydroxyindoleacetic acid in rat and human brain and biological fluids by reversed-phase HPLC with electrochemical detection (218) 663

Hydroxyl groups
Determination of hydroxyl groups and water content in silica by nuclear magnetic resonance spectroscopy (213) 33

Hydroxyphenylbenzotriazoles
Analysis of 2-hydroxybenzophenone and 2'-hydroxyphenylbenzotriazole UV stabilizers by HPLC (214) 237

Hydroxyprogesterone
HPLC of progesterone and its metabolites (219) 152

Hydroxytryptophan
Determination of 5-hydroxytryptophan, serotonin and 5-hydroxyindoleacetic acid in rat and human brain and biological fluids by reversed-phase HPLC with electrochemical detection (218) 663

Hymenin
Separation and identification of stereoisomers of sesquiterpene lactones by multiple development of thin-layer chromatograms (212) 379

Hypocholesteremic agents
HPLC analysis of derivatized hypocholesteremic agents from fermentation broths (212) 234

I

Immunoadsorbents
Use of $F(ab')_2$ antibody fragments in the synthesis of immunoadsorbents for preparing monospecific antigen (212) 179

Immunoglobulins
Use of $F(ab')_2$ antibody fragments in the synthesis of immunoadsorbents for preparing monospecific antigen (212) 179

Application of 1,1'-carbonyldiimidazole-activated matrices for the purification of proteins. III. The use of 1,1'-carbonyldiimidazole-activated agaroses in the biospecific affinity chromatographic isolation of serum antibodies (218) 509

Indoles
HPLC determination of biogenic amines. I. Use of aqueous acidic mobile phases with silica columns (212) 153

HPLC determination of biogenic amines. II. Comparison of detection methods (213) 429

Determination of 5-hydroxytryptophan, serotonin and 5-hydroxyindoleacetic acid in rat and human brain and biological fluids by reversed-phase HPLC with electrochemical detection (218) 663

Infectivity profiles
Application of a general equation for controlled pore glass permeation chromatography to an aggregating, spherical virus (211) 53

Injection techniques
Peak broadening or splitting caused by solvent flooding after splitless or cold on-column injection in capillary GC (213) 3

Sample transfer in splitless injections in capillary GC (214) 118

Direct desorption of traps for capillary column GC (214) 197

Automatic injection in high-resolution GC: a programmed temperature vaporizer as a general purpose injection system (217) 81

Application of the split-splitless injector to environmental analysis (217) 91

Properties of a capillary column injector with a controlled rate of injection (217) 99

Large-volume injection of samples dissolved in a non-eluting solvent. Application to the determination of antipyrine using normal-phase HPLC (217) 509

System for injection of continuously variable amounts of sample vapours for physico-chemical measurements by GC (219) 427

Inorganic anions
Methylation of inorganic anions for GC determination (211) 361

Inosine
High-performance displacement chromatography (218) 365

Insecticides
HPLC determination of carbaryl and 1-naphthol at residue levels in various water sources by direct injection and trace enrichment (211) 135

GC-MS method for the quantitative analysis of carbofuran in water (212) 366

HPLC of fenvalerate (216) 303

Instrumental parameters
Very-high-speed LC. II. Some instrumental factors influencing performance (217) 369

Insulins
Improved HPLC method for the analysis of insulins and related compounds (218) 597

Internal standard technique
Limits of the internal standard technique in chromatography (218) 73

Iodide
Methylation of inorganic anions for GC determination (211) 361

Iodothyronines
Interaction of heme proteins and thyroid hormone. II. Localization of the site on thyroid hormone that binds to hemoglobin (219) 148

Ion-exchange kinetics
Computer simulation of the kinetics of ion exchange accompanied by complex formation in ionites (212) 133

Ion-exchange resins
Rapid high-resolution separation of oligosaccharides on silver form cation-exchange resins (212) 51

Physico-chemical and chromatographic properties of new ion exchangers. I. CM-Trisacryl M (213) 389

Advantages of a new ion-exchange resin for the LC of biogenic amines: a methacrylic acid polymer cross-linked with butanediol-diacrylate (216) 137

Ionic solutes
Ion chromatography of inorganic and organic ionic species using refractive index detection (218) 57

Ionites
Computer simulation of the kinetics of ion exchange accompanied by complex formation in ionites (212) 133

Isoascorbic acid
Separation of ascorbic and isoascorbic acids by ion-pair reversed-phase chromatography (in French) (219) 338

Isocitric acid
Isotachophoretic analysis of organic acids produced by fermentation from *n*-alkanes (216) 313

Isocyanates
Determination of trace atmospheric isocyanate concentrations by reversed-phase HPLC using 1-(2-pyridyl)piperazine reagent (212) 93

GC determination of free toluene diisocyanate in flexible urethane foams (213) 162

Isoelectric focusing
Computer simulation and experimental validation of isoelectric focusing in ampholine-free systems (211) 313

Isoelectric focusing in agarose under denaturating conditions (215) 373

Isoelectric points
Isoelectric points and molecular weights of proteins. A new table (Review) (220) 115

Isoflavones
Separation of genistin, daidzin and their aglucones, and coumesterol by gradient HPLC (211) 166

Isomerization
Kinetics of *n*-butene isomerization studied by stopped-flow GC (213) 15

Isomer separation
Separation and identification of stereoisomers of sesquiterpene lactones by multiple development of thin-layer chromatograms (212) 379

Semi-preparative HPLC and spectroscopic characterisation of eight geometric isomers of leukotriene A methyl ester (214) 249

Selectivity and efficiency of separation of isomers of organic acids by clathrate chromatography (218) 465

HPLC of bile pigments: separation and characterization of the urobilinoids (218) 647

Determination of cross-contamination of the diastereomers ephedrine and pseudoephedrine

by HPLC, TLC and carbon-13 nuclear magnetic resonance spectroscopy (219) 81

Influence of intramolecular interactions on the GC separation of epimers of dicyclopentadiene derivatives (in German) (219) 113

Use of routine preparative HPLC in the separation of isomers (219) 140

Separation of diamantane-3-oneoxime stereoisomers by preparative HPLC (219) 179

Analytical and preparative separation of the *cis*- and *trans*-isomers of 4-(4'-*tert.*-butylcyclohexyl)-4-methylpentan-2-one by reversed-phase HPLC (219) 183

High-resolution GC with liquid crystal glass capillaries. III. Separation of hydrocarbon isomers on cholesteric mesophase (in German) (219) 225

Separation of the N-trifluoroacetyl dimethyl esters of leukotriene D and E isomers by semi-preparative HPLC (219) 325

GC separation of diastereomeric isoprenoids as molecular markers of oil pollution (219) 393

Isopentenyladenosine
Combined HPLC radioimmunoassay for cytokinins (214) 101

Isopilocarpine
HPLC analysis of pilocarpine hydrochloride, isopilocarpine, pilocarpic acid and isopilocarpic acid in eye-drop preparations (212) 331

Isoprenoids
GC separation of diastereomeric isoprenoids as molecular markers of oil pollution (219) 393

Isotachophoresis
Concept of the effective mobility of the hydrogen ion and its use in cationic isotachophoresis (219) 21

Isovanillin
Assay for catechol-O-methyltransferase by HPLC with fluorescence detection (211) 129

L

Lactam antibiotics
LC of cephalosporin C and α-amino acid mixtures on polyfunctional polystyrene resins (211) 79

Quantitative analysis of Cephamycin C in fermentation broths by means of thin-layer spectrodensitometry (213) 129

Lanthanide polymers
Applications of selective complexation by a europium(III) coordination polymer sorbent for the pre-fractionation of volatile compounds (217) 275

Laser-based detectors
HPLC studies of coal liquids by laser-based detectors (211) 95

Lecithin
Separation of some lecithin components by HPLC (in German) (216) 389

Lectins
Application of polyacrylamide gradient gel electrophoresis to the study of multiple components of concanavalin A and related lectins (211) 233

Polyacrylhydrazido-agarose: preparation via periodate oxidation and use for enzyme immobilization and affinity chromatography (215) 55

Consideration of the nature of the lectin-carbohydrate interaction (215) 351

Affinity chromatography for the purification of lectins (a review) (215) 361

Leukotrienes
Semi-preparative HPLC and spectroscopic characterisation of eight geometric isomers of leukotriene A methyl ester (214) 249

Separation of the N-trifluoroacetyl dimethyl esters of leukotriene D and E isomers by semi-preparative HPLC (219) 325

Ligand densities
Quantitative ether cleavage of ligands in hydrophobic agaroses: precise determination of the degree of substitution (215) 235

Ligand heterogeneity
Theory of affinity electrophoresis. Evaluation of the effects of protein multivalency, immobilized ligand heterogeneity and microdistribution and determination of effective concentration of immobilized ligand (216) 43

Lipase
Large-scale purification of staphylococcal lipase by hydrophobic interaction chromatography (216) 295

Lipids
Behaviour of biomolecules in water-organic solvent-inorganic salt two-phase ternary systems (216) 346

Analytical separation of green-plant and animal neutral lipids by TLC (216) 417

Lipophilic properties
Evaluation of lipophilic properties for a series of phenols, using reversed-phase HPLC and HPTLC (214) 59

Lipoproteins
Separation of apolipoproteins A-I and A-II by ion-paired reversed-phase HPLC (216) 285

Lipotropin
Partition chromatography and HPLC of β-lipotropin and synthetic β-endorphin analogues (215) 255

Liquid chromatography
The nomenclature of chromatography. II. Liquid chromatography (Review) (220) 29

Liquid crystals
Evaluation of liquid crystal smectic mesophases for GLC separations (214) 161

High-resolution GC with liquid crystal glass capillaries. III. Separation of hydrocarbon isomers on cholesteric mesophase (in German) (219) 225

Lymphocytes
Continuous-flow scanning of selected HPLC peak components by microprocessor control. Application to analysis of extracts from human lymphocytes (219) 255

Lysergic acid
Mechanism of zwitterion-pair chromatography. II. Ampicilline, lysergic acid, tryptophan and other solutes (218) 355

Lysozyme
Separation of the tryptic peptides from reduced, alkylated hen egg white lysozyme by HPLC (214) 185

M

Maleic anhydride
Direct GC determination of the products of catalytic air oxidation of *n*-butene-1 to maleic anhydride in the gaseous reaction mixture (214) 217

Maltitol syrups
GLC separation and determination of the components of maltitol syrups (216) 219

Maltosaccharides
HPLC of maltosaccharides (212) 126

GLC separation and determination of the components of maltitol syrups (216) 219

Malvidin glucoside
Separation of anthocyanin chalcones by HPLC (214) 222

Manning and Moore procedure
Determination of the optical purity of amino acids by HPLC. Modification of the Manning and Moore procedure (215) 279

Mass spectrometry
Middle molecule mass spectrometry (a review) (218) 21

Maytansinoids
HPLC separation and quantitation of maytansinoids in *Maytenus ilicifolia* (213) 340

Metabolic profiles
Classification of human cancer cells by means of capillary GC and pattern recognition analysis (217) 231

Metal chelates
Adsorption and desorption of proteins in metal chelate affinity chromatography. Purification of albumin (215) 333

Manganese(II)-bis(3-heptafluorobutyryl-1*R*-camphorate): a versatile agent for the resolution of racemic cyclic ethers by complexation GC (217) 51

Metal complexes
Coordination polymers as adsorbents and stationary phases in GC. VII. Alkaline earth metal di-*n*-hexylphosphinates: specific interactions (212) 29

High-performance paper electrophoresis. II. Comparison of separations obtained by high-voltage paper electrophoresis and high-performance paper electrophoresis (212) 374

Applications of HPLC in organometallic chemistry: ligand exchange in complexes of ylides (in German) (213) 47

Chromatographic study of optical resolution. VII. Directional ion-association model for the optical resolution of *cis*-$[Co(O)_2(N)_4]^+$ type of complexes by the antimony *d*-tartrate ion (213) 323

Applications of selective complexation by a europium(III) coordination polymer sorbent for the pre-fractionation of volatile compounds (217) 275

HPLC of *cis*-dichlorodiammineplatinum(II) using chemically-bonded and solvent-generated ion exchangers (217) 405

Chromatographic study of optical resolution. VIII. Theoretical study of the chromatographic behaviour of the enantiomers of racemic complex cations on a cation-exchange column (219) 29

HPLC of inorganic platinum(II) complexes using solvent-generated anion exchangers. II. The effect of electrolytes on solute retention (219) 235

Metal doping
Response enhancement by metal doping of a silicon-selective flame ionization detector for GC (213) 397

Metal ions
Separation of metal ions by paper electrophoresis in N-(phosphono)iminodisulphuric acid and glycine-N,N-bis(methylphosphonic acid) (in German) (211) 177

Metal salts
Use of metal salts as fluorescence-inducing reagents in TLC (217) 329

Metal traces
Stationary phase characterization in HPLC. A test for trace metal activity in octadecyl bonded silica gel (217) 399

Methacrylates
Some carriers for the immobilization of enzymes based on derivatized poly(vinyl alcohol) and on copolymers of methacrylates with different spacer lengths (215) 13

Methacrylic acid polymers
Advantages of a new ion-exchange resin for the LC of biogenic amines: a methacrylic acid polymer cross-linked with butanediol-diacrylate (216) 137

Methylazoxymethanol
Determination of the carcinogen methylazoxy-methyl-β-D-glucosiduronic acid in rat bile and urine (211) 403

Methylcarbazoles
The chemistry of carbazoles. IX. Substituent effect in the GLC of methylcarbazoles (211) 388

Methylcolchiceinamide
HPLC determination of N,N-dimethylcolchiceinamide and its metabolites, N-methylcolchiceinamide and colchiceinamide, in microbial culture (219) 321

Methylimidazole
Quantitative ion-pair extraction of 4(5)-methyl-imidazole from caramel colour and its determination by reversed-phase ion-pair LC (211) 213

Methyl pyridiniumaldoxime chloride
Determination of N-methyl pyridinium-2-aldoxime chloride and its hydrolytic by-products by ion-pair HPLC (212) 361

Mevinolin
HPLC analysis of derivatized hypocholesteremic agents from fermentation broths (212) 234

Microbicides
Speciation of aryloxyethoxyethyl benzyl dimethyl ammonium salts by glass capillary GC and HPLC (213) 463

Microbore columns
High-resolution reversed-phase LC utilizing microbore column concatenation (216) 9

Application of ultra-micro HPLC to trace analysis (218) 199

Recycling LC using microbore columns (219) 1

Microcoulometric detector
Detection of brominated and chlorinated organics by a GC microcoulometric detector. Effect of pyrolysis tube conditions (212) 115

Microsomes
Fractionation of microsomes of hepatocytes by isotachophoresis (in French) (213) 177

Middle molecule mass spectrometry
Middle molecule mass spectrometry (a review) (218) 21

Miotic agents
HPLC analysis of pilocarpine hydrochloride, isopilocarpine, pilocarpic acid and isopilocarpic acid in eye-drop preparations (212) 331

Mobile phase composition
LC with crown ether-containing mobile phases. I. Capacity factors of amino compounds on a hydrophobic stationary phase (211) 1

Use of organic modifiers in the mobile phase for the reversed-phase HPLC separation of steroidal hormones (212) 108

HPLC determination of biogenic amines. I. Use of aqueous acidic mobile phases with silica columns (212) 153

Hydrophobic chromatography with dynamically coated stationary phases. V. "Duplex soap chromatography" (212) 271

Solvent strength of multicomponent mobile phases in LSC. Binary-solvent mixtures and solvent localization (214) 1

Solvent strength of multicomponent mobile phases in LSC. Mixtures of three or more solvents (214) 21

Gradient elution in LC. XIV. Theory of ternary gradients in reversed-phase LC (214) 35

High-back-pressure LC. I. Development of micro-HPLC using liquefied alkanes as the mobile phase (216) 153

Non-ionic mobile phase dopants. I. Chiral charge-transfer acceptors and helicene resolution (216) 333

Behaviour of biomolecules in water-organic solvent-inorganic salt two-phase ternary systems (216) 346

Reversed-phase ion-pair chromatography with UV-absorbing ions in the mobile phase (218) 31

Computer-controlled single-pump solvent programmer for HPLC (218) 83

Systematic study of ternary solvent behaviour in reversed-phase LC (218) 261

HPLC of organic acids on bare silica (218) 285

Theoretical basis for systematic optimization of mobile phase selectivity in LSC. Solvent-solute localization effects (218) 299

Buffer-focusing chromatography using multi-component electrolyte elution systems (218) 483

HPLC of amino acids, peptides and proteins. XXXVI. Organic solvent modifier effects in the separation of unprotected peptides by reversed-phase LC (218) 497

Reversed-phase LCC of pyrimidine and purine derivatives. I. Unbuffered binary aqueous organic mobile phases on octadecylsilica (219) 245

Mobile phase pre-heater
Effect of mobile phase pre-heating on HPLC column performance: a new type of column thermostat (213) 243

Molecular dimensions
Comparison of dimensions obtained by size-exclusion chromatography and X-ray diffraction of rigid molecules (218) 473

Molecular retention indices
GC of homologous esters. XV. Molecular retention indices of aliphatic esters (219) 45

Molecular-sieve effects
The Faraday Society's discussion at Reading in 1949 and the exploitation of molecular-sieve effects for chemical separations (215) 1

Molecular weight distribution
Chromatography of polycarbamoyl sulphonates (anionic polyurethane derivatives) (214) 69

Molecular weights
Isoelectric points and molecular weights of proteins. A new table (Review) (220) 115

Monohydroxybiphenyls
GC behaviour of mono- and dihydroxybiphenyls on various silicone phases (211) 150

Monomethylhydrazine
Separation of hydrazine, monomethylhydrazine, 1,1-dimethylhydrazine and 1,2-dimethylhydrazine by HPLC with electrochemical detection (214) 229

Monosaccharides
Separation of monosaccharides as trimethylsilylated alditols on fused-silica capillary columns (213) 146

Rapid separation of malto-, xylo- and cello-oligosaccharides (DP 2-9) on cation-exchange resin using water as eluent (213) 151

Morphine
Application of statistical optimization methods to the separation of morphine, codeine, noscapine and papaverine in reversed-phase ion-pair chromatography (211) 201

Mucopolysaccharides
Hydrophobic interaction chromatography of mucopolysaccharides. Examination of fundamental conditions for fractionation of heparin on hydrophobic gels (213) 439

Mustards
TLC of some sulfur and nitrogen mustards (213) 173

Mutagenic agents
R_M and log P values of 5-nitroimidazoles (216) 93

Mycotoxins
HPLC of trichothecenes. I. Detection of T-2 toxin and HT-2 toxin (212) 370

Determination of patulin and penicillic acid in unroasted cocoa beans (214) 135

Rapid determination of aflatoxin M_1 in dairy products by reversed-phase HPLC (219) 156

Assay method for aflatoxin in milk (219) 162

N

Nadolol
Separation of nadolol diastereoisomers by reversed-phase HPLC (216) 374

Naphthoic acids
Selectivity and efficiency of separation of isomers of organic acids by clathrate chromatography (218) 465

Naphthol
HPLC determination of carbaryl and 1-naphthol at residue levels in various water sources by direct injection and trace enrichment (211) 135

Naproxen
Separation of acidic drugs in the µg/ml range in untreated blood plasma by direct injection on LC columns (218) 671

Neomycin
Determination of the relative amounts of the B and C components of neomycin by ion-exclusion chromatography using refractometric detection (211) 223

Neuroleptic drugs
Preparative purification of the peptide des-enkephalin γ-endorphin. Comparison of HPLC and counter-current chromatography (212) 356

Neuropeptides
HPLC and field desorption MS measurement of picomole amounts of endogenous neuropeptides in biologic tissue (217) 437

Nicotine alkaloids
Rapid method for the analysis of tobacco nicotine alkaloids (211) 111

Nitroanilines
Preparation and evaluation of N-(2-amino-ethyl)-γ-aminopropyltrimethoxysilane-treated silica columns for the HPLC analysis of some aromatic amines (212) 251

Nitrobenzoic acids
Selectivity and efficiency of separation of isomers of organic acids by clathrate chromatography (218) 465

Nitrocinnamic acids
Selectivity and efficiency of separation of isomers of organic acids by clathrate chromatography (218) 465

Nitrogen mustards
TLC of some sulfur and nitrogen mustards (213) 173

Nitrogen-selective detector
Nitrogen-selective detector for HPLC (216) 127

Nitroglycerin
Separation, identification and quantitation of nitroglycerin and its metabolic or hydrolysis products (216) 239

Nitroimidazoles
R_M and log P values of 5-nitroimidazoles (216) 93

Nitrophenols
HPLC determination of 4-nitroso- and 4-nitrophenols in the presence of phenol and alkylphenols (216) 361

Nitroprusside
HPLC determination of sodium nitroprusside (212) 339

Nitrosamines
Confirming the presence of N-nitrosamines in ambient air and cigarette smoke by converting to and photochemically altering their corresponding N-nitroamines (213) 253

Nitrosophenols
HPLC determination of 4-nitroso- and 4-nitrophenols in the presence of phenol and alkylphenols (216) 361

Nocodazole
HPLC determination of Nocodazole in liposomes and L 1210 cells (213) 170

Nomenclature
The nomenclature of chromatography. II. Liquid chromatography (Review) (220) 29

The nomenclature of chromatography. III. General rules for future revisions (Review) (220) 65

Non-electrolytes
Estimation of aqueous solubilities of organic non-electrolytes using LC retention data (218) 409

Noradrenaline
GC determination of some biogenic amines as their pentafluorobenzoyl derivatives in the picogram range and its applicability to biological materials (213) 413

Norepinephrine
Resolution of enantiomers of norepinephrine and epinephrine by reversed-phase HPLC (213) 327

Plasma catecholamines in hypertension and pheochromocytoma determined using ion-pair reversed-phase chromatography with amperometric detection. Investigation of the separation mechanism and clinical methodology (217) 523

Norverapamil
Rapid HPLC method for the measurement of verapamil and norverapamil in blood plasma or serum (218) 621

Noscapine
Application of statistical optimization methods to the separation of morphine, codeine, noscapine and papaverine in reversed-phase ion-pair chromatography (211) 201

Noxyptyline
GC determination of noxyptyline in substance, tablets and in biological material (219) 291

Nucleic acids
Aromatic interaction chromatography. Acriflavin gel for separation of nucleotides, oligonucleotides and nucleic acids (215) 243

Nucleobases
LC separation of purines, pyrimidines and their nucleosides on silica gel columns (211) 337

Nucleosides
LC separation of purines, pyrimidines and their nucleosides on silica gel columns (211) 337

Separation of cAMP from adenine nucleotides and nucleosides by electrophoresis on cellulose acetate membranes (211) 409

HPLC of the oligonucleotides (217) 479

Study of the performance of cation-exchange columns in open-tubular microcapillary LC (218) 189

Nucleotides
Separation of cAMP from adenine nucleotides and nucleosides by electrophoresis on cellulose acetate membranes (211) 409

Aromatic interaction chromatography. Acriflavin gel for separation of nucleotides, oligonucleotides and nucleic acids (215) 243

HPLC of the oligonucleotides (217) 479

Mechanism of zwitterion-pair chromatography. I. Nucleotides (218) 341

Improved method of resolving nucleotides by reversed-phase HPLC (219) 133

Use of periodate and methylamine for the quantitation of intracellular 5-fluoro-2'-deoxyuridine-5'-monophosphate by HPLC (219) 273

Nystatin
Identification and determination of the qualitative composition of nystatin using TLC and HPLC (216) 367

O

Octametaphosphate
Chromatographic study of the acidic hydrolysis of cyclic octametaphosphate (211) 87

Odorous volatiles
Determination of odorous volatiles in air using chromatographic profiles (216) 199

Oestrogens
Use of organic modifiers in the mobile phase for the reversed-phase HPLC separation of steroidal hormones (212) 108

HPLC of naturally occurring estrogens (212) 239

General method for the assay of oral contraceptives by HPLC (216) 269

Oils
Microextraction and GC analysis of selected petroleum hydrocarbons in water and fish tissue (212) 305

Fractionation of polar organic constituents in environmental samples using the lipophilic dextran gels Sephadex LH-20 and Sephasorb HP Ultrafine. Application to a weathered Ekofisk crude oil (214) 83

TLC analysis of aromatic hydrocarbons in crude oil and in petroleum products as their η^6-arene-η^5-cyclopentadienyliron hexafluorophosphates (219) 349

GC separation of diastereomeric isoprenoids as molecular markers of oil pollution (219) 393

Oligonucleotides
Aromatic interaction chromatography. Acriflavin gel for separation of nucleotides, oligonucleotides and nucleic acids (215) 243

HPLC of the oligonucleotides (217) 479

Oligopeptides
HPLC and field desorption MS measurement of picomole amounts of endogenous neuropeptides in biologic tissue (217) 437

Analysis of N-acetyl-N,O,S-permethylated peptides by combined LC-MS (218) 519

Oligosaccharides
Rapid high-resolution separation of oligosaccharides on silver form cation-exchange resins (212) 51

Rapid separation of malto-, xylo- and cello-oligosaccharides (DP 2-9) on cation-exchange resin using water as eluent (213) 151

Opiates
Application of statistical optimization methods to the separation of morphine, codeine, noscapine and papaverine in reversed-phase ion-pair chromatography (211) 201

Ion-exchange and straight-phase partition HPLC of the opiates on silica (212) 229

Fluorescence and UV detection of opiates separated by reversed-phase HPLC (213) 337

Optimization
Optimization of operating parameters in overpressured TLC (211) 45

Application of statistical optimization methods to the separation of morphine, codeine, noscapine and papaverine in reversed-phase ion-pair chromatography (211) 201

Optimisation in chromatography. Theory and application to the separation of aromatic acids in reversed-phase LC (213) 357

Very-high-speed LC. II. Some instrumental factors influencing performance (217) 369

Theoretical basis for systematic optimization of mobile phase selectivity in LSC. Solvent-solute localization effects (218) 299

Assessment and optimization of system parameters in size exclusion separation of proteins on diol-modified silica columns (218) 535

TLC as a pilot technique for the optimization of preparative column chromatography (218) 729

Orange peel oils
Role of water in qualitative and quantitative determination of polymethoxylated flavones by straight-phase HPLC: application to orange peel oils (211) 61

Organic acids
Alternate procedure for the preparation and separation of benzyl derivatives of organic acids (214) 131

Concentration and isolation of organic acids on graphitized carbon black (219) 263

Organic compounds
Evaluation of a liquid-liquid extraction technique for water pollutants (212) 187

Fractionation of polar organic constituents in environmental samples using the lipophilic dextran gels Sephadex LH-20 and Sephasorb HP Ultrafine. Application to a weathered Ekofisk crude oil (214) 83

Organic modifiers
Use of organic modifiers in the mobile phase for the reversed-phase HPLC separation of steroidal hormones (212) 108

HPLC of amino acids, peptides and proteins. XXXVI. Organic solvent modifier effects in the separation of unprotected peptides by reversed-phase LC (218) 497

Organic non-electrolytes
Estimation of aqueous solubilities of organic non-electrolytes using LC retention data (218) 409

Organic vapours
Dimethylformamide and carbon disulphide desorption efficiencies for organic vapours on gas-sampling charcoal tube analyses with a GC backflush technique (217) 317

Organochlorine compounds
Determination of polychlorinated alkanes via carbon skeleton capillary GC (213) 73

Application of the split-splitless injector to environmental analysis (217) 91

Organophosphorus compounds
Application of the split-splitless injector to environmental analysis (217) 91

Studies on the GC behaviour of organophosphorus esters (219) 297

Organosilicon compounds
Contributions to the chemistry of silicon-sulphur compounds. XXVI. Relationships between chromatographic behaviour and structure of triorganosilanes, triorganosilanethiols, hexaorganodisilthianes and triphenylthioalkylsilanes (219) 302

Organotin compounds
Simultaneous determination of trialkyltin homologues in biological materials (216) 209

Osmogenes
Determination of odorous volatiles in air using chromatographic profiles (216) 199

Oxirane carboxylic acids
Fluorescence labelling in trace analysis of biological samples. Simultaneous determination of free fatty acids and related carboxylic compounds (217) 491

Oxiranes
Manganese(II)-bis(3-heptafluorobutyryl-1*R*-camphorate): a versatile agent for the resolution of racemic cyclic ethers by complexation GC (217) 51

Oxocarboxylic acids
Analysis of the oxocarboxylic acid fraction in serum and urine as O-methyloximes by thermionic specific detection (217) 255

Oxytocin
Assay of the combined formulation of ergometrine and oxytocin by HPLC (214) 307

P

Packings
Characterization of Chromosorb porous polymer bead columns by GC retention values of light hydrocarbons and carbon dioxide (212) 261

Comparison of column packings for trace analysis of free amines by GLC (217) 177

GC behaviour of chemically bonded microparticulate silica (217) 191

Packing techniques
Rapid method for packing microparticulate columns packed with a chemically bonded stationary phase for HPLC (211) 378

Packing of Toyopearl columns for gel filtration. IV. Gravitational packing and influence of slurry reservoir size (211) 383

Pairing agents
Mechanism of zwitterion-pair chromatography. I. Nucleotides (218) 341

Papaverine
Application of statistical optimization methods to the separation of morphine, codeine, noscapine and papaverine in reversed-phase ion-pair chromatography (211) 201

Parthenin
Separation and identification of stereoisomers of sesquiterpene lactones by multiple development of thin-layer chromatograms (212) 379

Particle analysis
Comparison of sedimentation field flow fractionation with chromatographic methods for particulate and high-molecular-weight macromolecular characterizations (218) 217

Particle size
Application of a general equation for controlled pore glass permeation chromatography to an aggregating, spherical virus (211) 53

Partition coefficients
Parameterization of hydrophobic properties of aqueous polymeric biphasic systems and water-organic solvent systems (212) 13

Partition coefficients of amino acids and hydrophobic parameters π of their side-chains as measured by TLC (216) 79

R_M and log P values of 5-nitroimidazoles (216) 93

Comparison of conventional partitioning systems used for studying the hydrophobicity of polar organic compounds (216) 103

Partitioning
Chromatography in studies of quantitative structure-activity relationships (220) 71

Partitioning effects
GLSC: effects of adsorption and partitioning (213) 203

Pattern recognition
Classification of human cancer cells by means of capillary GC and pattern recognition analysis (217) 231

Patulin
Determination of patulin and penicillic acid in unroasted cocoa beans (214) 135

PCBs
Evaluation of liquid crystal smectic mesophases for GLC separations (214) 161

Peak broadening
Peak broadening or splitting caused by solvent flooding after splitless or cold on-column injection in capillary GC (213) 3

Microcapillary LC in open tubular columns with diameters of 10-50 μm. Potential application to chemical ionization MS detection (218) 137

Peak collection
Fraction collector controller for collecting LC peaks (212) 105

Peak dispersion
Dispersion of peaks by short straight open tubes in LC systems (218) 97

Effect of radial thermal gradients in elevated temperature HPLC (218) 123

Peak identification
Identification of ginsenosides from *Panax ginseng* in fractions obtained by HPLC by field desorption MS, multiple internal reflection IR spectroscopy and TLC (212) 37

Peak shape
Doubly peaked chromatograms (213) 355

Penicillamine
Rapid automated determination of D-penicillamine in plasma and urine by ion-exchange HPLC with electrochemical detection using a gold electrode (218) 615

Penicillic acid
Determination of patulin and penicillic acid in unroasted cocoa beans (214) 135

Pentachlorophenol
Determination of chlorinated phenols by HPLC (213) 83

Determination of pentachlorophenol and related compounds in animal materials by HPLC and GC (216) 229

Pentobarbital
Determination of 5-ethyl-5-(1-ethylpropyl)-barbituric acid in pentobarbital by HPLC (219) 431

Pepsin
Hydroxyalkyl methacrylate gels derivatized with epichlorohydrin as supports for large-scale and high-performance affinity chromatography (215) 165

Peptides
Factors affecting retention and resolution of peptides in HPLC (211) 15

TLC of closely related polypeptides on silanized silica gel (211) 29

Preparative purification of the peptide des-enkephalin γ-endorphin. Comparison of HPLC and counter-current chromatography (212) 356

Systematic separation of medium-sized biologically active peptides by HPLC (213) 275

Distinctive test with copper(II)-ninhydrin reagent for small α-peptides separated PC (213) 307

Analysis of protein and peptide mixtures. Evaluation of three sodium dodecyl sulphate-polyacrylamide gel electrophoresis buffer systems (213) 491

Separation of Dns-amino acids using reversed-phase HPLC: a sensitive method for determining N-termini of peptides and proteins (213) 501

Separation of the tryptic peptides from reduced, alkylated hen egg white lysozyme by HPLC (214) 185

Improved separation of basic peptides in anion-exchange chromatography (215) 109

Partition chromatography and HPLC of β-lipotropin and synthetic β-endorphin analogues (215) 255

Size-exclusion HPLC of protected peptides (215) 263

High-speed gel filtration of glycopolypeptides in 6 M guanidine hydrochloride (215) 289

Covalent chromatography as a means of isolating thiol peptides from large proteins. Application to human ceruloplasmin (215) 341

Behaviour of biomolecules in water-organic solvent-inorganic salt two-phase ternary systems (216) 346

Reversed-phase HPLC of dermorphins, opiate-like peptides from amphibian skins (216) 355

HPLC and field desorption MS measurement of picomole amounts of endogenous neuropeptides in biologic tissue (217) 437

HPLC of amino acids, peptides and proteins. XXXVI. Organic solvent modifier effects in the separation of unprotected peptides by reversed-phase LC (218) 497

Analysis of N-acetyl-N,O,S-permethylated peptides by combined LC-MS (218) 519

Improved HPLC method for the analysis of insulins and related compounds (218) 597

Separation of large polypeptides by HPLC (218) 603

HPLC of protein polypeptides on porous silica gel columns (TSK-GEL SW) in the presence of sodium dodecyl sulphate: comparison with SDS-polyacrylamide gel electrophoresis (219) 123

Pergolide
HPLC determination of pergolide and its metabolite, pergolide sulfoxide, in microbial extracts (219) 317

Permittivity detector
Improved high-frequency permittivity detector for the chromatography of alcohols and other polar species (212) 167

Peroxygen compounds
Sensitive, enzyme-catalyzed, chromogenic reagent for hydrogen peroxide and other peroxygen compounds on TLC plates (216) 413

Pesticides
HPLC determination of carbaryl and 1-naphthol at residue levels in various water sources by direct injection and trace enrichment (211) 135

Use of 214-nm and 229-nm discrete-line sources for the UV absorbance detection of some pesticides separated by HPLC (211) 144

GLC of *s*-triazines on a surface-bonded support in short columns (212) 85

GC-MS method for the quantitative analysis of carbofuran in water (212) 366

Quantitative determination of 1,2-dibromo-3-chloropropane in whole rat blood and drinking water by GC (213) 156

HPLC determination of azinphos methyl and azinphos methyl oxon in fruits and vegetables (214) 140

Determination of chlorophacinone in formulations by reversed-phase ion-pair chromatography (214) 335

Simultaneous determination of trialkyltin homologues in biological materials (216) 209

HPLC of fenvalerate (216) 303

Petroleum
Microextraction and GC analysis of selected petroleum hydrocarbons in water and fish tissue (212) 305

HPLC column switching techniques for rapid hydrocarbon group-type separations (218) 715

TLC analysis of aromatic hydrocarbons in crude oil and in petroleum products as their η^6-arene-η^5-cyclopentadienyliron hexafluorophosphates (219) 349

pH effects
Effect of pH on the GC behaviour of silica gel (219) 101

pH gradients
Buffer-focusing chromatography using multi-component electrolyte elution systems (218) 483

Phaeophytin
Rapid, quantitative separation of chlorophylls and their degradation products by HPLC (213) 349

Pharmaceuticals
Alkylation with alkyl halides as a derivatization method for the GC determination of acidic pharmaceuticals (Review) (220) 275

Phase transitions
Influence of the LC mobile phase on the phase transitions of alkyl-bonded silicas studied by GC (214) 202

Phenacetin
Large-volume injection of samples dissolved in a non-eluting solvent. Application to the determination of antipyrine using normal-phase HPLC (217) 509

Phenolic compounds
GC behaviour of mono- and dihydroxybiphenyls on various silicone phases (211) 150

Prediction of retention behaviour of some phenolic antioxidants (213) 363

C_{18} reversed-phase trace enrichment of chlorinated phenols, guaiacols and catechols in water (214) 327

Separation of plant polyphenolics by chromatography on a boronate resin (216) 261

High-performance displacement chromatography (218) 365

HPLC of flavonoids in barley and hops (218) 683

HPLC analysis of carvacrol and thymol in the essential oil of *Thymus capitatus* (219) 332

Phenolic lipids
Long-chain phenols. XXI. Quantitative analysis of the phenolic lipids in technical cashew nut-shell liquid, from *Anacardium occidentale,* by HPLC (213) 287

Phenols
Chromatographic behaviour of aromatic compounds on non-polar phases with adsorbed quaternary ammonium salts (212) 1

Determination of chlorinated phenols by HPLC (213) 83

Evaluation of lipophilic properties for a series of phenols, using reversed-phase HPLC and HPTLC (214) 59

HPLC determination of 4-nitroso- and 4-nitrophenols in the presence of phenol and alkylphenols (216) 361

Polarity of a poly(oxy aryl sulfonyl arylene) used as stationary phase in GLC (in French) (219) 213

Phenothiazines
LC of demoxepam and phenothiazines using a post-column photochemical reactor and fluorescence detection (217) 463

Phenylacetates
Chromatographic behaviour of aromatic compounds on non-polar phases with adsorbed quaternary ammonium salts (212) 1

Phenylalkylcarboxylic acids
Correlation of Hammet substituent constants with reversed-phase "ion-pair" LC adsorption parameters of charged organic compounds (217) 421

Phenylenediamines
Antioxidants and stabilizers. LXXXVI. Use of chromatography in the investigation of the mechanism of action of amine antidegradants (211) 299

Antioxidants and stabilizers. LXXXVII. The chromatographic behaviour of transformation products of an antioxidant, N,N'-diphenyl-1,4-phenylenediamine (216) 251

Phenylethylamines
Group-contribution approach to the behaviour of 2-phenylethylamines in reversed-phase HPLC (218) 639

Phenylpyrazolones
Application of the combined redox-complexation reaction to the detection of antipyrine and its derivatives in TLC (219) 189

Phosphate buffers
Purification of potassium phosphate for HPLC (214) 267

Phosphate esters
PC separation of phosphate esters, tricarboxylic cycle acids and amino acids in extracts from malaria parasites (214) 111

Phosphates
Chromatographic study of the acidic hydrolysis of cyclic octametaphosphate (211) 87

Studies on the GC behaviour of organo-phosphorus esters (219) 297

Study of the acidic hydrolysis of cyclic trimeta-phosphate by LC (219) 385

Phosphogluconate dehydrogenase
Purification of 6-phosphogluconate dehydrogenase from *Acer pseudoplatanus* L. using immobilised Procion Red HE-3B (219) 419

Phospholipids
Separation of some lecithin components by HPLC (in German) (216) 389

Phosphonates
Studies on the GC behaviour of organo-phosphorus esters (219) 297

Photoactive compounds
TLC assay of photoactive compounds (furocoumarins) using the fungus *Penicillium expansum* as a test organism (214) 349

Phthalaldehyde
Stability of the *o*-phthalaldehyde-histamine complex (212) 23

Phthalates
High-back-pressure LC. I. Development of micro-HPLC using liquefied alkanes as the mobile phase (216) 153

Application of ultra-micro HPLC to trace analysis (218) 199

Physico-chemical measurements
System for injection of continuously variable amounts of sample vapours for physico-chemical measurements by GC (219) 427

Phytoestrogens
Separation of genistin, daidzin and their aglucones, and coumesterol by gradient HPLC (211) 166

Pilocarpine
HPLC analysis of pilocarpine hydrochloride, isopilocarpine, pilocarpic acid and isopilo-carpic acid in eye-drop preparations (212) 331

Pinitol
Separation of pinitol and some other cyclitols by HPLC (211) 295

Plant hormones
Adsorption and reversed-phase partition HPLC of gibberellins (213) 507

Combined HPLC radioimmunoassay for cytokinins (214) 101

HPLC analysis of permethylated cytokinins (219) 167

Plant screening
Plant screening by chemical and chromatographic procedures under field conditions (213) 113

Plastics additives
Analysis of additives in plastics by high-performance size-exclusion chromatography (218) 703

Plate number
Temperature dependence of HETP for columns with small amounts of liquid stationary phase (217) 225

Platinum complexes
HPLC of *cis*-dichlorodiammineplatinum(II) using chemically-bonded and solvent-generated ion exchangers (217) 405

HPLC of inorganic platinum(II) complexes using solvent-generated anion exchangers. II. The effect of electrolytes on solute retention (219) 235

Poiseuille flow
Doubly peaked chromatograms (213) 355

Polarity criterion
New criterion for polarity of stationary phases in GC (217) 139

Polarity rating
Practical system for polarity rating of packed GLC columns (213) 317

Polarographic detector
Extra-column effects in polarographic *versus* UV detection in HPLC (218) 45

Pollutants
Glass capillary columns in the GC separation of aromatic amines. II. Application to samples from workplace atmospheres using nitrogen-selective detection (211) 103

GC behaviour of mono- and dihydroxybiphenyls on various silicone phases (211) 150

Detection of benzidines on thin-layer chromatograms with fluorescamine (211) 304

Determination of trace atmospheric isocyanate concentrations by reversed-phase HPLC using 1-(2-pyridyl)piperazine reagent (212) 93

Detection of brominated and chlorinated organics by a GC microcoulometric detector. Effect of pyrolysis tube conditions (212) 115

Evaluation of a liquid-liquid extraction technique for water pollutants (212) 187

C_{18} reversed-phase trace enrichment of chlorinated phenols, guaiacols and catechols in water (214) 327

Electron-capture GC determination of sulphide as a new pentafluorobenzyl derivative (219) 312

Polyacrylamide gels
Simple gel apparatus for horizontal polyacrylamide and agarose gel electrophoresis (211) 257

Polyacrylhydrazido-agarose
Polyacrylhydrazido-agarose: preparation via periodate oxidation and use for enzyme immobilization and affinity chromatography (215) 55

Polyamides
Analysis of resinous or fatty polyamides using GC and GPC (213) 55

Polyamines
Distinctive test with copper(II)-ninhydrin reagent for small α-peptides separated PC (213) 307

Polycarbamoyl sulphonates
Chromatography of polycarbamoyl sulphonates (anionic polyurethane derivatives) (214) 69

Polychlorinated alkanes
Determination of polychlorinated alkanes via carbon skeleton capillary GC (213) 73

Polychlorinated biphenyls
see PCBs

Polychlorinated dibenzodioxins
Determination of chlorinated dibenzo-*p*-dioxin contaminants in 2,4-D products by GC-MS techniques (217) 289

Polycyclic aromatic hydrocarbons
see also polynuclear aromatic hydrocarbons

Evaluation of liquid crystal smectic mesophases for GLC separations (214) 161

TLC analysis of aromatic hydrocarbons in crude oil and in petroleum products as their η^6-arene-η^5-cyclopentadienyliron hexafluorophosphates (219) 349

Polyethylene glycol
Analysis of polar compounds on polyethylene glycol glass capillary columns. The influence of KF and Na_3PO_4 on the change in partial molar free energy (217) 183

Polyethylene glycols
HPLC analysis of polyethylene glycols (211) 160

Polymer beads
Characterization of Chromosorb porous polymer bead columns by GC retention values of light hydrocarbons and carbon dioxide (212) 261

Sorption behaviour of several organic compounds on 6-deoxycyclodextrin polyurethane resins (219) 108

Polymer gels
Polymer gels (215) 7

Polymers
Concentration effects in gel permeation chromatography. VII. Viscosity phenomena and column geometry (211) 39

Some carriers for the immobilization of enzymes based on derivatized poly(vinyl alcohol) and on copolymers of methacrylates with different spacer lengths (215) 13

Polymeric supports bearing isonitrile functional groups for covalent fixation of biologically active molecules (a review) (215) 31

Coupling of proteins and other amines to Sepharose by bromine oxidation and reductive amination (215) 45

Scanning electron microscopy as an aid in GC. Study of porous polymers coated with stationary phases (217) 109

Applications of selective complexation by a europium(III) coordination polymer sorbent for the pre-fractionation of volatile compounds (217) 275

Middle molecule mass spectrometry (a review) (218) 21

Comparison of dimensions obtained by size-exclusion chromatography and X-ray diffraction of rigid molecules (218) 473

Polymethoxylated flavones
Role of water in qualitative and quantitative determination of polymethoxylated flavones by straight-phase HPLC: application to orange peel oils (211) 61

Polymyxins
Preparative-scale separation of polymyxins with an analytical HPLC system by using displacement chromatography (215) 295

Isolation and characterization of three new polymyxins in polymyxins B and E by HPLC (218) 653

Polynuclear aromatic hydrocarbons
see also polycyclic aromatic hydrocarbons

Sephadex LH-20 chromatography of extracts of marine sediment and biological samples for the isolation of polynuclear aromatic hydrocarbons (211) 284

High-performance micro packed flexible columns in LC (213) 25

High-back-pressure LC. I. Development of micro-HPLC using liquefied alkanes as the mobile phase (216) 153

Application of ultra-micro HPLC to trace analysis (218) 199

Polarity of a poly(oxy aryl sulfonyl arylene) used as stationary phase in GLC (in French) (219) 213

Poly(oxy aryl sulfonyl arylene)
Thermal stability of a poly(oxy aryl sulfonyl arylene) used as stationary phase in GLC (in French) (219) 197

Polarity of a poly(oxy aryl sulfonyl arylene) used as stationary phase in GLC (in French) (219) 213

Polypeptides
TLC of closely related polypeptides on silanized silica gel (211) 29

High-speed gel filtration of glycopolypeptides in 6 *M* guanidine hydrochloride (215) 289

HPLC of amino acids, peptides and proteins. XXXVI. Organic solvent modifier effects in the separation of unprotected peptides by reversed-phase LC (218) 497

Improved HPLC method for the analysis of insulins and related compounds (218) 597

Separation of large polypeptides by HPLC (218) 603

HPLC of protein polypeptides on porous silica gel columns (TSK-GEL SW) in the presence of sodium dodecyl sulphate: comparison with SDS-polyacrylamide gel electrophoresis (219) 123

Polyphenyl ethers
Evaluation of substituted polyphenyl ethers as polar phases in GC (217) 39

Polysaccharides
Affinity column for separation of pyruvylated and non-pyruvylated polysaccharides (213) 301

Investigation of the activation of various insoluble polysaccharides with 1,1'-carbonyldiimidazole and of the properties of the activated matrices (219) 361

Polystyrene resins
LC of cephalosporin C and α-amino acid mixtures on polyfunctional polystyrene resins (211) 79

Polyurethane resins
Sorption behaviour of several organic compounds on 6-deoxycyclodextrin polyurethane resins (219) 108

Polyurethanes
Chromatography of polycarbamoyl sulphonates (anionic polyurethane derivatives) (214) 69

Poly(vinyl alcohol)
Some carriers for the immobilization of enzymes based on derivatized poly(vinyl alcohol) and on copolymers of methacrylates with different spacer lengths (215) 13

Porphyrins
Analysis of free stool porphyrins by HPLC (217) 473

Post-column reactors
Fluorescence detection of chloroanilines in LC using a post-column reaction with fluorescamine. Comparison of reactor types and mixing tees (218) 3

Potassium phosphate
Purification of potassium phosphate for HPLC (214) 267

Precision
Experimental evaluation of the precision in time-based high-performance gel permeation chromatography measurements (216) 35

Profiles, chromatographic
Determination of odorous volatiles in air using chromatographic profiles (216) 199

Progesterone
Use of organic modifiers in the mobile phase for the reversed-phase HPLC separation of steroidal hormones (212) 108

Adsorption chromatographic separation of [^{125}I]progesterone-succinyl-tyrosine methyl ester (213) 511

HPLC of progesterone and its metabolites (219) 152

Progestogens
General method for the assay of oral contraceptives by HPLC (216) 269

Programmed flow preparation
Liquid column switching extraction and chromatography for programmed flow preparation (217) 387

Propionyl chloride
GLC analyses of chlorination products of propionyl chloride (213) 63

Prostaglandins
Trimethylsilylation reaction of prostaglandin-E methyl ester with various trimethylsilylating reagents (214) 177

Determination of prostaglandin E_2 in the cricket, *Teleogryllus commodus*, by reversed-phase HPLC (216) 380

Proteases
Resolution in high-performance liquid affinity chromatography. Dependence on eluite diffusion into the stationary phase (216) 169

Proteinases
Hydroxyalkyl methacrylate gels derivatized with epichlorohydrin as supports for large-scale and high-performance affinity chromatography (215) 165

Protein multivalency
Theory of affinity electrophoresis. Evaluation of the effects of protein multivalency, immobilized ligand heterogeneity and microdistribution and determination of effective concentration of immobilized ligand (216) 43

Proteins
see also enzymes

Improved single-column procedure for the amino acid analysis of collagen-like proteins (211) 175

Application of polyacrylamide gradient gel electrophoresis to the study of multiple components of concanavalin A and related lectins (211) 233

Separation and determination of thiamine-binding proteins in rats by HPLC (211) 290

Electrofocusing of stroma-free haemoglobin and its derivatives in agarose isoelectric focusing gels (211) 308

Determination of acetyl and formyl groups by GLC (211) 347

Hydrophobic interaction chromatography: a new method for sunflower protein fractionation (212) 199

Separation of cytochromes *c* by reversed-phase HPLC (212) 295

Characterization of polymerized stroma-free haemoglobin by chromatography, electrophoresis and electrofocusing in gels (213) 183

Physico-chemical and chromatographic properties of new ion exchangers. I. CM-Trisacryl M (213) 389

Analysis of protein and peptide mixtures. Evaluation of three sodium dodecyl sulphate-polyacrylamide gel electrophoresis buffer systems (213) 491

Separation of Dns-amino acids using reversed-phase HPLC: a sensitive method for determining N-termini of peptides and proteins (213) 501

Application of a three-dimensional drawing procedure to the evaluation of series of protein samples after analysis by gel electrophoresis and other methods (214) 126

Isolation of human haemopexin by bioaffinity chromatography on haeme-Sepharose (214) 317

Coupling of proteins and other amines to Sepharose by bromine oxidation and reductive amination (215) 45

Application of carrageenan beads for chromatographic purification of proteins (215) 93

Separation of membrane components by partition in detergent-containing polymer phase systems. Isolation of the light harvesting chlorophyll *a/b* protein (215) 131

Isolation of hormonal proteins and antibodies by affinity chromatography (215) 143

Hydrophobic chromatography on homologous series of alkylagaroses. A comparison of

charged and electrically neutral column materials (215) 211

High-performance liquid affinity chromatography of proteins on Cibacron Blue F3G-A bonded silica (215) 303

HPLC of macromolecules on agarose and its derivatives (215) 317

Adsorption and desorption of proteins in metal chelate affinity chromatography. Purification of albumin (215) 333

Covalent chromatography as a means of isolating thiol peptides from large proteins. Application to human ceruloplasmin (215) 341

Consideration of the nature of the lectin-carbohydrate interaction (215) 351

Affinity chromatography for the purification of lectins (a review) (215) 361

Isoelectric focusing in agarose under denaturating conditions (215) 373

Theory of affinity electrophoresis. Evaluation of the effects of protein multivalency, immobilized ligand heterogeneity and microdistribution and determination of effective concentration of immobilized ligand (216) 43

Resolution in high-performance liquid affinity chromatography. Dependence on eluite diffusion into the stationary phase (216) 169

High-performance liquid affinity chromatography of enzymes on silica-immobilised triazine dyes (216) 175

Evidence for a concentration-dependent polymerization of a commercial human serum albumin (216) 191

Application of ion-exchange chromatography for the production of human albumin (216) 279

Separation of apolipoproteins A-I and A-II by ion-paired reversed-phase HPLC (216) 285

Behaviour of biomolecules in water-organic solvent-inorganic salt two-phase ternary systems (216) 346

Buffer-focusing chromatography using multicomponent electrolyte elution systems (218) 483

Application of 1,1'-carbonyldiimidazole-activated matrices for the purification of proteins. III. The use of 1,1'-carbonyldiimidazole-activated agaroses in the biospecific affinity chromatographic isolation of serum antibodies (218) 509

Assessment and optimization of system parameters in size exclusion separation of proteins on diol-modified silica columns (218) 535

Comparison of short and ultrashort-chain alkylsilane-bonded silicas for the HPLC of proteins by hydrophobic interaction methods (218) 569

Analysis of glycoprotein hormones and other medically important proteins by high-performance gel filtration chromatography (218) 581

HPLC of ^{125}I-labelled proteins with on-line detection (218) 591

HPLC of protein polypeptides on porous silica gel columns (TSK-GEL SW) in the presence of sodium dodecyl sulphate: comparison with SDS-polyacrylamide gel electrophoresis (219) 123

Isoelectric focusing, gel chromatography and electrophoresis of pyridoxalated and polymerized stroma-free haemoglobin (219) 128

Interaction of heme proteins and thyroid hormone. II. Localization of the site on thyroid hormone that binds to hemoglobin (219) 148

Chromatofocusing of pyridoxalated and polymerized human haemoglobin (219) 436

Isoelectric points and molecular weights of proteins. A new table (Review) (220) 115

Protobiochemistry
Copolyamino acid fractionation and protobiochemistry (215) 115

Proton-donating solutes
Separation of proton-donating solutes by LC with a strong proton acceptor, tri-*n*-octylphosphine oxide, in the liquid stationary phase (218) 455

Pseudoephedrine
Determination of cross-contamination of the diastereomers ephedrine and pseudoephedrine by HPLC, TLC and carbon-13 nuclear magnetic resonance spectroscopy (219) 81

Psoralen
TLC assay of photoactive compounds (furocoumarins) using the fungus *Penicillium expansum* as a test organism (214) 349

Purines
LC separation of purines, pyrimidines and their nucleosides on silica gel columns (211) 337

Reversed-phase LCC of pyrimidine and purine derivatives. I. Unbuffered binary aqueous organic mobile phases on octadecylsilica (219) 245

Pyrethroids
HPLC of fenvalerate (216) 303

Pyridoxine
Automation of a complex HPLC system. Procedures and hardware for a vitamin B_6 model system (216) 338

Pyrimidines
LC separation of purines, pyrimidines and their nucleosides on silica gel columns (211) 337

Reversed-phase LCC of pyrimidine and purine derivatives. I. Unbuffered binary aqueous organic mobile phases on octadecylsilica (219) 245

Pyrrolizidine alkaloids
Preparative ion-pair HPLC and GC of pyrrolizidine alkaloids from comfrey (214) 257

Q

Quantitative structure-activity relationships
Chromatography in studies of quantitative structure-activity relationships (220) 71

Quaternary ammonium compounds
Speciation of aryloxyethoxyethyl benzyl dimethyl ammonium salts by glass capillary GC and HPLC (213) 463

R

Radial thermal gradients
Effect of radial thermal gradients in elevated temperature HPLC (218) 123

Recycling
Recycling LC using microbore columns (219) 1

Reflecting supports
Influence of reflecting supports of HPTLC silica gel 60 stationary phase on quantitative *in situ* measurements of thin-layer chromatograms (in German) (213) 453

Refractive index detection
Ion chromatography of inorganic and organic ionic species using refractive index detection (218) 57

Resins
LC of cephalosporin C and α-amino acid mixtures on polyfunctional polystyrene resins (211) 79

Rapid high-resolution separation of oligosaccharides on silver form cation-exchange resins (212) 51

Advantages of a new ion-exchange resin for the LC of biogenic amines: a methacrylic acid polymer cross-linked with butanediol-diacrylate (216) 137

Separation of plant polyphenolics by chromatography on a boronate resin (216) 261

Ligand-exchange chromatography of amino acid racemates on Separon gels containing L-proline or L-hydroxyproline groupings (217) 539

Sorption behaviour of several organic compounds on 6-deoxycyclodextrin polyurethane resins (219) 108

Resolution
Factors affecting retention and resolution of peptides in HPLC (211) 15

Resolution in high-performance liquid affinity chromatography. Dependence on eluite diffusion into the stationary phase (216) 169

High-resolution separations based on electrophoresis and electroosmosis (218) 209

Comparison of sedimentation field flow fractionation with chromatographic methods for particulate and high-molecular-weight macromolecular characterizations (218) 217

Retention behaviour
LC with crown ether-containing mobile phases. I. Capacity factors of amino compounds on a hydrophobic stationary phase (211) 1

Factors affecting retention and resolution of peptides in HPLC (211) 15

LC separation of purines, pyrimidines and their nucleosides on silica gel columns (211) 337

Chromatographic behaviour of aromatic compounds on non-polar phases with adsorbed quaternary ammonium salts (212) 1

Preparation and evaluation of N-(2-aminoethyl)-γ-aminopropyltrimethoxysilane-treated silica columns for the HPLC analysis of some aromatic amines (212) 251

Prediction of retention behaviour of some phenolic antioxidants (213) 363

GC data for chlorobenzenes, chloroanilines and N-sulphinylanilines. Relationship between molecular structure and retention based on retention index increments calculated by computer (214) 47

Study of retention behaviour of primary, secondary and tertiary anilines in normal- and reversed-phase LC (214) 291

GLC retention behaviour of hindered aliphatic esters (217) 125

Plasma catecholamines in hypertension and pheochromocytoma determined using ion-pair reversed-phase chromatography with amperometric detection. Investigation of the separation mechanism and clinical methodology (217) 523

Systematic study of ternary solvent behaviour in reversed-phase LC (218) 261

HPLC of organic acids on bare silica (218) 285

HPLC of amino acids, peptides and proteins. XXXVI. Organic solvent modifier effects in the separation of unprotected peptides by reversed-phase LC (218) 497

Influence of intramolecular interactions on the GC separation of epimers of dicyclopentadiene derivatives (in German) (219) 113

Studies on the GC behaviour of organophosphorus esters (219) 297

Contributions to the chemistry of silicon-sulphur compounds. XXVI. Relationships between chromatographic behaviour and structure of triorganosilanes, triorganosilanethiols, hexaorganodisilthianes and triphenylthioalkylsilanes (219) 302

Retention characteristics
Retention and selectivity characteristics of a non-polar perfluorinated stationary phase for LC (218) 443

Retention data
The chemistry of carbazoles. IX. Substituent effect in the GLC of methylcarbazoles (211) 388

Measurement of retention data on open tubular columns coupled in series (213) 189

Behaviour of stationary phases. I. Linear relationships between retention data of benzene derivatives (in French) (213) 223

Group-contribution approach to the behaviour of 2-phenylethylamines in reversed-phase HPLC (218) 639

HPLC of alkylbenzenes on silica (219) 53

Retention indices
Conversion of linear retention indices into logarithmic retention indices (211) 239

Prediction of retention indices of aromatic hydrocarbons on phases of different polarity (216) 321

Relationship between GC retention indices of homologous compounds and the number of carbon atoms and temperature (216) 326

GC of homologous esters. XV. Molecular retention indices of aliphatic esters (219) 45

GLC retention indices of 1318 substances of toxicological interest on SE-30 or OV-1 stationary phase (Review) (220) 195

Retentionless tubes
The height equivalent to a theoretical plate of retentionless rectangular tubes (216) 1

Retention mechanism
The role of hydrogen bonds in chromatography. I. Amino- and hydroxyanthraquinones (212) 139

Hydrophobic chromatography with dynamically coated stationary phases. V. "Duplex soap chromatography" (212) 271

Hydrophobic chromatography on homologous series of alkylagaroses. A comparison of charged and electrically neutral column materials (215) 211

Correlation of Hammet substituent constants with reversed-phase "ion-pair" LC adsorption parameters of charged organic compounds (217) 421

Development of modern bioaffinity chromatography (a review) (218) 241

Study on the retention of amines in reversed-phase ion-pair chromatography on bonded phases (218) 327

Mechanism of zwitterion-pair chromatography. I. Nucleotides (218) 341

Mechanism of zwitterion-pair chromatography. II. Ampicilline, lysergic acid, tryptophan and other solutes (218) 355

High-performance displacement chromatography (218) 365

Chromatographic characterization of silica surfaces (218) 395

Structural effects in enthalpy/entropy compensated and non-compensated behaviour in ion-pair reversed-phase HPLSC (218) 427

Assessment and optimization of system parameters in size exclusion separation of proteins on diol-modified silica columns (218) 535

Chromatographic study of optical resolution. VIII. Theoretical study of the chromatographic behaviour of the enantiomers of racemic complex cations on a cation-exchange column (219) 29

Retention parameters
Estimation of aqueous solubilities of organic non-electrolytes using LC retention data (218) 409

Ribosylzeatin
Combined HPLC radioimmunoassay for cytokinins (214) 101

R_M values
R_M values of steroids as an expression of their hydrophobic character. II. Effect of neighbouring groups on the hydrophobicity of substituents (216) 63

R_M values of steroids as an expression of their hydrophobic character. III. Interpretation of the slopes of the R_M *versus* acetone concentration curves (216) 69

R_M and log P values of 5-nitroimidazoles (216) 93

Rodenticides
Determination of chlorophacinone in formulations by reversed-phase ion-pair chromatography (214) 335

S

Saccharides
see also sugars and carbohydrates

Rapid high-resolution separation of oligosaccharides on silver form cation-exchange resins (212) 51

Separation of reduced disaccharides derived from glycosaminoglycans by HPLC (212) 65

HPLC of maltosaccharides (212) 126

New chromatographic system for the rapid analysis and preparation of colostrum sialyloligosaccharides (212) 313

Separation of monosaccharides as trimethylsilylated alditols on fused-silica capillary columns (213) 146

Rapid separation of malto-, xylo- and cellooligosaccharides (DP 2-9) on cation-exchange resin using water as eluent (213) 151

Affinity column for separation of pyruvylated and non-pyruvylated polysaccharides (213) 301

Separation and determination of phenyl isocyanate-derivatized carbohydrates and sugar alcohols by HPLC with UV detection (218) 65

Salicylic acid
Separation of acidic drugs in the µg/ml range in untreated blood plasma by direct injection on LC columns (218) 671

Sample transfer
Sample transfer in splitless injections in capillary GC (214) 118

Sampling techniques
Coupling of a low-pressure sampling technique to a gas chromatography (in French) (211) 247

Saponins
Identification of ginsenosides from *Panax ginseng* in fractions obtained by HPLC by field desorption MS, multiple internal reflection IR spectroscopy and TLC (212) 37

Sedative drugs
Comparison of C_n bonded silica gel TLC plates: conditions for use and separations of some barbiturates (214) 209

Analysis of barbiturates by GC (Review) (220) 253

Sedimentation field flow fractionation
Comparison of sedimentation field flow fractionation with chromatographic methods for particulate and high-molecular-weight macromolecular characterizations (218) 217

Selectivity
Retention and selectivity characteristics of a non-polar perfluorinated stationary phase for LC (218) 443

Separation efficiency
Microcapillary LC in open tubular columns with diameters of 10-50 µm. Potential application to chemical ionization MS detection (218) 137

High-resolution separations based on electrophoresis and electroosmosis (218) 209

Serotonin
HPLC determination of biogenic amines. II. Comparison of detection methods (213) 429

Determination of 5-hydroxytryptophan, serotonin and 5-hydroxyindoleacetic acid in rat and human brain and biological fluids by reversed-phase HPLC with electrochemical detection (218) 663

Sesquiterpene lactones
Separation and identification of stereoisomers of sesquiterpene lactones by multiple development of thin-layer chromatograms (212) 379

Shallot volatile oil
HPLC separation of shallot volatile oil (214) 234

Shogaols
GC and MS studies of ginger constituents. Identification of gingerdiones and new hexahydrocurcumin analogues (212) 75

Short open tubes
Dispersion of peaks by short straight open tubes in LC systems (218) 97

Sialic acids
Sialic acid quantitation by analytical isotachophoresis (214) 156

Sialyloligosaccharides
New chromatographic system for the rapid analysis and preparation of colostrum sialyloligosaccharides (212) 313

Silica
Determination of hydroxyl groups and water content in silica by nuclear magnetic resonance spectroscopy (213) 33

GC behaviour of chemically bonded microparticulate silica (217) 191

Chromatographic characterization of silica surfaces (218) 395

Effect of pH on the GC behaviour of silica gel (219) 101

Silica gel
HPLC of organic acids on bare silica (218) 285

Silicon-containing compounds
Response enhancement by metal doping of a silicon-selective flame ionization detector for GC (213) 397

Silicon-sulphur compounds
Contributions to the chemistry of silicon-sulphur compounds. XXVI. Relationships between chromatographic behaviour and structure of triorganosilanes, triorganosilanethiols, hexaorganodisilthianes and triphenylthioalkylsilanes (219) 302

Snake venoms
Single-step fractionation of *Vipera russelli* venom. A sensitive fluorimetric method to study the elution profile (216) 395

Sodium dodecyl sulphate
Determination of carbon chain distribution in alkyl sulfates by *in situ* hydrolysis-GC (219) 117

Sodium nitroprusside
HPLC determination of sodium nitroprusside (212) 339

Solanidines
Analysis of potato glycoalkaloids with radially compressed HPLC cartridges and ethanolamine in the mobile phase (219) 403

Solubilities of gases
Solubilities of gases in H_2O and ^2H_2O (216) 161

Solubility
Estimation of aqueous solubilities of organic non-electrolytes using LC retention data (218) 409

Solute diffusion
Resolution in high-performance liquid affinity chromatography. Dependence on eluite diffusion into the stationary phase (216) 169

Solvent localization
Solvent strength of multicomponent mobile phases in LSC. Binary-solvent mixtures and solvent localization (214) 1

Solvent programmer
Computer-controlled single-pump solvent programmer for HPLC (218) 83

Solvent recovery
Recovery of HPLC-grade acetonitrile by spinning-band distillation (213) 331

Solvent-solute localization effects
Theoretical basis for systematic optimization of mobile phase selectivity in LSC. Solvent-solute localization effects (218) 299

Solvent strength
Solvent strength of multicomponent mobile phases in LSC. Binary-solvent mixtures and solvent localization (214) 1

Solvent strength of multicomponent mobile phases in LSC. Mixtures of three or more solvents (214) 21

Solvent systems
Behaviour of biomolecules in water-organic solvent-inorganic salt two-phase ternary systems (216) 346

Sorbents
see also adsorbents

Hydroxyalkyl methacrylate gels derivatized with epichlorohydrin as supports for large-scale and high-performance affinity chromatography (215) 165

Comparison of the distribution constants of benzene in different chromatographic sorbent-gas systems, determined by direct measurement of sorption equilibria and calculated from GC retention data (217) 167

Applications of selective complexation by a europium(III) coordination polymer sorbent for the pre-fractionation of volatile compounds (217) 275

Ligand-exchange chromatography of amino acid racemates on Separon gels containing L-proline or L-hydroxyproline groupings (217) 539

Comparison of the distribution constants of acetone in different chromatographic sorbent-gas systems, determined by direct measurement of sorption equilibria and calculated from GC retention data (219) 37

Sorption behaviour of several organic compounds on 6-deoxycyclodextrin polyurethane resins (219) 108

Sorption traps
Direct desorption of traps for capillary column GC (214) 197

Spasmolytic drugs
Separation, identification and quantitation of nitroglycerin and its metabolic or hydrolysis products (216) 239

Spectrophotometric detectors
Use of 214-nm and 229-nm discrete-line sources for the UV absorbance detection of some pesticides separated by HPLC (211) 144

Split-splitless injector
Application of the split-splitless injector to environmental analysis (217) 91

Spray reagents
Distinctive test with copper(II)-ninhydrin reagent for small α-peptides separated PC (213) 307

Sensitive, enzyme-catalyzed, chromogenic reagent for hydrogen peroxide and other peroxygen compounds on TLC plates (216) 413

Stabilizers
Antioxidants and stabilizers. LXXXVI. Use of chromatography in the investigation of the mechanism of action of amine antidegradants (211) 299

Analysis of 2-hydroxybenzophenone and 2'-hydroxyphenylbenzotriazole UV stabilizers by HPLC (214) 237

Analysis of additives in plastics by high-performance size-exclusion chromatography (218) 703

Stationary phases
LC of cephalosporin C and α-amino acid mixtures on polyfunctional polystyrene resins (211) 79

GC behaviour of mono- and dihydroxybiphenyls on various silicone phases (211) 150

Capillary columns with immobilized stationary phases. I. A new simple preparation procedure (211) 243

An efficient chemically bonded reversed-phase thin-layer plate permitting the use of water-rich mobile phases (211) 262

GC separation of some enantiomers on optically active copper(II) complexes (211) 274

Chromatographic behaviour of aromatic compounds on non-polar phases with adsorbed quaternary ammonium salts (212) 1

Coordination polymers as adsorbents and stationary phases in GC. VII. Alkaline earth metal di-*n*-hexylphosphinates: specific interactions (212) 29

Preparation and evaluation of N-(2-aminoethyl)-γ-aminopropyltrimethoxysilane-treated silica columns for the HPLC analysis of some aromatic amines (212) 251

N-(1*R*, 3*R*)-*trans*-Chrysanthemoyl (*R*)-1-(α-naphthyl)ethylamine as a stationary phase for the separation of optical isomers by GC (213) 137

Capillary columns with immobilized stationary phases. II. Practical advantages and details of procedures (213) 211

Behaviour of stationary phases. I. Linear relationships between retention data of benzene derivatives (in French) (213) 223

Practical system for polarity rating of packed GLC columns (213) 317

Evaluation of liquid crystal smectic mesophases for GLC separations (214) 161

Influence of the LC mobile phase on the phase transitions of alkyl-bonded silicas studied by GC (214) 202

Comparison of C_n bonded silica gel TLC plates: conditions for use and separations of some barbiturates (214) 209

Evaluation of substituted polyphenyl ethers as polar phases in GC (217) 39

New results in the GC separation of enantiomers of hydroxy acids and carbohydrates (217) 71

Scanning electron microscopy as an aid in GC. Study of porous polymers coated with stationary phases (217) 109

New criterion for polarity of stationary phases in GC (217) 139

GC behaviour of chemically bonded microparticulate silica (217) 191

Stationary phase characterization in HPLC. A test for trace metal activity in octadecyl bonded silica gel (217) 399

Chromatographic characterization of silica surfaces (218) 395

Retention and selectivity characteristics of a non-polar perfluorinated stationary phase for LC (218) 443

Separation of proton-donating solutes by LC with a strong proton acceptor, tri-*n*-octylphosphine oxide, in the liquid stationary phase (218) 455

Selectivity and efficiency of separation of isomers of organic acids by clathrate chromatography (218) 465

Comparison of short and ultrashort-chain alkylsilane-bonded silicas for the HPLC of proteins by hydrophobic interaction methods (218) 569

Thermal stability of a poly(oxy aryl sulfonyl arylene) used as stationary phase in GLC (in French) (219) 197

Polarity of a poly(oxy aryl sulfonyl arylene) used as stationary phase in GLC (in French) (219) 213

High-resolution GC with liquid crystal glass capillaries. III. Separation of hydrocarbon isomers on cholesteric mesophase (in German) (219) 225

LC of sugars on silica-based stationary phases (Review) (220) 313

Steroids
Use of organic modifiers in the mobile phase for the reversed-phase HPLC separation of steroidal hormones (212) 108

LC measurement of cortisol in methylene chloride extracts of aqueous solutions (212) 221

HPLC of naturally occurring estrogens (212) 239

Adsorption chromatographic separation of [^{125}I]progesterone-succinyl-tyrosine methyl ester (213) 511

R_M values of steroids as an expression of their hydrophobic character. II. Effect of neighbouring groups on the hydrophobicity of substituents (216) 63

R_M values of steroids as an expression of their hydrophobic character. III. Interpretation of the slopes of the R_M *versus* acetone concentration curves (216) 69

General method for the assay of oral contraceptives by HPLC (216) 269

HPTLC determination of fluorescence-labelled cortisol (217) 349

Analysis of isomeric ethynylestradiol glucuronides in urine (217) 453

HPLC of progesterone and its metabolites (219) 152

Sterols
Capillary column GLC analysis of cholesterol derivatives. Application to the autoxidation products of cholesterol (211) 392

Stimulant drugs
Sensitive method for the HPLC determination of amphetamine in urinary extracts (213) 166

Streptomycin
Determination of streptomycin sulfate and dihydrostreptomycin sulfate by HPLC (219) 89

Structure-activity relationships
R_M and log P values of 5-nitroimidazoles (216) 93

Structure-retention relationships
GC data for chlorobenzenes, chloroanilines and N-sulphinylanilines. Relationship between molecular structure and retention based on retention index increments calculated by computer (214) 47

GC of homologous esters. XV. Molecular retention indices of aliphatic esters (219) 45

Substances of toxicological interest
GLC retention indices of 1318 substances of toxicological interest on SE-30 or OV-1 stationary phase (Review) (220) 195

Substituent constants
Correlation of Hammet substituent constants with reversed-phase "ion-pair" LC adsorption parameters of charged organic compounds (217) 421

Sugar alcohols
Separation and determination of phenyl isocyanate-derivatized carbohydrates and sugar alcohols by HPLC with UV detection (218) 65

Sugars
see also saccharides and carbohydrates

Continuous TLC of sugars of clinical interest in samples of urine impregnated on paper (217) 357

LC of sugars on silica-based stationary phases (Review) (220) 313

Sulphide
Methylation of inorganic anions for GC determination (211) 361

Electron-capture GC determination of sulphide as a new pentafluorobenzyl derivative (219) 312

Sulphinylanilines
GC data for chlorobenzenes, chloroanilines and N-sulphinylanilines. Relationship between molecular structure and retention based on retention index increments calculated by computer (214) 47

Sulphonic acids
HPLC of organic acids on bare silica (218) 285

Sulphur compounds
LC with UV absorbance and polarographic detection of ethylenethiourea and related sulfur compounds. Application to rat urine analysis (212) 245

Sulphur imides
Separation of cyclic sulphur-nitrogen compounds by HPLC. LXXVIII (216) 399

Sulphur mustards
TLC of some sulfur and nitrogen mustards (213) 173

Supports
Determination of hydroxyl groups and water content in silica by nuclear magnetic resonance spectroscopy (213) 33

Influence of reflecting supports of HPTLC silica gel 60 stationary phase on quantitative *in situ* measurements of thin-layer chromatograms (in German) (213) 453

Polymeric supports bearing isonitrile functional groups for covalent fixation of biologically active molecules (a review) (215) 31

Coupling of proteins and other amines to Sepharose by bromine oxidation and reductive amination (215) 45

Hydroxyalkyl methacrylate gels derivatized with epichlorohydrin as supports for large-scale and high-performance affinity chromatography (215) 165

Scanning electron microscopy as an aid in GC. Study of porous polymers coated with stationary phases (217) 109

GC behaviour of chemically bonded microparticulate silica (217) 191

Investigation of the activation of cross-linked agarose with carbonylating reagents and the preparation of matrices for affinity chromatography purifications (219) 353

Investigation of the activation of various insoluble polysaccharides with 1,1'-carbonyldiimidazole and of the properties of the activated matrices (219) 361

Surface silanols
Determination of hydroxyl groups and water content in silica by nuclear magnetic resonance spectroscopy (213) 33

Role of surface groups in affecting the chromatographic performance of certain types of fused-silica glass capillary columns. I (217) 3

Surfactants
Hydrophobic chromatography with dynamically coated stationary phases. V. "Duplex soap chromatography" (212) 271

Sympathomimetic drugs
Group-contribution approach to the behaviour of 2-phenylethylamines in reversed-phase HPLC (218) 639

T

Testosterone
Use of organic modifiers in the mobile phase for the reversed-phase HPLC separation of steroidal hormones (212) 108

HPLC of progesterone and its metabolites (219) 152

Tetrachlorophenols
Determination of pentachlorophenol and related compounds in animal materials by HPLC and GC (216) 229

Tetrahydrofurans
Manganese(II)-bis(3-heptafluorobutyryl-1*R*-camphorate): a versatile agent for the resolution of racemic cyclic ethers by complexation GC (217) 51

Thermal gradients
Effect of radial thermal gradients in elevated temperature HPLC (218) 123

Thermal lens calorimetry
Thermal lens calorimetry. Application to chromatographic detection (218) 15

Thermionic detector
Micro-column HPLC and flame-based detection principles (218) 179

Thermostats
Effect of mobile phase pre-heating on HPLC column performance: a new type of column thermostat (213) 243

Thiamine-binding proteins
Separation and determination of thiamine-binding proteins in rats by HPLC (211) 290

Thin-layer plates
An efficient chemically bonded reversed-phase thin-layer plate permitting the use of water-rich mobile phases (211) 262

Influence of reflecting supports of HPTLC silica gel 60 stationary phase on quantitative *in situ* measurements of thin-layer chromatograms (in German) (213) 453

Thiocyanate
Methylation of inorganic anions for GC determination (211) 361

Thioimidazole
LC with UV absorbance and polarographic detection of ethylenethiourea and related sulfur compounds. Application to rat urine analysis (212) 245

Thiol peptides
Covalent chromatography as a means of isolating thiol peptides from large proteins. Application to human ceruloplasmin (215) 341

Thiols
Photometric monitoring of thiols by means of a thiomercurimetric detector (211) 252

Thiomercurimetric detector
Photometric monitoring of thiols by means of a thiomercurimetric detector (211) 252

Thymol
HPLC analysis of carvacrol and thymol in the essential oil of *Thymus capitatus* (219) 332

Thyroid hormones
Interaction of heme proteins and thyroid hormone. II. Localization of the site on thyroid hormone that binds to hemoglobin (219) 148

Time-based GPC
Experimental evaluation of the precision in time-based high-performance gel permeation chromatography measurements (216) 35

Tobacco
Rapid method for the analysis of tobacco nicotine alkaloids (211) 111

Toluene diisocyanate
GC determination of free toluene diisocyanate in flexible urethane foams (213) 162

Toxicological interesting substances
GLC retention indices of 1318 substances of toxicological interest on SE-30 or OV-1 stationary phase (Review) (220) 195

Toyopearl columns
Packing of Toyopearl columns for gel filtration. IV. Gravitational packing and influence of slurry reservoir size (211) 383

Trace analysis
Determination of trace atmospheric isocyanate concentrations by reversed-phase HPLC using 1-(2-pyridyl)piperazine reagent (212) 93

Fractionation of polar organic constituents in environmental samples using the lipophilic dextran gels Sephadex LH-20 and Sephasorb HP Ultrafine. Application to a weathered Ekofisk crude oil (214) 83

Comparison of column packings for trace analysis of free amines by GLC (217) 177

Determination of C_2-C_5 hydrocarbons in the atmosphere at low parts per 10^9 to high parts per 10^{12} levels (217) 301

Determination of trace amounts of hydrogen halides in the atmosphere by collection in sampling tubes and subsequent derivatization and GC (217) 311

Application of ultra-micro HPLC to trace analysis (218) 199

Comparison of the distribution constants of acetone in different chromatographic sorbent-gas systems, determined by direct measurement of sorption equilibria and calculated from GC retention data (219) 37

Tranquilizers
Analysis of xylazine in biological material by GC using packed and capillary columns (216) 350

LC of demoxepam and phenothiazines using a post-column photochemical reactor and fluorescence detection (217) 463

Transformer oils
Determination of 2,6-di-(*tert.* -butyl)-4-methylphenol in new and aged transformer oils using HPLC (in French) (213) 481

Trialkyltin compounds
Simultaneous determination of trialkyltin homologues in biological materials (216) 209

Triazine dyes
High-performance liquid affinity chromatography of enzymes on silica-immobilised triazine dyes (216) 175

Triazines
GLC of *s*-triazines on a surface-bonded support in short columns (212) 85

Tricarboxylic acids
PC separation of phosphate esters, tricarboxylic cycle acids and amino acids in extracts from malaria parasites (214) 111

Isotachophoretic analysis of organic acids produced by fermentation from *n*-alkanes (216) 313

Trichothecenes
HPLC of trichothecenes. I. Detection of T-2 toxin and HT-2 toxin (212) 370

Trimetaphosphate
Study of the acidic hydrolysis of cyclic trimetaphosphate by LC (219) 385

Trimethylolpropane
HPLC separation of the intermediate products in the synthesis of trimethylolpropane (216) 402

Trimethylphosphorodithioate
The determination of O,S,S-trimethylphosphorodithioate in the plasma and various tissues of rats using high-resolution GC with nitrogen-phosphorus detection (219) 285

Trimethylsilylation
Trimethylsilylation reaction of prostaglandin-E methyl ester with various trimethylsilylating reagents (214) 177

Triorganosilanes
Contributions to the chemistry of silicon-sulphur compounds. XXVI. Relationships between chromatographic behaviour and structure of triorganosilanes, triorganosilanethiols, hexaorganodisilthianes and triphenylthioalkylsilanes (219) 302

Trioxalen
TLC assay of photoactive compounds (furocoumarins) using the fungus *Penicillium expansum* as a test organism (214) 349

Tripdiolide
Rapid TLC assay of tripdiolide using fluorimetric detection (214) 152

Tryptic peptides
Separation of the tryptic peptides from reduced, alkylated hen egg white lysozyme by HPLC (214) 185

Tryptophan
Mechanism of zwitterion-pair chromatography. II. Ampicilline, lysergic acid, tryptophan and other solutes (218) 355

Tswett
New data on M.S. Tswett's life and work (Review) (220) 1

Tubes
Dispersion of peaks by short straight open tubes in LC systems (218) 97

T-2 toxin
HPLC of trichothecenes. I. Detection of T-2 toxin and HT-2 toxin (212) 370

T4D virus
Characterization of T4D virus by sedimentation field-flow fractionation (215) 323

U

Ultraviolet detector
Extra-column effects in polarographic *versus* UV detection in HPLC (218) 45

Urobilinoids
HPLC of bile pigments: separation and characterization of the urobilinoids (218) 647

Urushiol
Separation of urushiol by HPLC on an 8% octa,decylsilane chemically bonded silica gel column with electrochemical detection. Analysis of urushiol in the sap of lac trees (*Rhus vernicifera*) and that in the Japanese lac-making process (214) 343

UV-absorbing ions
Reversed-phase ion-pair chromatography with UV-absorbing ions in the mobile phase (218) 31

V

Vanillate hydroxylase
Purification and partial characterization of vanillate hydroxylase (decarboxylating) from *Sporotrichum pulverulentum* (215) 99

Vanillin
Assay for catechol-O-methyltransferase by HPLC with fluorescence detection (211) 129

Vaporizer
Automatic injection in high-resolution GC: a programmed temperature vaporizer as a general purpose injection system (217) 81

Vasodilator drugs
Separation, identification and quantitation of nitroglycerin and its metabolic or hydrolysis products (216) 239

Venoms
Single-step fractionation of *Vipera russelli* venom. A sensitive fluorimetric method to study the elution profile (216) 395

Verapamil
Rapid HPLC method for the measurement of verapamil and norverapamil in blood plasma or serum (218) 621

Very-high-speed LC
Very-high-speed LC. II. Some instrumental factors influencing performance (217) 369

Vinca alkaloids
HPLC of *Vinca rosea* alkaloids and the correlation of plate height and molecular weight (214) 95

Viruses
Application of a general equation for controlled pore glass permeation chromatography to an aggregating, spherical virus (211) 53

Characterization of T4D virus by sedimentation field-flow fractionation (215) 323

Viscosity phenomena
Concentration effects in gel permeation chromatography. VII. Viscosity phenomena and column geometry (211) 39

Vitamin B_6
Automation of a complex HPLC system. Procedures and hardware for a vitamin B_6 model system (216) 338

Vitamin C
Separation of ascorbic and isoascorbic acids by ion-pair reversed-phase chromatography (in French) (219) 338

Vitamins
Resolution of fat-soluble vitamins in HPLC with methanol-containing mobile phases (211) 155

Volatile compounds
Applications of selective complexation by a europium(III) coordination polymer sorbent for the pre-fractionation of volatile compounds (217) 275

Voltammetric detectors
Carbon pastes for voltammetric detectors in HPLC (213) 41

Voltaren
Simultaneous determination of diclofenac sodium and its hydroxy metabolites by capillary column GC with electron-capture detection (217) 263

W

Water
GC determination of water in acetone (212) 121

Determination of hydroxyl groups and water content in silica by nuclear magnetic resonance spectroscopy (213) 33

Solubilities of gases in H_2O and ^2H_2O (216) 161

X

Xanthinin
Separation and identification of stereoisomers of sesquiterpene lactones by multiple development of thin-layer chromatograms (212) 379

Xanthotoxin
TLC assay of photoactive compounds (furocoumarins) using the fungus *Penicillium expansum* as a test organism (214) 349

Xylazine
Analysis of xylazine in biological material by GC using packed and capillary columns (216) 350

Y

Ylides
Applications of HPLC in organometallic chemistry: ligand exchange in complexes of ylides (in German) (213) 47

Z

Zimelidine
Straight-phase ion-pair chromatography of zimelidine and similar divalent amines. II. The chromatographic system (211) 181

Zirconium derivatives
Use of metal salts as fluorescence-inducing reagents in TLC (217) 329

Zwitterionic solutes
Mechanism of zwitterion-pair chromatography. I. Nucleotides (218) 341

Mechanism of zwitterion-pair chromatography. II. Ampicilline, lysergic acid, tryptophan and other solutes (218) 355

PUBLICATION SCHEDULE FOR 1982

Journal of Chromatography (incorporating *Chromatographic Reviews*) and *Journal of Chromatography, Biomedical Applications*

MONTH	J	F	M	A	M	J	J	A	S	O	N	D
Journal of Chromatography	234/1 234/2 235/1 235/2	236/1 236/2	237/1 237/2 237/3	238/1 238/2 239	240/1 240/2 241/1	The publication schedule for further issues will be published later.						
Chromatographic Reviews		251/1		251/2								
Biomedical Applications	227/1	227/2	228	229/1	229/2							

INFORMATION FOR AUTHORS

(Detailed *Instructions to Authors* were published in Vol. 209, No. 3, pp. 501-504. A free reprint can be obtained by application to the publisher.)

Types of Contributions. The following types of papers are published in the *Journal of Chromatography* and the section on *Biomedical Applications:* Regular research papers (Full-length papers), Short communications and Notes. Short communications are preliminary announcements of important new developments and will, whenever possible, be published with maximum speed. Notes are usually descriptions of short investigations and reflect the same quality of research as Full-length papers, but should preferably not exceed four printed pages. For review articles, see page 2 of cover under Submission of Papers.

Submission. Every paper must be accompanied by a letter from the senior author, stating that he is submitting the paper for publication in the *Journal of Chromatography*. Please do not send a letter signed by the director of the institute or the professor unless he is one of the authors.

Manuscripts. Manuscripts should be typed in double spacing on consecutively numbered pages of uniform size. The manuscript should be preceded by a sheet of manuscript paper carrying the title of the paper and the name and full postal address of the person to whom the proofs are to be sent. Authors of papers in French or German are requested to supply an English translation of the title of the paper. As a rule, papers should be divided into sections, headed by a caption (*e.g.,* Summary, Introduction, Experimental, Results, Discussion, etc.). All illustrations, photographs, tables, etc., should be on separate sheets.

Introduction. Every paper must have a concise introduction mentioning what has been done before on the topic described, and stating clearly what is new in the paper now submitted.

Summary. Full-length papers and Review articles should have a summary of 50–100 words which clearly and briefly indicates what is new, different and significant. In the case of French or German articles an additional summary in English, headed by an English translation of the title, should also be provided. (Short communications and Notes are published without a summary.)

Illustrations. The figures should be submitted in a form suitable for reproduction, drawn in Indian ink on drawing or tracing paper. Each illustration should have a legend, all the *legends* being typed (with double spacing) together on a *separate sheet*. If structures are given in the text, the original drawings should be supplied. Coloured illustrations are reproduced at the author's expense, the cost being determined by the number of pages and by the number of colours needed. The written permission of the author and publisher must be obtained for the use of any figure already published. Its source must be indicated in the legend.

References. References should be numbered in the order in which they are cited in the text, and listed in numerical sequence on a separate sheet at the end of the article. Please check a recent issue for the lay-out of the reference list. Abbreviations for the titles of journals should follow the system used by *Chemical Abstracts*. Articles not yet published should be given as "in press", "submitted for publication", "in preparation" or "personal communication".

Proofs. One set of proofs will be sent to the author to be carefully checked for printer's errors. Corrections must be restricted to instances in which the proof is at variance with the manuscript. "Extra corrections" will be inserted at the author's expense.

Reprints. Fifty reprints of Full-length papers, Short communications and Notes will be supplied free of charge. Additional reprints can be ordered by the authors. An order form containing price quotations will be sent to the authors together with the proofs of their article.

News. News releases of new products and developments, and information leaflets of meetings should be addressed to: The Editor of the News Section, Journal of Chromatography/Journal of Chromatography, Biomedical Applications, Elsevier Scientific Publishing Company, P.O. Box 330, 1000 AH Amsterdam, The Netherlands.

Advertisements. Advertisement rates are available from the publisher on request. The Editors of the journal accept no responsibility for the contents of the advertisements.

Modern Thin-Layer Chromatography

can only be fully exploited with a complete system of instruments for sample application, chromatographic development and evaluation.

Here is a complete system for conventional TLC

1. Sample Application





or

CAMAG Micro-Applicator

variable volume sample application 0.5–2.3 µl (spotwise), reproducibility 1%.

CAMAG LINOMAT III*

application of larger sample volumes (approx. 2–500 µl), in the form of narrow bands, reproducibility 1%.

2. Chromatogram Development



CAMAG
Twin-Trough Chamber

versatile,
economical in solvent consumption,
reproducible conditions

3. Quantitative Evaluation



CAMAG TLC/HPTLC Scanner*

Available in two options: low-priced filter version or monochromator equipped.

Scanning modes: absorbance, fluorescence, fluorescence quenching.

Automatic scanning of all tracks of a complete chromatogram, or the selected segment.

Ask for full information on COMPLETE CAMAG SYSTEMS for High-Performance TLC also.

The units marked with an * are, by the way, compatible with HPTLC anyway.

If you are planning to set up a new TLC laboratory, or to complement your existing TLC equipment, ask the CAMAG specialist. He will help.

CAMAG Sonnenmattstr. 11 · CH-4132 Muttenz/Switzerland · Tel. (061) 61 34 34 · Telex 62 649